

1996

Invariant Subspaces of the Harmonic Dirichlet Space with Large Co-Dimension

William T. Ross

University of Richmond, wross@richmond.eduFollow this and additional works at: <http://scholarship.richmond.edu/mathcs-faculty-publications>Part of the [Algebra Commons](#)

Recommended Citation

Ross, William T. "Invariant Subspaces of the Harmonic Dirichlet Space with Large Co-Dimension." *Proceedings of the American Mathematical Society* 124, no. 6 (1996): 1841-1846.

This Article is brought to you for free and open access by the Math and Computer Science at UR Scholarship Repository. It has been accepted for inclusion in Math and Computer Science Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact scholarshiprepository@richmond.edu.

**INVARIANT SUBSPACES OF THE HARMONIC DIRICHLET
 SPACE WITH LARGE CO-DIMENSION**

WILLIAM T. ROSS

(Communicated by Palle E. T. Jorgensen)

Abstract. In this paper, we comment on the complexity of the invariant subspaces (under the bilateral Dirichlet shift $f \rightarrow \zeta f$) of the harmonic Dirichlet space D . Using the sampling theory of Seip and some work on invariant subspaces of Bergman spaces, we will give examples of invariant subspaces $F \subset D$ with $\dim(F/\zeta F) = n$, $n \in \mathbf{N} \cup \{\infty\}$. We will also generalize this to the Dirichlet classes D_a , $0 < a < \infty$, as well as the Besov classes $B_{p,\theta}^a$, $1 < p < \infty$, $0 < a < 1$.

1. Introduction

For a space X of functions on the unit circle $\mathbf{T} = \{z \in \mathbf{C} : |z| = 1\}$ for which the shift operator

$$T : X \rightarrow X, \quad (Tf)(\zeta) = \zeta f(\zeta)$$

is continuous and invertible (e.g. $L^p(\mathbf{T}, |d\zeta|)$, $C^n(\mathbf{T})$, $C^\infty(\mathbf{T})$, $W_n^p(\mathbf{T})$ the Sobolev classes, $B_p^a(\mathbf{T})$ the Besov classes, D_a the Dirichlet classes, $BL_{p,\theta}^n(\mathbf{T})$ the Triebel-Lizorkin classes), the problem of characterizing the subspaces (closed linear manifolds) $F \subset X$ for which $TF \subset F$, the so called *invariant subspaces*, is a very difficult and open problem. There are two types of invariant subspaces to consider: *simply invariant* (or 1-invariant) $TF \subsetneq F$, and *2-invariant* $TF = F$. The 2-invariant subspaces are often described by their zero sets in \mathbf{T} [1], [10], [11], [12], while the 1-invariant subspaces are known to be much more complicated and for most of the classes mentioned above, a complete characterization seems a long way off [3], [6], [8], [9].

In this paper, we focus our attention on the *harmonic Dirichlet space* D of functions $f \in L^2(\mathbf{T}, |d\zeta|)$ with finite norm

$$\|f\|^2 = \sum_{n=-\infty}^{\infty} (1 + |n|) |\hat{f}(n)|^2,$$

where $\{\hat{f}(n)\}$ are the Fourier coefficients of f . The 2-invariant subspaces of D can be characterized by their zero sets on \mathbf{T} [12], while the simply invariant subspaces of D are much more complicated. In this paper, we remark on the complexity of the invariant subspaces F with $D_A \subset F \subset D$, where D_A is the analytic Dirichlet

Received by the editors October 31, 1994 and, in revised form, December 9, 1994.

1991 *Mathematics Subject Classification*. Primary 30H05; Secondary 30C15.

Key words and phrases. Dirichlet spaces, invariant subspaces, co-dimension, Bergman spaces.

This research was supported in part by a grant from the National Science Foundation.

space $\{f \in D : \hat{f}(n) = 0 \ \forall n < 0\}$, by examining their *index*. Here for an invariant subspace $F \subset D$, we define the *index* (or sometimes called the *co-dimension*) to be

$$\text{ind } F = \dim(F/\zeta F).$$

For the analytic Dirichlet space D_A , every non-zero invariant subspace has index 1 [13]. In stark contrast to this, it was observed in [1] that there are examples of invariant subspaces $D_A \subset F \subset D$ with $\text{ind } F = n$ for any $n \in \mathbf{N} \cup \{\infty\}$. We will give a new and more direct proof of this fact as well as specific examples of these types of subspaces. To do this, we will employ the natural one-to-one correspondence between the invariant subspaces $D_A \subset F \subset D$ and the invariant subspaces (under $f \rightarrow zf$) of the Bergman space $L^2(D)$. We will then use some recent results of Seip, Hedenmalm, and Richter [4], [5], [14] which yield specific examples of invariant subspaces of the Bergman space with large index. We then transfer this information back to the Dirichlet space to obtain the following theorem.

Theorem 1.1. *Given $n \in \mathbf{N} \cup \{\infty\}$, there are sequences $A_j, 0 \leq j < n$, of points in D such that*

$$F = \text{span}_{0 \leq j < n} \{ \bar{B}_a H^2 \cap D : a \in A_j \} \tag{1}$$

is a simply invariant subspace of D with $\text{ind } F = n$.

Note that $H^2 = \{f \in L^2 : \hat{f}(n) = 0 \ \forall n < 0\}$ is the usual Hardy space on the circle and

$$\bar{B}_a(\zeta) = \frac{\bar{a} - a\zeta}{|a|1 - \bar{a}\zeta}, \quad \zeta \in \mathbf{T}. \tag{1.1}$$

We also use $\text{span } X$ to denote the closed linear span of X , and \bar{z} to denote the complex conjugate of a complex number z .

We will remark at the end that a similar result also holds for the Dirichlet spaces D_α and the Besov classes $B_p^\alpha(\mathbf{T})$.

2. Bergman spaces

The sequences A_j in Theorem 1.1 will be zero sequences of the Bergman space, and thus we begin our discussion with some basic Bergman space facts. The *Bergman space* $L^2_a(D)$ is the space of analytic functions f on D such that

$$\int_D |f(z)|^2 dA(z) = \sum_{n=0}^{\infty} \frac{|a_n|^2}{n+1} < \infty.$$

Here dA is area measure on D normalized so it has mass 1, and $\{a_n\}$ are the power series coefficients of f . It is well known that $L^2_a(D)$ is a closed subspace of $L^2(D, dA)$ and the operator $f \rightarrow zf$ is continuous. The subspaces $M \subset L^2_a(D)$ with $zM \subset M$ (also called *invariant subspaces*) are tremendously complicated and poorly understood. As before, we define the *index* (or *co-dimension*) of an invariant subspace $M \subset L^2_a(D)$ to be $\text{ind } M = \dim(M/zM)$. It is known [2] that given $n \in \mathbf{N} \cup \{\infty\}$ there exists an invariant subspace $M \subset L^2_a(D)$ with $\text{ind } M = n$. Moreover, using sampling theory, Hedenmalm, Richter, and Seip [4], [5] have been able to generate specific examples of these types of invariant subspaces by using zero-based invariant subspaces.

Given a sequence $A \subset D$, we let

$$I(A) = \{f \in L^2_a(D) : f|_A = 0\}.$$

In the case where $I(A) \neq 0$ one can show that $I(A)$ is a closed invariant subspace of $L^2(D)$ with $\text{ind } I(A) = 1$. We now state the Seip, Hedenmalm, Richter result [4], [5].

Theorem 2.1 (Hedenmalm, Richter, Seip). *Given $n \in \mathbb{N} \cup \{\infty\}$ there are sequences A_j , $0 \leq j < n$, of points in D such that*

$$M = \text{span}\{I(A_j) : 0 \leq j < n\}$$

is an invariant subspace of $L^2(D)$ with $\text{ind } M = n$.

3. The correspondence

As mentioned in the introduction, there will be a natural correspondence between the invariant subspaces $D_A \subset F \subset D$ and the invariant subspaces of the Bergman space. This type of correspondence was observed by Makarov [9] in the $C^\infty(\mathbb{T})$ case where he observed a natural correspondence between the invariant subspaces $C_A^\infty(\mathbb{T}) \subset F \subset C^\infty(\mathbb{T})$ and the invariant subspaces (under $f \rightarrow zf$) of $A^{-\infty}$, the analytic distributions. Here $C_A^\infty(\mathbb{T}) = \{f \in C^\infty : \hat{f}(n) = 0 \ \forall n < 0\}$, and $A^{-\infty}$ are the distributions g with $\hat{g}(n) = 0$ for all $n < 0$. We make a similar observation for the Dirichlet space and also prove an interesting index formula.

The dual space of D , which we will denote by D^I , is the space of distributions g with

$$\sum_{n=-\infty}^{\infty} T_\infty \frac{|\hat{g}(n)|^2}{|n|+1} < \infty.$$

The pairing between D and D^I is given by

$$(f, g) = \sum_{n=-\infty}^{\infty} \hat{f}(n)\hat{g}(n), \quad f \in D, g \in D^I.$$

Let D_A^I be the analytic distributions of D^I

$$D_A^I = \{g \in D^I : \hat{g}(n) = (\zeta^n, g) = 0 \ \forall n < 0\}.$$

Making $\{\hat{g}(n)\}$ the power series coefficients of an analytic function $g(z)$ on the disk, we see that D_A^I can be naturally identified with the Bergman space $L^2_{\mathbb{D}}$ and moreover

$$(3.1) \quad (f, g) = \lim_{r \rightarrow 1^-} \int_{\mathbb{T}} \overline{f(r\zeta)} g(r\zeta) \frac{|d\zeta|}{2\pi}, \quad f \in D, g \in D_A^I.$$

Theorem 3.1. (1) *If $D_A \subset F \subset D$ is invariant, then $M = (\zeta F)^\perp$ is an invariant subspace of $L^2_{\mathbb{D}}$.*

(2) *If $M \subset L^2_{\mathbb{D}}$ is invariant, then $F = \overline{\zeta(M)}^\perp$ is invariant and $D_A \subset F \subset D$. $\text{ind } F$.*

Proof. (1) Letting $g \in M$, we see that $(\zeta f, g) = 0$ for all $f \in F$. But since $\zeta^n \in F$ for all $n \geq 0$, then $(\zeta^{-n-1}, g) = \hat{g}(-n-1) = 0$ for all $n \geq 0$, i.e. $g \in D_A^I = L^2_{\mathbb{D}}$. Also, by (3.1),

$$(\zeta f, zg) = (\zeta(\zeta f), g) = 0 \quad \forall f \in F$$

since F is invariant. Thus $zg \in M$, which makes M an invariant subspace of $L^2_{\mathbb{D}}$. $(M)^\perp$ and so

$$(\zeta(\zeta f), \bar{g}) = (\zeta f, \overline{zg}) = 0 \quad \forall g \in M$$

since M is invariant. Thus $\zeta f \in \zeta(M)^\perp = F$. Also, $M \subset L^2_a(D) = D^I_A$ and so

$$0 = (\zeta^{n+1}, g) = (\zeta \zeta^n, g) \quad \forall n \geq 0, g \in M,$$

which implies $\zeta^n \in \overline{\zeta(M)}^\perp = F$. Thus $D_A \subset F \subset D$ and F is invariant.

(3) To prove the index formula, first we notice from the Hahn-Banach theorem and basic linear algebra that

$$\begin{aligned} \dim(F/\zeta F) &= \dim((\zeta F)^\perp/F^\perp) \\ &= \dim(\overline{M}/\overline{\zeta(M)})^\perp \\ &= \dim(M/(\zeta(M^\perp))^\perp). \end{aligned}$$

The proof will be finished once we show that

$$(3.2) \quad (\zeta(M^\perp))^\perp = zM.$$

To this end, we let $g \in M$; then $zg \in M$ and for all $f \in M^\perp$,

$$(\zeta f, zg) = (\zeta \zeta f, g) = (f, g) = 0.$$

Thus $zg \in (\zeta(M^\perp))^\perp$.

For the other direction we observe that

$$D_A \subset F = \zeta(M)^\perp$$

and so $\zeta D_A \subset M^\perp$. Thus if $g \in (\zeta(M^\perp))^\perp$, then

$$0 = (g, \zeta \zeta) = (g, 1) = \tilde{g}(0).$$

Next we observe (using the invariance of M) that if $h \in M^\perp$,

$$0 = (h, k) = (h, zk) = (\zeta h, k) \quad \forall k \in M.$$

Thus $\zeta h \in M^\perp$. Hence $0 = (g, \zeta \zeta h) = (g, h)$ for all $h \in M^\perp$. Thus $g \in (M^\perp)^\perp = M$. Now using the fact that $g \in L^2_a(D)$ and $g(0) = 0$, we get that $\underset{z}{1}g \in L^2_a(D)$ and so, by (3.1),

$$0 = (g, \zeta h) = \underset{z}{1}g, \zeta h = \underset{z}{1}g, h \quad \forall h \in M^\perp.$$

Thus $\underset{z}{1}g \in M$ and hence $g \in zM$. So (3.2) has been established and the proof is complete. □

Recall from (1.1) that for $w \in D$, $B_w(\zeta)$ is the single Blaschke factor with zero at w . Using the fact that $I(\{w\}) = \text{span}\{z^n B_w(z) : n \geq 0\}$, and the F. and M. Riesz theorem, one can prove the following.

Lemma 3.2. *If $M = I(\{w\})$, then $F = \overline{\zeta(M)}^\perp = \overline{B_w}H^2 \cap D$.*

We are now ready to prove our main theorem.

Proof of Theorem 1.1. By Theorem 2.1, the invariant subspace

$$M = \text{span}\{I(A_j) : 0 \leq j < n\}$$

has index n . Thus by Theorem 3.1, $F = \zeta(\mathcal{M})^\perp$ also has index n . We now identify F :

$$\begin{aligned}
 F &= \zeta(\mathcal{M})^\perp \\
 &= \bar{\zeta} \left(\text{span} \{ \overline{I(A_j)} : 0 \leq j < n \} \right)^\perp \\
 &= \zeta \left(\left(\overline{I(A_j)} \right)^\perp \right)_{0 \leq j < n} \\
 &= \zeta \left(\text{span} \{ \overline{I(\{a_j\})} \} \right)^\perp \quad \text{since mult. by } \bar{\zeta} \text{ is invertible} \\
 &= \zeta \left(\text{span} \{ I(\{\bar{a}_j\})^\perp : a \in A_j \} \right)_{0 \leq j < n} \\
 &= \zeta \left(\text{span} \{ \zeta(I(\{\bar{a}_j\})^\perp) : a \in A_j \} \right)_{0 \leq j < n} \quad \text{since mult. by } \zeta \text{ is cont. and invertible} \\
 &= \zeta \left(\text{span} \{ \overline{B_a H^p} \cap D : a \in A_j \} \right)_{0 \leq j < n} \quad \text{by Lemma 3.2.} \quad \square
 \end{aligned}$$

4. Generalizations

In this last section, we remark that the analog of Theorem 1.1 is also true for the Dirichlet classes D_α , $0 < \alpha < \infty$, of $f \in L^2(\mathbb{T}, |d\zeta|)$ with

$$\sum_{n=-\infty}^{\infty} (1 + |n|)^\alpha |\hat{f}(n)|_2 < \infty,$$

and the Besov classes B_p^α , $1 < p < \infty$, $0 < \alpha < 1$, of f with

$$\int_0^\pi \int_0^{2\pi} \frac{|f(e^{i(\theta+h)}) - f(e^{i\theta})|_p}{h^{1+\alpha p}} d\theta dh < \infty.$$

Here the appropriate Bergman spaces to consider are the weighted Bergman spaces A_p^α , $0 < \alpha < \infty$, $1 < p < \infty$, of analytic functions on \mathbb{D} with

$$\int_{\mathbb{D}} |f(z)|^p (1 - |z|)^{\alpha p - 1} dA(z) < \infty.$$

The analog of Theorem 2.1 is true for A_p^α where, of course, $I(A) = \{f \in A_p^\alpha : f|_A = 0\}$ [4], [5]. For D_α , the analytic distributions (in the dual) can be identified with the weighted Bergman space $A_2^{\alpha/2}$ via (3.1) [7]. For B_p^α , the analytic distributions (in the dual) can be identified with A_q^α where $q = p(p-1)^{-1}$ via (3.1). For both these spaces, the analog of Theorem 3.1 remains true. Thus we have the following.

Theorem 4.1. *Let $X = D_\alpha$, $0 < \alpha < \infty$, or B_p^α , $1 < p < \infty$, $0 < \alpha < 1$, and N . Then there are sequences A_j , $0 \leq j < n$, of points in \mathbb{D} so that*

$$F = \zeta \left(\text{span}_X \{ \overline{B_a H^p} \cap X : a \in A_j \} \right)_{0 \leq j < n}$$

is a simply invariant subspace of X with $\text{ind } F = n$.

References

- [1] A. Aleman, S. Richter, and W.T. Ross, 'Bergman spaces on disconnected domains, *Canad. J. Math.* (to appear).
- [2] H. Bercovici, C. Foias, and C. Pearcy, *Dual algebras with applications to invariant subspaces and dilation theory*, CBMS Regional Conf. Ser. in Math., no. 56, Amer. Math. Soc., Providence, RI, 1985. MR **87g**:47091
- [3] M. Hasumi and T.P. Srinivasan, 'Invariant subspaces of continuous functions' , *Canad. J. Math.* **22** (1965), 643 - 651. MR **31**:3871
- [4] H. Hedenmalm, 'An invariant subspace of the Bergman space having the co-dimension two property' , *J. Reine Angew. Math.* **443** (1993), 1 - 9. MR **94k**:30092
- [5] H. Hedenmalm, S. Richter, K. Seip, 'Zero sequences and invariant subspaces in the Bergman space' , preprint.
- [6] H. Helson, *Lectures on invariant subspaces*, New York and London, 1964. MR **30**:1409
- [7] S. Khrushchev and V. Peller, 'Hankel operators, best approximation, and stationary Gaussian processes' , *Russian Math. Surveys.* **37** (1982), 61 - 144. MR **84e**:47036
- [8] N.G. Makarov, 'Sets of 1-invariant and $\bar{1}$ -invariant subspaces (smooth functions)' , *Soviet Math. Dokl.* **25** (1982), 191 - 194. MR **83h**:46053
- [9] N.G. Makarov, 'Invariant subspaces of the space $C^\infty(\mathbb{T})$ ', *Math. USSR Sbornik*, **47** (1984), 1 - 26. MR **84d**:30088
- [10] B. Malgrange, *Ideals of differentiable functions*, Tata Inst. Fund. Res., Bombay and Oxford Univ. Press, London, 1967. MR **35**:3446
- [11] Y.U. Netrusov 'Spectral synthesis in spaces of smooth functions' , *Russian Acad. Dokl. Math.*, **46** (1993), 135 - 138. MR **94c**:46068
- [12] S. Richter, W. T. Ross, and C. Sundberg, 'Hyperinvariant subspaces of the harmonic Dirichlet space' , *J. Reine Angew. Math.* **448** (1994), 1 - 26. MR **95e**:47045
- [13] S. Richter and A. Shields, 'Bounded analytic functions in the Dirichlet space' , *Math. Z.* **198** (1988), 151 - 159. MR **89c**:46039
- [14] K. Seip, 'Beurling type density theorems in the unit disk' , *Invent. Math.* **113** (1993), 26 - 39. MR **94g**:30033

Department of Mathematics, University of Richmond, Richmond, Virginia 23173
E-mail address: rossb@mathcs.urich.edu