

2013

Model Spaces: A Survey

William T. Ross

University of Richmond, wross@richmond.edu

Stephan Ramon Garcia

Follow this and additional works at: <http://scholarship.richmond.edu/mathcs-faculty-publications>



Part of the [Algebra Commons](#)

This is a pre-publication author manuscript of the final, published article.

Recommended Citation

Ross, William T. and Garcia, Stephan Ramon, "Model Spaces: A Survey" (2013). *Math and Computer Science Faculty Publications*. 123.
<http://scholarship.richmond.edu/mathcs-faculty-publications/123>

This Post-print Article is brought to you for free and open access by the Math and Computer Science at UR Scholarship Repository. It has been accepted for inclusion in Math and Computer Science Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact scholarshiprepository@richmond.edu.

MODEL SPACES: A SURVEY

STEPHAN RAMON GARCIA AND WILLIAM T. ROSS

ABSTRACT. This is a brief and gentle introduction, aimed at graduate students, to the subject of model subspaces of the Hardy space.

1. INTRODUCTION

In the summer of 2013, the authors gave a series of lectures and mini-courses in Montréal, Lens, and Helsinki on the topic of model spaces. In preparing for these lectures, we discovered the need for an easy introduction to model spaces suitable for the graduate students who formed the intended audience for our lectures. The standard texts on the subject [14, 67–69] are thorough and encyclopedic, but are sufficiently intimidating that the beginner might find this whole beautiful subject out of reach. The purpose of this survey is to give the novice a friendly, albeit incomplete, introduction to model spaces.

Model spaces are Hilbert spaces of the form $(uH^2)^\perp$, where u is an inner function, H^2 is the classical Hardy space on the open unit disk \mathbb{D} , and \perp denotes the orthogonal complement in H^2 . On a functional analysis level, model spaces are the orthogonal complements of the nontrivial invariant subspaces of the unilateral shift $Sf = zf$ on H^2 . These subspaces were characterized as uH^2 by Beurling in his famous 1949 paper [16]. As such, the spaces $(uH^2)^\perp$ are the invariant subspaces of the backward shift operator $S^*f = (f - f(0))/z$ on H^2 . However, unlike the spaces uH^2 which are simple to understand (i.e., all H^2 multiples of the inner function u), the model spaces $(uH^2)^\perp$ are much more troublesome. For instance, it is not immediately clear which functions actually belong $(uH^2)^\perp$ or what properties these functions have.

A major breakthrough in the study of model spaces occurred in 1970, with the publication of the seminal paper of Douglas, Shapiro, and Shields [31]. Extending some partial results of earlier authors, they showed that functions in $(uH^2)^\perp$ have analytic continuations in the same neighborhoods of points on the unit circle as does u . However, a generic inner function u need not have an analytic continuation across any point of the unit circle \mathbb{T} and thus a new type of continuation was needed. Fortunately this type of continuation, called a *pseudocontinuation*, was discovered and formalized in

First author partially supported by National Science Foundation Grant DMS-1265973.

two earlier papers [85, 86] of Shapiro and indeed turned out to be the determining characterization of functions in $(uH^2)^\perp$. In fact, a notion of continuation more general than analytic continuation was already being discussed, in various forms (rational approximation for example), often never actually being called a “continuation.” For example an earlier paper of Tumarkin [96] discussed functions in $(uH^2)^\perp$ via controlled approximation by rational functions.

From these beginnings, the function theoretic aspects of model spaces have truly grown. We now understand much of the subtle relationship between the boundary behavior of functions in $(uH^2)^\perp$ and the angular derivative of u through the papers of Ahern and Clark [3, 4].

On an operator theory level, work of Sz.-Nagy and Foias showed that the compression of the unilateral shift S to a model space $(uH^2)^\perp$ is a model for a certain class of contractions [93]. This seminal work has been studied by many people and continues to have relevance in operator theory [75]. Clark later examined unitary rank-one perturbations of this compressed shift and was able to come up with an exact spectral realization of this unitary operator using a family of measures that now bear his name [29]. These measures enjoy truly fascinating properties that have been harnessed by Aleksandrov [6, 9], Poltoratski [73], Sarason [72, 82, 83], and others to explore fine boundary properties of the inner function u as well as completeness of families of reproducing kernels in $(uH^2)^\perp$. These measures have even appeared in a somewhat different form in mathematical physics [90, 91].

Model spaces have also been helpful in examining nearly invariant subspaces (subspaces that contain f/z whenever they contain a function f satisfying $f(0) = 0$) [80]. These nearly invariant subspaces have been useful in developing characterizations of the invariant subspaces for the shift $f \mapsto zf$ on Hardy spaces of planar domains with holes or slits [10, 11, 56].

Though model spaces have many connections to old ideas in analysis (analytic continuation, factorization, pseudocontinuation, etc.), they continue to be relevant. Recent work of Makarov and Poltoratski [64] show that a spectral realization of certain selfadjoint Schrödinger operators can be realized through model spaces and Clark measures. Model spaces also make important connections to the subject of complex symmetric operators [39–43], a certain class of Hilbert space operators that are frequently modeled by compressions of Toeplitz operators to model spaces [28, 50, 92].

As the subject of model spaces is quite vast, we again emphasize that these notes are not meant to be an encyclopedic treatment. These notes are instead meant to give the beginning student a reason to want to study this material and to provide the means for them to take their first few steps into this rich and fertile territory.

2. PRELIMINARIES

Before proceeding, we start with a brief review of Hardy space theory. The material presented in this section is now considered classical and can be found in many standard texts [33, 52, 57, 59]. A relatively new text that might be more suitable for a student who is new to Hardy spaces is [66].

2.1. Lebesgue spaces. Let m denote normalized Lebesgue measure on the unit circle \mathbb{T} (i.e., $m = d\theta/2\pi$) and let $L^2 := L^2(\mathbb{T}, m)$ denote the space of m -measurable (i.e., Lebesgue measurable) functions $f : \mathbb{T} \rightarrow \mathbb{C}$ such that

$$\|f\| := \left(\int_{\mathbb{T}} |f(\zeta)|^2 dm(\zeta) \right)^{\frac{1}{2}}$$

is finite. As such, L^2 is a Hilbert space endowed with the inner product

$$\langle f, g \rangle := \int_{\mathbb{T}} f(\zeta) \overline{g(\zeta)} dm(\zeta).$$

A simple calculation using the fact that $m(\mathbb{T}) = 1$ shows that the family of functions $\{\zeta \mapsto \zeta^n : n \in \mathbb{Z}\}$ is an orthonormal basis for L^2 . The coefficients

$$\widehat{f}(n) := \langle f, \zeta^n \rangle = \int_{\mathbb{T}} f(\zeta) \overline{\zeta^n} dm(\zeta)$$

of a function f in L^2 with respect to this basis are called the (complex) *Fourier coefficients* of f . In light of Parseval's Identity

$$\|f\|^2 = \sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2,$$

we see that the L^2 norm of f coincides with the norm of the sequence $\{\widehat{f}(n) : n \in \mathbb{Z}\}$ of Fourier coefficients in the space $\ell^2(\mathbb{Z})$ of all square-summable sequences on \mathbb{Z} . We therefore identify the Hilbert spaces L^2 and $\ell^2(\mathbb{Z})$ via $f \leftrightarrow \{\widehat{f}(n) : n \in \mathbb{Z}\}$.

We also require the space $L^\infty := L^\infty(\mathbb{T})$ of all essentially bounded functions on \mathbb{T} which, when equipped with the norm

$$\|f\|_\infty := \text{ess-sup}_{\zeta \in \mathbb{T}} |f(\zeta)|,$$

becomes a Banach algebra. We also remark that for any φ in L^∞ , the multiplication operator $f \mapsto \varphi f$ on L^2 is bounded and has operator norm equal to $\|\varphi\|_\infty$.

2.2. Hardy spaces. For an analytic function f on \mathbb{D} the integral means

$$\int_{\mathbb{T}} |f(r\zeta)|^2 dm(\zeta)$$

are increasing as a function of r on $(0, 1)$. Indeed, if $f(z) = \sum_{n=0}^{\infty} a_n z^n$, then

$$\int_{\mathbb{T}} |f(r\zeta)|^2 dm(\zeta) = \sum_{n=0}^{\infty} |a_n|^2 r^{2n},$$

which is clearly increasing in r . This leads us to define the *Hardy space* H^2 as those f for which

$$\|f\| := \lim_{r \rightarrow 1^-} \left(\int_{\mathbb{T}} |f(r\zeta)|^2 dm(\zeta) \right)^{\frac{1}{2}} \quad (2.1)$$

is finite. It is no accident that we use $\|\cdot\|$ to denote both the norm in L^2 and in H^2 . Indeed, classical work of Fatou and Riesz show that the *radial limit*¹

$$f(\zeta) := \lim_{r \rightarrow 1^-} f(r\zeta) \quad (2.2)$$

exists for m -almost every ζ in \mathbb{T} and it turns out that the H^2 norm of f and the L^2 norm of its boundary function, defined in (2.2), coincide. In this manner, we often regard H^2 as a closed subspace of L^2 . As such, H^2 inherits the structure of a Hilbert space from L^2 .

Let H^∞ denote the space of all bounded analytic functions on \mathbb{D} , endowed with the supremum norm

$$\|f\|_\infty := \sup_{z \in \mathbb{D}} |f(z)|.$$

In a similar manner, the radial boundary function of an H^∞ function belongs to L^∞ and one has a corresponding maximum modulus type result

$$\sup_{z \in \mathbb{D}} |f(z)| = \text{ess-sup}_{\zeta \in \mathbb{T}} |f(\zeta)|,$$

that allows us to view H^∞ as a Banach subalgebra of L^∞ . Moreover, H^∞ happens to be the multiplier algebra for H^2 , meaning that the operator of multiplication by an analytic function φ on H^2 (i.e., $f \mapsto \varphi f$) is bounded if and only if φ belongs to H^∞ . The norm of this multiplication operator is precisely $\|\varphi\|_\infty$.

The inner product on the Hardy space H^2 is given by

$$\langle f, g \rangle = \int_{\mathbb{T}} f(\zeta) \overline{g(\zeta)} dm(\zeta) = \sum_{n=0}^{\infty} a_n \overline{b_n},$$

where $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$ denote typical elements of H^2 . In other words, we have a natural identification of H^2 with the sequence space $\ell^2(\mathbb{N})$, where each f in H^2 is identified with its sequence of Taylor coefficients $\{a_n\}_{n \geq 0}$.

Of great importance is the manner in which H^2 sits inside of L^2 . If $f(z) = \sum_{n=0}^{\infty} a_n z^n$ belongs to H^2 , then the almost everywhere defined boundary function f has an associated Fourier series

$$f \sim \sum_{n=0}^{\infty} a_n \zeta^n$$

¹It turns out that each f in H^2 has a finite *non-tangential limit* at ζ for almost every ζ . By this we mean that the limit of $f(z)$ exists as z approaches ζ in every *Stolz region* $\{z \in \mathbb{D} : |z - \zeta| < \alpha(1 - |z|)\}$, $\alpha > 1$.

which belongs to the first component in the direct sum

$$L^2 = H^2 \oplus \overline{zH^2},$$

where

$$\overline{zH^2} = \{\overline{zh} : h \in H^2\}.$$

In terms of Fourier coefficients,

$$H^2 = \bigvee \{z^n : n \geq 0\}, \quad \overline{zH^2} = \bigvee \{z^n : n \leq -1\},$$

where \bigvee denotes the closed linear span in L^2 . In particular, note that

$$a_n = \begin{cases} \widehat{f}(n) & \text{if } n \geq 0, \\ 0 & \text{if } n < 0, \end{cases}$$

and that the polynomials in z are dense in H^2 . These results are summarized in the following diagram:

$$\begin{array}{ccc} H^2 & \xleftrightarrow{\text{identified}} & \ell^2(\mathbb{N}) \\ \cap & & \cap \\ L^2 & \xleftrightarrow{\text{identified}} & \ell^2(\mathbb{Z}) \end{array}$$

2.3. The Cauchy-Szegő kernel. In light of the inequality

$$|f(\lambda)| \leq \sum_{n=0}^{\infty} |a_n| |\lambda|^n \leq \left(\sum_{n=0}^{\infty} |a_n|^2 \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} |\lambda|^{2n} \right)^{\frac{1}{2}} = \frac{\|f\|}{\sqrt{1-|\lambda|^2}},$$

which holds for all λ in \mathbb{D} and all f in H^2 , it follows that for fixed $\lambda \in \mathbb{D}$ the point evaluation functionals $f \mapsto f(\lambda)$ are bounded on H^2 and hence, by the Riesz Representation Theorem, must be of the form

$$f(\lambda) = \langle f, c_\lambda \rangle \tag{2.3}$$

for some c_λ in H^2 . In fact, it is not hard to show that

$$c_\lambda(z) = \frac{1}{1 - \overline{\lambda}z}, \quad \lambda \in \mathbb{D}, \tag{2.4}$$

which is called the *Cauchy-Szegő kernel* or, perhaps more frequently, the *Cauchy kernel*. In more general terms, one says that the Cauchy-Szegő kernel is the *reproducing kernel* for H^2 . Of great interest to us are other reproducing kernels, which satisfy equations analogous to (2.3) on various subspaces of H^2 (see Section 5).

Before proceeding, we should also remark that the reproducing formula (2.3) is simply a restatement of the identity

$$f(\lambda) = \sum_{n=0}^{\infty} a_n \lambda^n = \langle (a_0, a_1, a_2, \dots), (1, \overline{\lambda}, \overline{\lambda}^2, \dots) \rangle_{\ell^2(\mathbb{N})}.$$

When written as a contour integral, (2.3) reduces to the Cauchy Integral Formula

$$f(\lambda) = \int_{\mathbb{T}} \frac{f(\zeta)}{1 - \lambda\bar{\zeta}} dm(\zeta)$$

for H^2 functions (recall from (2.2) that for each f in H^2 , the almost everywhere defined boundary function $\zeta \mapsto f(\zeta)$ belongs to L^2 , allowing the preceding integral to be well-defined). Along similar lines, for fixed $n \geq 0$ we have

$$f^{(n)}(\lambda) = \langle f, c_\lambda^{(n)} \rangle,$$

where

$$c_\lambda^{(n)}(z) = \frac{n! \bar{\lambda}^n}{(1 - \bar{\lambda}z)^{n+1}}$$

is the n th derivative of c_λ with respect to the variable z .

Proposition 2.5. *The set $\{c_\lambda : \lambda \in \mathbb{D}\}$ is linearly independent.*

Proof. If $\lambda_1, \lambda_2, \dots, \lambda_n$ are distinct elements of \mathbb{D} and

$$\sum_{i=1}^n \alpha_i c_{\lambda_i} = 0,$$

then

$$\sum_{i=1}^n \bar{\alpha}_i f(\lambda_i) = 0$$

holds for all f in H^2 . The Lagrange Interpolation Theorem provides us with a polynomial $p(z)$ satisfying $p(\lambda_i) = \alpha_i$ so that $\sum_{i=1}^n |\alpha_i|^2 = 0$. Thus $\alpha_i = 0$ for $i = 1, 2, \dots, n$. \square

2.4. Canonical factorization. Although H^2 is a linear space, it is its multiplicative structure that reveals its true function-theoretic depth. We recall here the main ingredients necessary to describe the canonical factorization of H^2 functions.

Definition 2.6. An *inner function* is a bounded analytic function u on \mathbb{D} such that $|u(\zeta)| = 1$ for almost every ζ in \mathbb{T} .²

The simplest nontrivial example of an inner function is a Möbius transformation of the form

$$e^{i\theta} \frac{w - z}{1 - \bar{w}z},$$

²The reader is reminded that whenever we use the term “boundary function” or write $f(\zeta)$ for $f \in H^2$ and $\zeta \in \mathbb{T}$, we are referring to the almost everywhere defined radial (nontangential) limit in (2.2).

where $|w| < 1$ and $0 \leq \theta < 2\pi$, which is easily seen to be an automorphism of \mathbb{D} mapping \mathbb{T} onto \mathbb{T} . More generally, if $\{z_n\}_{n \geq 1}$ is a sequence of points in $\mathbb{D} \setminus \{0\}$, repeated according to multiplicity, then the *Blaschke condition*

$$\sum_{n=1}^{\infty} (1 - |z_n|) < \infty \quad (2.7)$$

is necessary and sufficient for the convergence (uniformly on compact subsets of \mathbb{D}) of the corresponding *Blaschke product*

$$B(z) := z^m \prod_{n=1}^{\infty} \frac{|z_n|}{z_n} \frac{z_n - z}{1 - \bar{z}_n z}, \quad (2.8)$$

where m denotes a nonnegative integer. With some work, one can show that every Blaschke product is an inner function [33, Thm. 2.4] (the only thing left to check is that the boundary function is unimodular almost everywhere). The importance of these functions stems from the fact that the Blaschke condition (2.7) completely characterizes the zero sets for H^2 functions.

Theorem 2.9. *A sequence $\{z_n\}_{n \geq 1} \subset \mathbb{D}$, repeated according to multiplicity, is the zero set of a nonconstant H^2 function if and only if it satisfies the Blaschke condition (2.7).*

Other examples of inner functions are furnished by the following construction. For a positive, finite, singular (with respect to m) Borel measure μ on \mathbb{T} , we claim that the analytic function

$$S_\mu(z) := \exp \left(- \int \frac{\zeta + z}{\zeta - z} d\mu(\zeta) \right), \quad z \in \mathbb{D}, \quad (2.10)$$

is inner. Such a function is known as a *singular inner function*. First notice that for any $z \in \mathbb{D}$,

$$\begin{aligned} |S_\mu(z)| &= \exp \left[\Re \left(- \int \frac{\zeta + z}{\zeta - z} d\mu(\zeta) \right) \right] \\ &= \exp \left(- \int_{\mathbb{T}} \frac{1 - |z|^2}{|\zeta - z|^2} d\mu(\zeta) \right) \\ &\leq 1 \end{aligned}$$

since both the measure μ and the *Poisson kernel*

$$P_z(\zeta) := \frac{1 - |z|^2}{|\zeta - z|^2}, \quad (\zeta \in \mathbb{T}, z \in \mathbb{D})$$

are nonnegative. Since S_μ belongs to H^∞ , it follows that S_μ has nontangential boundary values m -a.e. on \mathbb{T} . To show that these boundary values are almost everywhere unimodular, we require some basic facts from the

theory of harmonic functions. Let

$$(D\mu)(w) := \lim_{t \rightarrow 0^+} \frac{\mu((e^{-it}w, e^{it}w))}{2t} \quad (w \in \mathbb{T})$$

denote the *symmetric derivative* of μ on \mathbb{T} , where $(e^{-it}w, e^{it}w)$ denotes the circular arc subtended by the points $e^{-it}w$ and $e^{it}w$. We also have the identity

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} P_{rw}(\zeta) d\mu(\zeta) = (D\mu)(w). \quad m\text{-a.e. } w \in \mathbb{T}. \quad (2.11)$$

Since μ is singular, it follows that $D\mu = 0$ holds m -almost everywhere and so it now follows from the identity

$$|S_\mu(z)| = \exp\left(-\int_{\mathbb{T}} P_z(\zeta) d\mu(\zeta)\right)$$

that S has unimodular boundary values m -almost everywhere. For instance, if $\mu = \delta_1$ denotes the point mass at $\zeta = 1$, then

$$S_{\delta_1}(z) = \exp\left(\frac{z+1}{z-1}\right).$$

This type of inner function is often called an *atomic inner function*.

Theorem 2.12. *Every inner function u can be factored uniquely as*

$$u = e^{i\gamma} B_\Lambda S_\mu,$$

where $\gamma \in [0, 2\pi)$, Λ is a Blaschke sequence, and μ is a positive singular measure on \mathbb{T} . Conversely, any such product is inner.

The reader is warned that, as an abuse of language, one frequently permits a function of the form $e^{i\gamma} B_\Lambda$ (resp. $e^{i\gamma} S_\mu$) to be called a Blaschke product (resp. a singular inner function). For the sake of convenience, we adopt this common practice here in these notes.

Definition 2.13. Let u_1, u_2 be inner functions and $f \in H^2$.

- (i) We say that u_1 *divides* u_2 , written $u_1|u_2$, if $u_2/u_1 \in H^\infty$.
- (ii) We say that u_1 *divides* f , written $u_1|f$, if $f/u_1 \in H^2$.
- (iii) We say that u_1 and u_2 are *relatively prime* if the only inner divisors of both u_1 and u_2 are constant functions of unit modulus.

For example, if u_1 and u_2 are Blaschke products with simple zeros, then u_1 divides u_2 if and only if the zero set of u_1 is contained in the zero set of u_2 . Moreover, u_1 is relatively prime to u_2 if and only if u_1 and u_2 have no common zeros.

Definition 2.14. An *outer function* is an analytic function F on \mathbb{D} of the form

$$F(z) = e^{i\gamma} \exp\left(\int_{\mathbb{T}} \frac{\zeta+z}{\zeta-z} \varphi(\zeta) dm(\zeta)\right), \quad (2.15)$$

where γ is a real constant and φ is a real-valued function in L^1 .

The significance of the somewhat unwieldy formula in (2.15) lies in the fact that

$$\log |F(z)| = \int_{\mathbb{T}} P_z(\zeta) \varphi(\zeta) dm(\zeta),$$

from which it follows that $\log |F(z)|$ equals the harmonic extension to \mathbb{D} of the boundary function $\varphi : \mathbb{T} \rightarrow \mathbb{R}$ so that $\varphi = \log |F|$ a.e. on \mathbb{T} . In particular, an outer function is determined up to a unimodular constant factor by its modulus on \mathbb{T} . On the other hand, it is possible to show that every f in H^2 that does not vanish identically satisfies

$$\int_{\mathbb{T}} \log |f(\zeta)| dm(\zeta) > -\infty$$

and hence one can form the outer function (2.15) corresponding to the boundary data $\varphi = \log |f|$ [33, Thm. 2.2]. Putting this all together, we can now state the canonical factorization for H^2 functions [33, Thm. 2.8].

Theorem 2.16. *Every function f in $H^2 \setminus \{0\}$ has a unique factorization of the form*

$$f = BSF \tag{2.17}$$

where B is a Blaschke product, S is a singular inner function, and F is an outer function in H^2 . Conversely, any product of the form (2.17) belongs to H^2 .

2.5. Bounded type. In order to adequately discuss the cyclic vectors for the backward shift operator, we need two more additional classes of meromorphic functions on \mathbb{D} .

Definition 2.18. Let f be a meromorphic function on \mathbb{D} .

- (i) We say f is of *bounded type* if f can be written as quotient of two bounded analytic functions. The set of functions of bounded type is denoted by N (often called the Nevanlinna class).
- (ii) We say f is in the *Smirnov class* if it is the quotient of two bounded analytic functions where the denominator is an outer function. The Smirnov class is denoted by N^+ . Notice in this case that N^+ is a space of analytic functions on \mathbb{D} since the denominator is outer and outer functions have no zeros on \mathbb{D} .

Remark 2.19. It is known that $H^2(\mathbb{D}) \subset N^+$ and that every $f \in N$ has finite non-tangential limits almost everywhere [33, Ch. 2]. Furthermore, if $f \in N^+$ and the boundary function belongs to L^2 , then $f \in H^2$. This fact is no longer true for $f \in N^3$

³Even if f is analytic on \mathbb{D} , this is no longer true. Just consider the function $f(z) = \exp(-\frac{z+1}{z-1})$, which is the reciprocal of the atomic inner function defined earlier. This function belongs to N , is analytic on \mathbb{D} , has unimodular boundary values, but does not belong to H^2 since some integral estimates will show that f does not have bounded integral means.

2.6. Beurling's Theorem. Of supreme importance in the world of operator-related function theory are the shift operators. Chief among these is the *unilateral shift* $S : H^2 \rightarrow H^2$ defined by

$$[Sf](z) = zf(z),$$

or, in terms of Taylor coefficients, by

$$S(a_0, a_1, \dots) = (0, a_0, a_1, \dots).$$

Because of its ubiquity in the realm of operator theory, one often refers to S as *the* shift operator. One easily sees that S is an isometry that is not unitary. The adjoint S^* of the unilateral shift is the *backward shift* $S^* : H^2 \rightarrow H^2$ given by

$$[S^*f](z) = \frac{f(z) - f(0)}{z},$$

or, in terms of Taylor coefficients, by

$$S^*(a_0, a_1, \dots) = (a_1, a_2, \dots).$$

If u is inner, then the operator $f \mapsto uf$ is an isometry on H^2 and thus uH^2 is a subspace (i.e., a closed linear manifold) of H^2 . Moreover, assuming that u is a nonconstant inner function, that uH^2 is a nontrivial invariant subspace for the operator S . A celebrated theorem of Beurling [16] (also see any of the standard texts mentioned above for a complete proof and further discussions and generalizations) says that these are all of them.

Theorem 2.20 (Beurling's Theorem). *The nontrivial invariant subspaces of H^2 for the unilateral shift S are precisely the subspaces*

$$uH^2 := \{uh : h \in H^2\},$$

where u is an inner function. Moreover, f is cyclic for S , i.e.,

$$\bigvee \{f, Sf, S^2f, \dots\} = \bigvee \{qf : q \text{ is a polynomial}\} = H^2,$$

if and only if f is an outer function.

From an operator theoretic perspective, Beurling's Theorem is notable for providing an explicit description of the lattice of invariant subspaces for the unilateral shift operator. Indeed,

$$u_1H^2 \subseteq u_2H^2 \iff u_1/u_2 \in H^\infty. \quad (2.21)$$

For our purposes, however, it is the invariant subspaces for the backward shift that are of greatest importance. These are the so-called *model spaces* that are the primary focus of our investigations.

3. MODEL SPACES

The details of much of the following material can be found in the original sources, which we attempt to quote whenever possible, as well as the texts [26,27,67,68,75,76]. On the other hand, many of the results discussed below are part of the folklore on the subject and occasionally proper references cannot be readily identified.

3.1. Basic properties. We are now ready to introduce our primary object of study.

Definition 3.1. If u is an inner function, then the corresponding *model space* is

$$\mathcal{K}_u := (uH^2)^\perp = H^2 \ominus uH^2. \quad (3.2)$$

The definition above is somewhat unenlightening since the function theoretic properties of the model space \mathcal{K}_u do not immediately present themselves. A more direct description of \mathcal{K}_u via the boundary values of these functions comes from the following result.

Proposition 3.3. For inner u , the model space \mathcal{K}_u is the set of functions f in H^2 such that $f = \overline{g}zu$ almost everywhere on \mathbb{T} for some g in H^2 . In other words,

$$\mathcal{K}_u = H^2 \cap \overline{uzH^2},$$

where the right hand side is regarded as a set of functions on \mathbb{T} .

Proof. For each f in H^2 , we see that

$$\langle f, uh \rangle = 0, \forall h \in H^2 \iff \langle \overline{u}f, h \rangle = 0, \forall h \in H^2 \iff \overline{u}f \in \overline{zH^2}.$$

Since $u\overline{u} = 1$ almost everywhere on \mathbb{T} , we see that an H^2 function f belongs to the orthocomplement of uH^2 if and only if f belongs to $\overline{uzH^2}$ (i.e., $f = \overline{g}zu$ for some g in H^2).⁴ \square

Just as the Beurling-type subspaces uH^2 constitute the nontrivial invariant subspaces for the unilateral shift on H^2 , the subspaces \mathcal{K}_u play an analogous role for the backward shift.

Corollary 3.4. The model spaces \mathcal{K}_u , where u is inner, are precisely the proper invariant subspaces of H^2 for the backward shift

$$f(z) \mapsto \frac{f(z) - f(0)}{z}, \quad (a_0, a_1, a_2, \dots) \mapsto (a_1, a_2, a_3, \dots). \quad (3.5)$$

The following corollary is an immediate consequence of (2.21) (recall what it means for one inner function to *divide* another from Definition 2.13).

Corollary 3.6. If u_1, u_2 are inner functions, then

$$u_1|u_2 \iff \mathcal{K}_{u_1} \subseteq \mathcal{K}_{u_2}.$$

⁴In fact, g actually belongs to \mathcal{K}_u as well (see Section 7).

Remark 3.7. We pause here to mention the mildly surprising fact, not widely known, that if u is not a finite Blaschke product, then \mathcal{K}_u contains a linearly ordered chain of S^* -invariant subspaces of uncountable length. This is obvious if u has a singular inner factor S_μ since $\mathcal{K}_{S_{\alpha\mu}} \subseteq \mathcal{K}_u$ for $0 \leq \alpha \leq 1$. On the other hand, if u is an infinite Blaschke product, then things are not so clear. Suppose that $u = \prod_{n=1}^{\infty} b_n$ where each b_n is a Blaschke factor. Let $I \subseteq \mathbb{R}$ be a nonempty interval and let $\tau : \mathbb{Q} \cap I \rightarrow \mathbb{N}$ be a bijection. For each α in I , define $I_\alpha = \{q \in \mathbb{Q} \cap I : q < \alpha\}$ and notice that each α corresponds to a distinct Blaschke product $u_\alpha = \prod_{n \in \tau(I_\alpha)} b_n$ that divides u . This yields the desired chain of S^* -invariant subspaces.

Building upon the proof of Proposition 3.3, we have the following important result [31, Thm. 3.1.5].

Proposition 3.8. *A function f in H^2 is noncyclic for S^* if and only if there exists a function g in H^2 and an inner function u such that*

$$f = \overline{g}zu \quad (3.9)$$

almost everywhere on \mathbb{T} . If u and the inner factor of g are relatively prime, then the S^ -invariant subspace of H^2 generated by f is \mathcal{K}_u itself:*

$$\bigvee \{f, S^*f, S^{*2}f, \dots\} = \mathcal{K}_u.$$

Remark 3.10. We will see another description of \mathcal{K}_u as well as the noncyclic vectors for S^* when we discuss pseudocontinuations in Section 6.

Being invariant under the backward shift operator, it is not surprising that the spaces \mathcal{K}_u are also invariant under certain functions of the backward shift. In what follows, we let $P : L^2 \rightarrow H^2$ denote the *Riesz projection*

$$P(\dots, a_{-1}, a_0, a_1, a_2, \dots) = (a_0, a_1, a_2, \dots), \quad (3.11)$$

the orthogonal projection which returns the “analytic part” of a Fourier series in L^2 . For instance, $P(1 + 2 \cos \theta) = P(e^{-i\theta} + 1 + e^{i\theta}) = 1 + e^{i\theta} = 1 + \zeta$. We also remark that, as an orthogonal projection, the operator P is self-adjoint and hence satisfies $\langle Pf, g \rangle = \langle f, Pg \rangle$ for all f, g in L^2 .

Definition 3.12. For φ in $L^\infty(\mathbb{T})$ the *Toeplitz operator* $T_\varphi : H^2 \rightarrow H^2$ with *symbol* φ is defined by

$$T_\varphi(f) = P(\varphi f).$$

The study of Toeplitz operators is a vast subject and we refer the reader to the texts [32,66] for the basics and to [18] for an encyclopedia on the subject. Note that when φ belongs to H^∞ , the Toeplitz operator $T_\varphi f = \varphi f$ is just a multiplication operator. A simple calculation shows that T_z and $T_{\bar{z}}$ are precisely the forward and backward shift operators on H^2 . The operator $T_z : H^2 \rightarrow H^2$ enjoys an H^∞ -functional calculus given by $\varphi(T_z) = T_\varphi$ for φ in H^∞ [93]. Being $T_{\bar{z}}$ -invariant already, it should come as little surprise that the spaces \mathcal{K}_u are also invariant under conjugate-analytic Toeplitz operators (i.e., Toeplitz operators of the form $T_{\bar{\varphi}}$ where φ belongs to H^∞).

Proposition 3.13. *If $\varphi \in H^\infty$ and u is inner then $T_{\overline{\varphi}}\mathcal{K}_u \subseteq \mathcal{K}_u$.*

Proof. For each f in \mathcal{K}_u we have

$$\langle T_{\overline{\varphi}}f, uh \rangle = \langle P(\overline{\varphi}f), uh \rangle = \langle \overline{\varphi}f, P(uh) \rangle = \langle \overline{\varphi}f, uh \rangle = \langle f, u(\varphi h) \rangle = 0. \quad \square$$

Proposition 3.13 shows that the spaces \mathcal{K}_u enjoy the so-called *F-property*.⁵

Proposition 3.14. *If f belongs to \mathcal{K}_u and θ is an inner function that divides f , i.e., f/θ belongs to H^2 , then f/θ also belongs to \mathcal{K}_u . In particular, the outer factor of any function in \mathcal{K}_u also belongs to \mathcal{K}_u .*

Proof. Simply observe that $T_{\overline{\theta}}f = P(\overline{\theta}f) = P(f/\theta) = f/\theta$ since f/θ is in H^2 . By the preceding proposition, f/θ belongs to \mathcal{K}_u . \square

3.2. Finite dimensional model spaces. The simplest examples of model spaces are those corresponding to finite Blaschke products

$$u(z) = \prod_{j=1}^n \frac{z - \lambda_j}{1 - \overline{\lambda_j}z}. \quad (\lambda_j \in \mathbb{D})$$

Indeed, these are the only model spaces that are finite-dimensional and whose elements can be completely characterized in an explicit fashion. To do this, one should first notice that the Cauchy-Szegő kernel c_λ (from (2.4)) belongs to \mathcal{K}_u whenever λ is a zero of u . Indeed, if $u(\lambda) = 0$, then clearly $\langle uh, c_\lambda \rangle = u(\lambda)h(\lambda) = 0$ for all h in H^2 .

Proposition 3.15. *If u is a finite Blaschke product with zeros $\lambda_1, \lambda_2, \dots, \lambda_n$, repeated according to their multiplicity, then*

$$\mathcal{K}_u = \left\{ \frac{a_0 + a_1z + \dots + a_{n-1}z^{n-1}}{(1 - \overline{\lambda_1}z)(1 - \overline{\lambda_2}z) \dots (1 - \overline{\lambda_n}z)} : a_0, a_1, \dots, a_{n-1} \in \mathbb{C} \right\}. \quad (3.16)$$

In particular, \mathcal{K}_{z^n} is the space of all polynomials of degree $\leq n - 1$.

Proof. We provide the proof in the special case where the zeros λ_i are distinct. Since $\langle uh, c_{\lambda_i} \rangle = u(\lambda_i)h(\lambda_i) = 0$ for all h in H^2 , it follows that

$$\text{span}\{c_{\lambda_1}, c_{\lambda_2}, \dots, c_{\lambda_n}\} \subseteq \mathcal{K}_u.$$

If $f(\lambda_i) = \langle f, c_{\lambda_i} \rangle = 0$ for all i , then $u|f$ and hence f belongs to uH^2 . Thus

$$\text{span}\{c_{\lambda_1}, c_{\lambda_2}, \dots, c_{\lambda_n}\}^\perp \subseteq \mathcal{K}_u^\perp.$$

Since $\mathcal{K}_u = \text{span}\{c_{\lambda_1}, c_{\lambda_2}, \dots, c_{\lambda_n}\}$ the result follows by simple algebra. To be more specific, any linear combination of the Cauchy kernels $\{c_{\lambda_j} : 1 \leq j \leq n\}$ can be expressed as a rational function of the type prescribed in (3.16). Conversely, any expression of the type encountered in (3.16) can be decomposed, via partial fractions, into a linear combination of the functions $c_{\lambda_1}, c_{\lambda_2}, \dots, c_{\lambda_n}$. \square

⁵In general, a set \mathcal{C} of functions contained in H^2 has the *F-property* if whenever θ divides f , then $f/\theta \in \mathcal{C}$. Good sources for this are [88, 89].

If λ has multiplicity m as a zero of u , then one must also include the functions

$$c_\lambda, c'_\lambda, c''_\lambda, \dots, c_\lambda^{(m-1)}$$

in place of c_λ in the preceding proof. Along similar lines, the proof of Proposition 3.15 and the preceding comment provides us with the following useful fact.

Proposition 3.17. *Suppose that u is the finite Blaschke product with distinct zeros $\lambda_1, \lambda_2, \dots, \lambda_n$ with respective multiplicities m_1, m_2, \dots, m_n , then*

$$\mathcal{K}_u = \text{span}\{c_{\lambda_i}^{(\ell_i-1)} : 1 \leq i \leq n, 1 \leq \ell_i \leq m_i\}.$$

In fact, the preceding observation makes it clear why $\dim \mathcal{K}_u < \infty$ occurs if and only if u is a finite Blaschke product. If u has a factor that is an infinite Blaschke product, then Proposition 2.5 ensures that \mathcal{K}_u contains an infinite, linearly independent set (namely the Cauchy kernels corresponding to the distinct zeros of u). On the other hand, if u is a singular inner function, then $u^{1/n}$ is an inner function that divides u whence $\mathcal{K}_{u^{1/n}} \subseteq \mathcal{K}_u$ for $n \geq 1$. We may find an infinite orthonormal sequence in \mathcal{K}_u by selecting unit vectors f_n in $\mathcal{K}_{u^{1/n}} \ominus \mathcal{K}_{u^{1/(n+1)}}$, from which it follows that $\dim \mathcal{K}_u = \infty$. For the general case we point out the following decomposition that is interesting in its own right.

Proposition 3.18. *If $\{u_j\}_{j \geq 1}$ is a possibly finite sequence of inner function such that $u = \prod_{j \geq 1} u_j$ exists, then*

$$\mathcal{K}_u = \mathcal{K}_{u_1} \oplus \bigoplus_{n \geq 2} \left(\prod_{j=1}^{n-1} u_j \right) \mathcal{K}_{u_n}.$$

In particular, if u and v are inner functions, then

$$\mathcal{K}_{uv} = \mathcal{K}_u \oplus u\mathcal{K}_v. \quad (3.19)$$

We refer the reader to [3, 61] for further details, although we provide a proof of the special case (3.19) in Subsection 5.1.

3.3. Three unitary operators. The following three transformations from one model space to another are often useful.

Proposition 3.20. *Suppose u is a fixed inner function.*

(i) *If $w \in \mathbb{D}$, then*

$$f \mapsto \frac{\sqrt{1-|w|^2}}{1-\bar{w}u} f$$

defines a unitary operator from \mathcal{K}_u onto $\mathcal{K}_{\frac{u-w}{1-\bar{w}u}}$.

(ii) *If φ is a disk automorphism, i.e., $\varphi(z) = \zeta(z-a)(1-\bar{a}z)^{-1}$ for some $a \in \mathbb{D}$ and $\zeta \in \mathbb{T}$, then*

$$f \mapsto \sqrt{\varphi'}(f \circ \varphi)$$

defines a unitary operator from \mathcal{K}_u onto $\mathcal{K}_{u \circ \varphi}$.

(iii) If $u^\#(z) := \overline{u(\bar{z})}$, then, in terms of boundary functions, the map

$$f(\zeta) \mapsto \bar{\zeta} f(\bar{\zeta}) u^\#(\zeta)$$

is a unitary operator from \mathcal{K}_u into $\mathcal{K}_{u^\#}$.

The first unitary operator is due to Crofoot [30] (a more detailed discussion of these so-called *Crofoot transforms* can be found in [83, Sec. 13]). When dealing with a model space \mathcal{K}_u where u has a nontrivial Blaschke factor, one can often make the simplifying assumption that $u(0) = 0$. Also of great importance is the fact that Crofoot transforms intertwine the conjugations (see Section 7) on the corresponding model spaces [83, Lem. 3.1]. The second unitary operator is clearly unitary on H^2 (change of variables formula). Showing that its restriction to \mathcal{K}_u has the correct range is a little tricky and the proof of this can be found in [28, Prop. 4.1]. The third unitary operator depends on a discussion on conjugations that we take up in more detail in Section 7. Indeed, on the face of it, the map does not even seem to take analytic functions to analytic functions. However, when one thinks of model space functions in terms of their boundary values as in Proposition 3.3, everything works out. The proof can be found in [28, Lemma 4.3].

4. MODEL OPERATORS

One of the main reasons that model spaces are worthy of study in their own right stems from the so-called *model theory* developed by Sz.-Nagy and Foiaş, which shows that a wide range of Hilbert space operators can be realized concretely as restrictions of the backward shift operator to model spaces. These ideas have since been generalized in many directions (e.g., de Branges-Rovnyak spaces, vector-valued Hardy spaces, etc.) and we make no attempt to provide an encyclopedic account of the subject, referring the read instead to the influential texts [14, 67, 69, 75, 93].

4.1. Contractions. In the following, we let \mathcal{H} denote a separable complex Hilbert space. If T is an arbitrary bounded operator on \mathcal{H} , then we may assume that $\|T\| \leq 1$ (i.e., T is a *contraction*). As such, T enjoys a decomposition of the form $T = K \oplus U$ (see [93, p. 8] for more details) where U is a unitary operator and K is a *completely nonunitary (CNU) contraction* (i.e., there does not exist a reducing subspace for K upon which K is unitary).

Since the structure and behavior of unitary operators is well-understood, via the spectral theorem, the study of arbitrary bounded Hilbert space operators can therefore be focused on CNU contractions. With a few additional hypotheses, one can obtain a concrete *functional model* for such operators. In light of the fact that the following theorem justifies, to an extent, the further study of model subspaces, we feel obliged to provide a complete proof. Moreover, the proof itself is surprisingly simple and is worthy of admiration for its own sake.

Theorem 4.1 (Sz.-Nagy-Foiaş). *If T is a contraction on a Hilbert space that satisfies*

- (i) $\|T^n \mathbf{x}\| \rightarrow 0$ for all $\mathbf{x} \in \mathcal{H}$,
- (ii) $\text{rank}(I - T^*T) = \text{rank}(I - TT^*) = 1$,

then there exists an inner function u such that T is unitarily equivalent to $S^|_{\mathcal{K}_u}$, where S^* is the backward shift operator on H^2 .*

Proof. We let \cong denote the unitary equivalence of Hilbert spaces or their operators. Since the defect operator $D = \sqrt{I - T^*T}$ has rank 1, we see that $\text{ran } D \cong \mathbb{C}$ so that

$$\tilde{\mathcal{H}} := \bigoplus_{n=1}^{\infty} \text{ran } D \cong H^2. \quad (4.2)$$

It follows that for each $n \in \mathbb{N}$ we have

$$\begin{aligned} \sum_{j=0}^n \|DT^j \mathbf{x}\|^2 &= \sum_{j=0}^n \langle (I - T^*T)^{\frac{1}{2}} T^j \mathbf{x}, (I - T^*T)^{\frac{1}{2}} T^j \mathbf{x} \rangle \\ &= \sum_{j=0}^n \langle (I - T^*T) T^j \mathbf{x}, T^j \mathbf{x} \rangle \\ &= \sum_{j=0}^n \left(\langle T^j \mathbf{x}, T^j \mathbf{x} \rangle - \langle T^* T T^j \mathbf{x}, T^j \mathbf{x} \rangle \right) \\ &= \sum_{j=0}^n (\|T^j \mathbf{x}\|^2 - \|T^{j+1} \mathbf{x}\|^2) \\ &= \|\mathbf{x}\|^2 - \|T^{n+1} \mathbf{x}\|^2. \end{aligned}$$

Since, by hypothesis, $\|T^n \mathbf{x}\| \rightarrow 0$ for each $\mathbf{x} \in \mathcal{H}$, we conclude that

$$\sum_{j=0}^{\infty} \|DT^j \mathbf{x}\|^2 = \|\mathbf{x}\|^2 \quad (\mathbf{x} \in \mathcal{H})$$

and hence the operator $\Phi : \mathcal{H} \rightarrow H^2$ defined by

$$\Phi \mathbf{x} = (D\mathbf{x}, DT\mathbf{x}, DT^2\mathbf{x}, DT^3\mathbf{x}, \dots)$$

is an isometric embedding of \mathcal{H} into H^2 (here we have identified a function in H^2 with its sequence of Taylor coefficients). Since Φ is an isometry, its image

$$\Phi \mathcal{H} = (D\mathcal{H}, DT\mathcal{H}, DT^2\mathcal{H}, \dots)$$

is closed in H^2 and clearly S^* -invariant. Therefore, by Corollary 3.4, $\text{ran } \Phi = \mathcal{K}_u$ for some u (the possibility that $\text{ran } \Phi = H^2$ is ruled out because the following argument would show that $T \cong S^*$, violating the assumption that $\text{rank}(I - TT^*) = 1$). Now observe that

$$\Phi T\mathbf{x} = (DT\mathbf{x}, DT^2\mathbf{x}, DT^3\mathbf{x}, \dots) = S^* \Phi \mathbf{x}. \quad (\mathbf{x} \in \mathcal{H})$$

Letting $U : \mathcal{H} \rightarrow \text{ran } \mathcal{K}_u$ denote the unitary operator obtained from Φ by reducing its codomain from H^2 to \mathcal{K}_u , it follows that

$$UT = (S^*|_{\mathcal{K}_u})U.$$

Thus T is unitarily equivalent to the restriction of S^* to \mathcal{K}_u . □

The case of higher defect indices (i.e., $\text{rank}(I - T^*T) = \text{rank}(I - TT^*) \geq n$, $n > 1$), is treated by moving to vector-valued Hardy spaces $H^2(\mathcal{E})$ and operator-valued inner functions. However, in making such a move one sacrifices a large variety of tools and techniques inherited from classical function theory (making the theory of model spaces more difficult and less interesting from our perspective). For instance, the multiplication of operator-valued inner functions is no longer commutative and the corresponding factorization theory is more complicated.

4.2. Spectrum of an inner function. In light of the Sz.-Nagy-Foiaş Theorem, the restriction of the backward shift to the spaces \mathcal{K}_u is of premier importance. However, it turns out that the *compressed shift*, the compression $S_u : \mathcal{K}_u \rightarrow \mathcal{K}_u$ of the unilateral shift to \mathcal{K}_u , is more prevalent in the literature. Here

$$S_u f = P_u(zf),$$

where P_u denotes the orthogonal projection from L^2 onto \mathcal{K}_u (see Subsection 5.1). In reality, the distinction alluded to above is artificial since the operator S_u is unitarily equivalent to the restriction of the backward shift to the space $\mathcal{K}_{u^\#}$ where $u^\#(z) = \overline{u(\bar{z})}$. In fact the unitary operator which intertwines S_u and $S^*|_{\mathcal{K}_{u^\#}}$ is the one given in Proposition 3.20. See [28, Lemma 4.5] for more details on this.

There happens to be two convenient ways to describe the spectrum $\sigma(S_u)$ of S_u . In fact, one can show that $\sigma(S_u)$ coincides with the so-called *spectrum* of the inner function u , as defined below (this result is sometimes referred to as the Livšic-Möller Theorem).

Definition 4.3. The *spectrum* of an inner function is the set

$$\sigma(u) := \left\{ \lambda \in \mathbb{D}^- : \liminf_{z \rightarrow \lambda} |u(z)| = 0 \right\}. \tag{4.4}$$

It follows from the preceding definition, for instance, that every zero of u in \mathbb{D} belongs to $\sigma(u)$. Moreover, any limit point of zeros of u must also lie in $\sigma(u)$. On the other hand, if u is the singular inner function associated to the singular measure μ , then the nontangential limit of u is zero μ -almost everywhere [52, Thm. 6.2]. Even further, if $\lambda \in \mathbb{D}$ and $u(\lambda) = 0$ then $c_\lambda(z) = (1 - \bar{\lambda}z)^{-1}$ belongs to \mathcal{K}_u and satisfies $S^*c_\lambda = \bar{\lambda}c_\lambda$. In other words, $\bar{\lambda}$ belongs to $\sigma_p(S^*|_{\mathcal{K}_u})$, the point spectrum (i.e., set of eigenvalues) of $S^*|_{\mathcal{K}_u}$. These observations suggest the following important theorem that provides us with a convenient description of the spectrum of an inner function in terms of its canonical factorization.

Theorem 4.5. *If $u = B_\Lambda S_\mu$, where B_Λ is a Blaschke product with zero sequence Λ and S_μ is a singular inner function with corresponding singular measure μ , then*

- (i) $\sigma(S_u) = \sigma(u) = \Lambda^- \cup \text{supp } \mu$,
- (ii) $\sigma_p(S^*|\mathcal{K}_u) = \{\bar{\lambda} : \lambda \in \Lambda\}$,
- (iii) $\sigma_p(S_u) = \Lambda$,

where Λ^- denotes the closure of Λ .

5. REPRODUCING KERNELS

As a closed subspace of the reproducing kernel Hilbert space⁶ H^2 , each model space \mathcal{K}_u itself possesses a reproducing kernel. In this section, we identify these kernels and explore their basic properties.

5.1. Basic properties. Recalling that the Cauchy kernels $c_\lambda = (1 - \bar{\lambda}z)^{-1}$ are the reproducing kernels for the Hardy space (see Subsection 2.3), let us attempt to compute the corresponding reproducing kernels for \mathcal{K}_u . We first observe that if $f = uh$ belongs to uH^2 , then

$$f(\lambda) = u(\lambda)h(\lambda) = u(\lambda)\langle h, c_\lambda \rangle = u(\lambda)\langle f\bar{u}, c_\lambda \rangle = \langle f, \overline{u(\lambda)}uc_\lambda \rangle,$$

from which it follows that the reproducing kernel for uH^2 is given by

$$\overline{u(\lambda)}u(z)c_\lambda(z).$$

If f belongs to \mathcal{K}_u , then it follows that

$$\begin{aligned} f(\lambda) &= \langle f, c_\lambda \rangle \\ &= \langle f, c_\lambda \rangle - u(\lambda)\langle f, uc_\lambda \rangle \\ &= \langle f, (1 - \overline{u(\lambda)}u)c_\lambda \rangle. \end{aligned}$$

Moreover, the function $(1 - \overline{u(\lambda)}u)c_\lambda$ belongs to \mathcal{K}_u since

$$\begin{aligned} \langle uh, (1 - \overline{u(\lambda)}u)c_\lambda \rangle &= u(\lambda)h(\lambda) - u(\lambda)\langle uh, uc_\lambda \rangle \\ &= u(\lambda)h(\lambda) - u(\lambda)\langle h, c_\lambda \rangle \\ &= u(\lambda)h(\lambda) - u(\lambda)h(\lambda) \\ &= 0 \end{aligned}$$

for all h in H^2 . Putting this all together we see that

$$f(\lambda) = \langle f, k_\lambda \rangle, \quad (f \in \mathcal{K}_u)$$

⁶A Hilbert space \mathcal{H} of analytic functions on a domain $\Omega \subset \mathbb{C}$ is called a *reproducing kernel Hilbert space* if for each $\lambda \in \Omega$ there is a function $K_\lambda \in \mathcal{H}$ for which $f(\lambda) = \langle f, K_\lambda \rangle_{\mathcal{H}}$ for all $f \in \mathcal{H}$. All of the classical Hilbert spaces of analytic functions on the disk (e.g., Hardy, Bergman, Dirichlet, etc.) are reproducing kernel Hilbert spaces and understanding the properties of the kernels often yields valuable information about the functions in \mathcal{H} . A few good sources for this are [2, 12, 70].

where

$$k_\lambda(z) = \frac{1 - \overline{u(\lambda)}u(z)}{1 - \overline{\lambda}z}. \quad (5.1)$$

The function k_λ is called the *reproducing kernel* for \mathcal{K}_u . When dealing with more than one model space at a time, the notation k_λ^u is often used to denote the reproducing kernel for \mathcal{K}_u .

From the general theory of reproducing kernels [2, 12, 70], it follows that if e_1, e_2, \dots is any orthonormal basis for \mathcal{K}_u , then

$$k_\lambda(z) = \sum_{n \geq 1} \overline{e_n(\lambda)} e_n(z), \quad (5.2)$$

the sum converging in the norm of \mathcal{K}_u [2]. The following example illustrates both expressions (5.1) and (5.2).

Example 5.3. If $u = z^n$, then $\mathcal{K}_u = \text{span}\{1, z, z^2, \dots, z^{n-1}\}$ and

$$k_\lambda(z) = \frac{1 - \overline{\lambda}^n z^n}{1 - \overline{\lambda}z} = 1 + \overline{\lambda}z + \overline{\lambda}^2 z^2 + \dots + \overline{\lambda}^{n-1} z^{n-1}.$$

We are now in a position to provide an elegant derivation of the decomposition (3.19). Indeed, for inner functions u and v , divide the trivial equality

$$1 - \overline{u(\lambda)v(\lambda)}u(z)v(z) = (1 - \overline{u(\lambda)}u(z)) + \overline{u(\lambda)}u(z)(1 - \overline{v(\lambda)}v(z))$$

by $1 - \overline{\lambda}z$ to get the identity

$$k_\lambda^{uv} = k_\lambda^u + \overline{u(\lambda)}u(z)k_\lambda^v.$$

However, the preceding is simply a restatement, in terms of reproducing kernels, of the fact that $\mathcal{K}_{uv} = \mathcal{K}_u \oplus u\mathcal{K}_v$.

Since

$$|k_\lambda(z)| \leq \frac{2}{1 - |\lambda|} \quad (5.4)$$

for each z in \mathbb{D} , it follows that each k_λ belongs to $\mathcal{K}_u \cap H^\infty$. Moreover, the kernel functions k_λ are among the few readily identifiable functions that belong to \mathcal{K}_u . As such, they provide invaluable insight into the structure and properties of model spaces. In fact, kernel functions often wind up being “test functions” for various statements about model spaces. As an example of what we mean by this, suppose that the quantity

$$\|k_\lambda\|^2 = \frac{1 - |u(\lambda)|^2}{1 - |\lambda|^2}$$

remains bounded as $\lambda \rightarrow \zeta \in \mathbb{T}$ along some path $\Gamma \subset \mathbb{D}$, then since

$$|f(\lambda)| = |\langle f, k_\lambda \rangle| \leq \|f\| \|k_\lambda\|,$$

we see that every f in \mathcal{K}_u is bounded along Γ . We will see more of these types of results in Section 6.

The orthogonal projection

$$P_u : L^2 \rightarrow \mathcal{K}_u \quad (5.5)$$

arises frequently in the study of Hankel operators, model operators and truncated Toeplitz operators. This important operator can be expressed in a simple manner using reproducing kernels.

Proposition 5.6. *For each f in L^2 and λ in \mathbb{D} ,*

$$(P_u f)(\lambda) = \langle f, k_\lambda \rangle. \quad (5.7)$$

Proof. Using the selfadjointness of P_u we get

$$\langle f, k_\lambda \rangle = \langle f, P_u k_\lambda \rangle = \langle P_u f, k_\lambda \rangle = (P_u f)(\lambda). \quad \square$$

Rewriting (5.7) as an integral we obtain

$$(P_u f)(\lambda) = \int_{\mathbb{T}} f(\zeta) \frac{1 - u(\lambda)\overline{u(\zeta)}}{1 - \lambda\bar{\zeta}} dm(\zeta),$$

which can be used to estimate or control the behavior of $P_u f$. In particular, the preceding formula highlights the dependence of $P_u f$ on the behavior of u itself.

5.2. Density results. As mentioned earlier, in general, the model spaces \mathcal{K}_u contain few readily identifiable functions. One therefore relies heavily upon the kernel functions k_λ in the study of model spaces since they are among the few explicitly describable functions contained in \mathcal{K}_u . As the following proposition shows, one can always find a collection of kernels whose span is dense in the whole space.

Proposition 5.8. *If Λ is a subset of \mathbb{D} such that either (i) Λ has an accumulation point in \mathbb{D} , or (ii) $\sum_{\lambda \in \Lambda} (1 - |\lambda|)$ diverges, then for any inner function u*

$$\bigvee \{k_\lambda : \lambda \in \Lambda\} = \mathcal{K}_u.$$

Proof. The containment \subseteq is obvious. If $f \perp k_\lambda$ for all $\lambda \in \Lambda$, then f vanishes on Λ . (i) If Λ has an accumulation point in \mathbb{D} , then the Identity Theorem implies that $f \equiv 0$. (ii) If $\sum_{\lambda \in \Lambda} (1 - |\lambda|)$ diverges, then $f \equiv 0$ since the zero set of a nonzero H^2 function must satisfy the Blaschke condition (2.7). \square

A somewhat more general result is provided by [44, Thm. 1]:

Proposition 5.9. *If u is a nonconstant inner function, then there exists a subset $E \subset \mathbb{D}$ of area measure zero such that for each w in $\mathbb{D} \setminus E$, the inverse image $u^{-1}(\{w\})$ is nonempty, $u'(\lambda) \neq 0$ for all λ in $u^{-1}(\{w\})$, and*

$$\bigvee \{k_\lambda : \lambda \in u^{-1}(\{w\})\} = \mathcal{K}_u.$$

Remark 5.10. We will take up the general question of whether a sequence of kernel functions has a dense linear span in \mathcal{K}_u when we discuss completeness problems in Section 9.

Since each kernel function k_λ belongs to H^∞ by (5.4), an immediate consequence of either of the preceding propositions is the following useful result.

Proposition 5.11. *The linear manifold $\mathcal{K}_u \cap H^\infty$ is dense in \mathcal{K}_u .*

A second proof can be obtained by noting that

$$\mathcal{K}_u = \bigvee \{S^{*n}u : n \geq 1\}$$

and that each of the backward shifts $S^{*n}u$ of u belongs to H^∞ . In certain sophisticated applications one requires a dense subset of \mathcal{K}_u consisting of functions having a certain degree of smoothness on \mathbb{D}^- . This next result is a restatement of Proposition 3.15 for infinite Blaschke products.

Proposition 5.12. *If u is a Blaschke product with simple zeros $\lambda_1, \lambda_2, \dots$, then*

$$\bigvee \{k_{\lambda_1}, k_{\lambda_2}, \dots\} = \mathcal{K}_u.$$

Thus \mathcal{K}_u contains a dense subset whose elements are continuous on \mathbb{D}^- and whose boundary functions are infinitely differentiable on \mathbb{T} .

Proof. Indeed, suppose that f is a function in \mathcal{K}_u that satisfies $\langle f, k_{\lambda_n} \rangle = 0$ for all n . This implies that $f(\lambda_n) = 0$ for all n , whence $u|f$ so that f belongs to uH^2 . In other words, f must be identically zero. Since

$$k_{\lambda_n}(z) = \frac{1 - \overline{u(\lambda_n)}u(z)}{1 - \overline{\lambda_n}z} = \frac{1}{1 - \overline{\lambda_n}z} = c_{\lambda_n}(z), \quad (5.13)$$

the second statement follows immediately. \square

If the zeros of u are not simple, then the preceding statement is true if one also includes the appropriate derivatives of the kernel functions in the spanning set. When u is not a Blaschke product, finding a dense set of functions in \mathcal{K}_u , each of which is continuous on \mathbb{D}^- , is much more difficult. A deep result in this direction is due to A.B. Aleksandrov [7], who proved the following astonishing result (see [26] for a discussion of this).

Theorem 5.14 (Aleksandrov). *For an inner function u , the set $\mathcal{K}_u \cap \mathcal{A}$ is dense in \mathcal{K}_u . Here \mathcal{A} denotes the disk algebra, the Banach algebra of all H^∞ functions that are continuous on \mathbb{D}^- .*

This theorem is remarkable due to the fact that \mathcal{K}_u often does not contain a single readily identifiable function that is continuous on \mathbb{D}^- . For example, if u is the singular inner function

$$u(z) = \exp\left(\frac{z+1}{z-1}\right),$$

then it is not at all obvious that \mathcal{K}_u contains any functions that are continuous on \mathbb{D}^- , let alone a dense set of them. See [34] for some related results concerning when \mathcal{K}_u contains smoother functions than those in \mathcal{A} .

5.3. Cauchy-Szegő bases. If u is an inner function and $u(\lambda) = 0$, then (5.13) tells us that $k_\lambda(z) = c_\lambda(z) = (1 - \bar{\lambda}z)^{-1}$. Thus the kernel functions corresponding to zeros of u are extremely simple functions to work with since they do not depend explicitly upon u . In certain situations, these functions can be used to construct bases for model spaces (in a sense to be made precise shortly). However, no two Cauchy kernels are orthogonal, so one cannot hope to obtain an orthonormal basis of kernel functions.⁷ We therefore need a somewhat more flexible definition [25].

Definition 5.15. A linearly independent sequence \mathbf{x}_n in a Hilbert space \mathcal{H} is called a *Riesz basis* for \mathcal{H} if $\bigvee\{\mathbf{x}_1, \mathbf{x}_2, \dots\} = \mathcal{H}$ and there exist constants $M_1, M_2 > 0$ such that

$$M_1 \sum_{i=1}^n |a_i|^2 \leq \left\| \sum_{i=1}^n a_i \mathbf{x}_i \right\|^2 \leq M_2 \sum_{i=1}^n |a_i|^2$$

for all finite numerical sequences a_1, a_2, \dots, a_n .

The following seminal result of L. Carleson tells us when the normalized reproducing kernels corresponding to the zero set of a Blaschke product forms a Riesz basis for \mathcal{K}_u [67, p. 133]

Theorem 5.16. *If u is a Blaschke product with simple zeros $\lambda_1, \lambda_2, \dots$, then the normalized kernels*

$$\frac{k_{\lambda_n}}{\|k_{\lambda_n}\|} = \frac{\sqrt{1 - |\lambda_n|^2}}{1 - \bar{\lambda}_n z}$$

form a Riesz basis for \mathcal{K}_u if and only if there exists a $\delta > 0$ such that

$$\delta < \prod_{\substack{j=1 \\ j \neq i}}^{\infty} \left| \frac{\lambda_i - \lambda_j}{1 - \bar{\lambda}_j \lambda_i} \right|. \quad (i = 1, 2, \dots)$$

There is a generalization of this result where one can somewhat relax the condition that the λ_n are the zeros of u [58].

5.4. Takenaka-Malmquist-Walsh bases. Unlike the Hardy space H^2 itself, the model spaces \mathcal{K}_u do not come pre-equipped with a canonical orthonormal basis. It turns out that an orthonormal basis for \mathcal{K}_u , where u is a Blaschke product, can be obtained by orthogonalizing the kernel functions corresponding to the zeros of u .

For w in \mathbb{D} , we let

$$b_w(z) = \frac{z - w}{1 - \bar{w}z}. \quad (5.17)$$

If u is a Blaschke product with simple zeros $\lambda_1, \lambda_2, \dots$, then observe that

$$\langle b_{\lambda_1} k_{\lambda_2}, k_{\lambda_1} \rangle = b_{\lambda_1}(\lambda_1) k_{\lambda_2}(\lambda_1) = 0.$$

⁷One can sometimes obtain orthonormal bases consisting of *boundary kernels* (see Section 8).

Similarly, we have

$$\langle b_{\lambda_1} b_{\lambda_2} k_{\lambda_3}, k_{\lambda_1} \rangle = \langle b_{\lambda_1} b_{\lambda_2} k_{\lambda_3}, k_{\lambda_2} \rangle = 0. \quad (5.18)$$

This process suggests the following iterative definition: If u is a Blaschke product with zeros $\lambda_1, \lambda_2, \dots$, let

$$v_1(z) := \frac{\sqrt{1 - |\lambda_1|^2}}{1 - \overline{\lambda_1}z},$$

$$v_k(z) = \left(\prod_{i=1}^{k-1} b_{\lambda_i} \right) \frac{\sqrt{1 - |\lambda_k|^2}}{1 - \overline{\lambda_k}z}. \quad (k \geq 2)$$

One can show that $\{v_n : n \geq 1\}$ is an orthonormal basis for \mathcal{K}_u that is the Gram-Schmidt orthonormalization of the kernels $k_{\lambda_1}, k_{\lambda_2}, \dots$.

The terminology here is not completely standard. It seems that these bases first appeared in Takenaka's 1925 paper involving finite Blaschke products [94]. The classic text [67] considers the general case where u is a potentially infinite Blaschke product, referring to this result as the Malmquist-Walsh Lemma. In light of Takenaka's early contribution to the subject, the authors chose in [49] to refer to such a basis for \mathcal{K}_u as a *Takenaka-Malmquist-Walsh (TMW) basis*.

Another important family of orthonormal bases of \mathcal{K}_u is (sometimes) provided by the Aleksandrov-Clark spectral theory of rank-one unitary perturbations of model operators. The so-called *modified Aleksandrov-Clark bases* are particularly important in the study of finite-dimensional model spaces. To discuss these bases, we first require a few words about boundary behavior and angular derivatives.

6. BOUNDARY BEHAVIOR

6.1. Pseudocontinuations. Functions in model spaces enjoy certain "continuation" properties across \mathbb{T} . These types of continuation properties appear in many places [76] and in many different settings but they all go under the broad heading of "generalized analytic continuation." The type of generalized analytic continuation that is relevant to model spaces is called *pseudocontinuation* and it was first explored by H. S. Shapiro [85, 86]. In what follows, we let

$$\mathbb{D}_e := \{|z| > 1\} \cup \{\infty\}$$

denote the *extended exterior disk*, the complement of the closed unit disk in the extended complex plane.

Definition 6.1. Let f and \tilde{f} be meromorphic functions on \mathbb{D} and \mathbb{D}_e , respectively. If the nontangential limiting values of f (from \mathbb{D}) agree with the nontangential limiting values of \tilde{f} (from \mathbb{D}_e) almost everywhere on \mathbb{T} , then we say that f and \tilde{f} are *pseudocontinuations* of one another.

Remark 6.2. Pseudocontinuations are unique in the sense that if F and G are meromorphic functions on \mathbb{D}_e that are both pseudocontinuations of f , then $F = G$. This follows from the Privalov uniqueness theorem [59, p. 62], which states that if f and g are meromorphic on \mathbb{D} with equal nontangential limits on any set of positive Lebesgue measure on \mathbb{T} , then $f = g$. This is why the definition of pseudocontinuation is stated in terms of nontangential limits as opposed to radial limits, where Privalov's theorem is no longer true. However, in the context we will apply this definition, all the functions involved will be of *bounded type*, that is to say the quotient of two bounded analytic functions (recall Definition 2.18), where the nontangential limits exist almost everywhere.

By Privalov's uniqueness theorem again, we can show that pseudocontinuation is compatible with analytic continuation in that if f (on \mathbb{D}) has a pseudocontinuation F (on \mathbb{D}_e) and f also has an analytic continuation F_1 across some neighborhood of a point on \mathbb{T} , then $F_1 = F$.

Example 6.3. Let us provide a few instructive examples (see [76] for further details).

- (i) Any inner function u has a pseudocontinuation to \mathbb{D}_e defined by

$$\tilde{u}(z) := \frac{1}{u(1/\bar{z})}. \quad (6.4)$$

In fact, this pseudocontinuation is one of bounded type (written PCBT), being a quotient of two bounded analytic functions on \mathbb{D}_e . Pseudocontinuations of bounded type will play an important role momentarily.

- (ii) If f is a rational function whose poles lie in \mathbb{D}_e , then f is PCBT.
 (iii) Since pseudocontinuations must be compatible with analytic continuations (see the previous remark), functions with isolated branch points on \mathbb{T} , such as $\sqrt{1-z}$, do not possess pseudocontinuations.
 (iv) The function $f(z) = \exp z$ is not pseudocontinuable. Although it is analytically continuable to \mathbb{C} , it is not meromorphic on \mathbb{D}_e due to its essential singularity at ∞ .
 (v) The classical gap theorems of Hadamard [53] and Fabry [37] show that analytic functions on \mathbb{D} with lacunary power series do not have analytic continuations across any point of the unit circle. Such functions do not have pseudocontinuations either [1, 8, 86].

The following seminal result demonstrates the direct connection between membership in a model space and pseudocontinuity. In what follows,

$$H^2(\mathbb{D}_e) := \{f(1/z) : f \in H^2\} \quad (6.5)$$

denotes the Hardy space of the extended exterior disk.

Theorem 6.6 (Douglas, Shapiro, Shields [31]). *A function $f \in H^2$ belongs to \mathcal{K}_u if and only if f/u has a pseudocontinuation $F_u \in H^2(\mathbb{D}_e)$ satisfying $F_u(\infty) = 0$.*

Proof. Suppose that f belongs to \mathcal{K}_u . By Proposition 3.3, $f = \overline{g\zeta}u$ almost everywhere on \mathbb{T} . Define the function F_u on \mathbb{D}_e by

$$F_u(z) := \frac{1}{z} \overline{g\left(\frac{1}{\bar{z}}\right)} \quad (6.7)$$

and note by (6.5) that F_u is in $H^2(\mathbb{D}_e)$ and $F_u(\infty) = 0$. Moreover, by the identity $f = \overline{g\zeta}u$ almost everywhere on \mathbb{T} we see that $F_u(\zeta) = f(\zeta)/u(\zeta)$ for almost every $\zeta \in \mathbb{T}$, i.e., F_u is a pseudocontinuation of f/u . This proves one direction.

For the other direction, suppose that f/u has a pseudocontinuation $F_u \in H^2(\mathbb{D}_e)$ with $F_u(\infty) = 0$. Then, again, by (6.5) we have

$$F_u(z) = \frac{1}{z} h\left(\frac{1}{z}\right)$$

for some $h \in H^2$ and $f(\zeta)/u(\zeta) = F_u(\zeta)$ for almost every ζ in \mathbb{T} . Define a function g on \mathbb{D} by $g(z) := \overline{h(\bar{z})}$ and observe that $g \in H^2$. From here we see that $f(\zeta) = u(\zeta)\zeta\overline{g(\bar{\zeta})}$ for almost every ζ in \mathbb{T} from which it follows, via Proposition 3.3, that f belongs to \mathcal{K}_u . \square

With a little more work, one can also determine the *cyclic vectors* for S^* , those f in H^2 for which

$$\bigvee \{S^{*n}f : n = 0, 1, 2, \dots\} = H^2.$$

Corollary 6.8. *A function f in H^2 is not cyclic for S^* if and only if f is PCBT.*

Proof. We will prove only one direction. Indeed, suppose f is non-cyclic for S^* . Then f must belong to some model space \mathcal{K}_u . By the previous theorem, f/u has a pseudocontinuation F_u that belongs to $H^2(\mathbb{D}_e)$. But since u already has a natural pseudocontinuation \tilde{u} given by (6.4), we see that f has a pseudocontinuation $\tilde{u}F_u$ that is of bounded type⁸. \square

Along with Example 6.3 this corollary shows that inner functions and rational function are noncyclic for S^* whereas functions like $\sqrt{1-z}$ and e^z are cyclic.

In principle the cyclic vector problem for S^* is solved (i.e., f is noncyclic for S^* if and only if f is PCBT). However, this solution is not as explicit as the solution to the cyclic vector problem for S (i.e., f is cyclic for S if and only if f is outer). Outer functions are, in a sense, readily identifiable. On the other hand, PCBT functions are not so easily recognized. We refer the reader to some papers which partially characterize the noncyclic vectors

⁸We are using the well-known fact here that F_u (or any function in $H^2(\mathbb{D}_e)$) is of bounded type (see Remark 2.19).

for S^* by means of growth of Taylor coefficients [60,84], the modulus of the function [31,60], and gaps in the Taylor coefficients [1,8,31].

6.2. Analytic continuation. Recall from (4.4) and Theorem 4.5 that if $u = B_\Lambda S_\mu$, where B_Λ is a Blaschke product with zero set Λ and S_μ is a singular inner function with corresponding singular measure μ , then the *spectrum* of u is the set

$$\sigma(u) = \Lambda^- \cup \text{supp } \mu.$$

The relevance of $\sigma(u)$ to the function theoretic properties of \mathcal{K}_u lies in the following observation [67, p. 65], [27, p. 84].

Proposition 6.9. *Every f in \mathcal{K}_u can be analytically continued to*

$$\widehat{\mathbb{C}} \setminus \{1/\bar{z} : z \in \sigma(u)\},$$

where $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ denotes the extended complex plane.

Proof. A sketch of the proof goes as follows. By [52, Sect. 2.6] u has an analytic continuation across $\mathbb{T} \setminus \sigma(u)$. By Theorem 6.6, f/u has a pseudo-continuation F_u in $H^2(\mathbb{D}_e)$ for any $f \in \mathcal{K}_u$. Let J be an open arc contained in $\mathbb{T} \setminus \sigma(u)$. We know that u has an analytic continuation across J . We also know that the integral means

$$\int_J \left| \frac{f}{u}(r\zeta) \right| dm(\zeta)$$

and

$$\int_J |F_u(s\zeta)| dm(\zeta)$$

are uniformly bounded in $0 < r < 1$ and $s > 1$. One can now take a triangle Δ cut by J and show (by cutting Δ with $(1-\epsilon)J$ and $(1+\epsilon)J$ and using the boundedness of the above integral means) that the contour integral of the function defined by f/u on \mathbb{D} and F_u on \mathbb{D}_e integrates to zero on Δ . Now apply Morera's theorem⁹. \square

6.3. Nontangential limits. Functions in H^2 possess nontangential limits almost everywhere on \mathbb{T} . However, one can easily see that for each fixed ζ in \mathbb{T} there exists an H^2 function that does not have a finite nontangential limit at ζ (e.g., $f(z) = \log(\zeta - z)$). In sharp contrast to this, for a given model space \mathcal{K}_u , there might exist points ζ in \mathbb{T} such that each function in \mathcal{K}_u possesses a nontangential limit at ζ . We begin with a definition.

Definition 6.10. If u is inner and $\zeta \in \mathbb{T}$, then u has an *angular derivative in the sense of Carathéodory* (ADC) at ζ if the nontangential limits of u and u' exist at ζ and $|u(\zeta)| = 1$.

⁹There is a precise version of this type of Morera's theorem found in [52].

A number of equivalent conditions for the existence of an angular derivative (called the Julia-Carathéodory theorem) can be found in [87, p. 57]. The precise relationship between angular derivatives and model spaces is contained in the following important result.

Theorem 6.11 (Ahern-Clark [3,4]). *For an inner function $u = B_\Lambda S_\mu$, where B_Λ is a Blaschke product with zeros $\Lambda = \{\lambda_n\}_{n=1}^\infty$, repeated according to multiplicity, S_μ is a singular inner function with corresponding singular measure μ , and $\zeta \in \mathbb{T}$, the following are equivalent:*

- (i) Every $f \in \mathcal{K}_u$ has a nontangential limit at ζ .
- (ii) For every $f \in \mathcal{K}_u$, $f(\lambda)$ is bounded as $\lambda \rightarrow \zeta$ nontangentially.
- (iii) u has an ADC at ζ .
- (iv) $k_\zeta(z) = \frac{1 - \overline{u(\zeta)}u(z)}{1 - \bar{\zeta}z} \in H^2$,
- (v) $\sum_{n \geq 1} \frac{1 - |\lambda_n|^2}{|\zeta - \lambda_n|^2} + \int_{\mathbb{T}} \frac{d\mu(\xi)}{|\xi - \zeta|^2} < \infty$.

It is worth comparing condition (v) of the preceding theorem to an old result of Frostman [38] that says that an inner function u (and all its divisors) has a non-tangential limits of modulus one at ζ in \mathbb{T} whenever

$$\sum_{n \geq 1} \frac{1 - |\lambda_n|}{|\zeta - \lambda_n|} + \int_{\mathbb{T}} \frac{d\mu(\xi)}{|\xi - \zeta|} < \infty.$$

We should also mention that there are inner functions (in fact Blaschke products) u for which $|u'(\zeta)| = \infty$ at every point of \mathbb{T} [38].

There are also results about *tangential* limits of functions from model spaces [15,20]. When condition (v) fails then there are functions in \mathcal{K}_u that do not have nontangential limits at ζ . In particular, there are functions f in \mathcal{K}_u for which $|f(r\zeta)|$ is unbounded as $r \rightarrow 1^-$. However, in [54] there are results that give bounds (both upper and lower) on the growth of $|f(r\zeta)|$ as a function of r .

7. CONJUGATION

Each model space \mathcal{K}_u comes equipped with a conjugation, a certain type of conjugate-linear operator that generalizes complex conjugation on \mathbb{C} . Not only does this conjugation cast a new light on pseudocontinuations, it also interacts with a number of important linear operators that act upon model spaces. Most of the material in this section can be found in [39,46].

7.1. Basic properties. We say that a conjugate-linear function $C : \mathcal{H} \rightarrow \mathcal{H}$ on a complex Hilbert space \mathcal{H} is a *conjugation* if C is isometric (i.e., $\|Cx\| =$

$\|x\|$ for all x in \mathcal{H}) and involutive (i.e., $C^2 = I$). In light of the polarization identity, it turns out that the isometric condition is equivalent to the assertion that $\langle Cx, Cy \rangle = \langle y, x \rangle$ for all x, y in \mathcal{H} .

The structure of conjugations is remarkably simple. If C is a conjugation on \mathcal{H} , then there exists an orthonormal basis $\{e_n\}$ of \mathcal{H} such that $Ce_n = e_n$ for all n [40, Lem. 1]. We refer to such a basis as a *C-real* orthonormal basis of \mathcal{H} . As a consequence, any conjugation is unitarily equivalent to the *canonical conjugation*

$$(z_1, z_2, \dots) \mapsto (\overline{z_1}, \overline{z_2}, \dots)$$

on an ℓ^2 -space of the appropriate dimension.

In order to understand the natural conjugation on \mathcal{K}_u , we must first remind the reader that $\mathcal{K}_u = H^2 \cap u\overline{z}H^2$, as a space of functions on \mathbb{T} (see Proposition 3.3).

Proposition 7.1. *The conjugate-linear operator $C : \mathcal{K}_u \rightarrow \mathcal{K}_u$, defined in terms of boundary functions on \mathbb{T} by $Cf = \overline{fzu}$, is a conjugation. In particular, $|f| = |Cf|$ almost everywhere on \mathbb{T} so that f and Cf share the same outer factor.*

Proof. Since $|u| = 1$ almost everywhere on \mathbb{T} , it follows that C is conjugate-linear, isometric, and involutive. We need only prove that C maps \mathcal{K}_u into \mathcal{K}_u . Since f is orthogonal to uH^2 , it follows that

$$\begin{aligned} \langle Cf, \overline{zh} \rangle &= \langle \overline{fzu}, \overline{zh} \rangle \\ &= \int_{\mathbb{T}} \overline{f(\zeta)} \zeta u(\zeta) \zeta h(\zeta) dm(\zeta) \\ &= \int_{\mathbb{T}} u(\zeta) h(\zeta) \overline{f(\zeta)} dm(\zeta) \\ &= \langle uh, f \rangle \\ &= 0 \end{aligned}$$

for all h in H^2 . In other words, Cf belongs to H^2 . Similarly,

$$\langle Cf, uh \rangle = \langle \overline{fzu}, uh \rangle = \langle \overline{fz}, h \rangle = 0,$$

from which it follows that Cf belongs to \mathcal{K}_u . \square

Example 7.2. If $u(z) = z^n$, then $\mathcal{K}_u = \bigvee \{1, z, \dots, z^{n-1}\}$ and the conjugation C assumes the form

$$C(a_0 + a_1z + \dots + a_{n-1}z^{n-1}) = \overline{a_{n-1}} + \overline{a_{n-2}}z + \dots + \overline{a_0}z^{n-1}.$$

Now suppose that u is a finite Blaschke product having zeros $\lambda_1, \lambda_2, \dots, \lambda_n$, repeated according to multiplicity. Referring back to Proposition 3.15 for an explicit description of \mathcal{K}_u , one can show that

$$C \left(\frac{a_0 + a_1z + \dots + a_{n-1}z^{n-1}}{\prod_{i=1}^n (1 - \overline{\lambda_i}z)} \right) = \frac{\overline{a_{n-1}} + \overline{a_{n-2}}z + \dots + \overline{a_0}z^{n-1}}{\prod_{i=1}^n (1 - \overline{\lambda_i}z)}.$$

In other words, each element of \mathcal{K}_u can be represented as a rational function whose denominator is $\prod_{i=1}^n (1 - \overline{\lambda_i}z)$ and whose numerator is a polynomial

of degree $\leq n - 1$. The conjugation C acts on \mathcal{K}_u by reversing the order of the polynomial in the numerator and conjugating its coefficients.

Example 7.3. A straightforward computation reveals that C sends reproducing kernels to difference quotients:

$$\begin{aligned} [Ck_\lambda](z) &= \overline{\left(\frac{1 - \overline{u(\lambda)}u(z)}{1 - \overline{\lambda}z} \right)} \overline{z}u(z) \\ &= \frac{1 - u(\lambda)\overline{u(z)}}{1 - \lambda\overline{z}} \cdot \frac{u(z)}{z} \\ &= \frac{u(z) - u(\lambda)}{z - \lambda}, \end{aligned}$$

where, as usual, we consider all of the functions involved as functions on \mathbb{T} (i.e., so that $z\overline{z} = 1$). For each λ , the kernel function k_λ is an outer function since it is the quotient of the two outer functions $1 - \overline{u(\lambda)}u(z)$ and $1 - \overline{\lambda}z$ [33, Ex. 1, Ch. 3]. In light of Proposition 7.1, we expect that the difference quotient Ck_λ is simply an inner multiple of k_λ . This is indeed the case as the following computation shows.

$$\begin{aligned} [Ck_\lambda](z) &= \frac{u(z) - u(\lambda)}{z - \lambda} \\ &= \frac{u(z) - u(\lambda)}{1 - \overline{u(\lambda)}u(z)} \cdot \frac{1 - \overline{\lambda}z}{z - \lambda} \cdot \frac{1 - \overline{u(\lambda)}u(z)}{1 - \overline{\lambda}z} \\ &= \frac{b_{u(\lambda)}(u(z))}{b_\lambda(z)} k_\lambda(z), \end{aligned}$$

where $b_w(z) = (z - w)/(1 - \overline{w}z)$ is the disk automorphism (5.17).

Example 7.4. If ζ is a point on \mathbb{T} for which u has an ADC (see Subsection 6.3), then the boundary kernel k_ζ , as defined in Theorem 6.11, belongs to \mathcal{K}_u . In this case, something remarkable occurs. Since ζ and $u(\zeta)$ are of unit modulus, it follows that

$$\begin{aligned} [Ck_\zeta](z) &= \frac{u(z) - u(\zeta)}{z - \zeta} \\ &= \frac{u(\zeta)}{\zeta} \cdot \frac{1 - \overline{u(\zeta)}u(z)}{1 - \overline{\zeta}z} \\ &= \overline{\zeta}u(\zeta)u_\zeta(z). \end{aligned}$$

For either branch of the square root, it follows that the outer function

$$(\overline{\zeta}u(\zeta))^{\frac{1}{2}} k_\zeta$$

belongs to \mathcal{K}_u and is fixed by C .

Under certain circumstances, it is possible to construct C -real orthonormal bases for \mathcal{K}_u using boundary kernels. This is relatively straightforward

for finite dimensional model spaces, as the following example demonstrates (see Section 8 for a discussion of the general setting).

Example 7.5. Let u be a finite Blaschke product with n zeros repeated according to multiplicity. For the sake of simplicity, suppose that $u(0) = 0$. For each fixed α in \mathbb{T} , the equation $u(\zeta) = \alpha$ has precisely n distinct solutions $\zeta_1, \zeta_2, \dots, \zeta_n$ on \mathbb{T} . The functions

$$e_j(z) = \frac{e^{\frac{i}{2}(\arg \alpha - \arg \zeta_j)} (1 - \overline{\alpha}u(z))}{\sqrt{|u'(\zeta_j)|} (1 - \overline{\zeta_j}z)}$$

for $j = 1, 2, \dots, n$ form a C -real orthonormal basis of \mathcal{K}_u .

7.2. Associated inner functions. Fix an inner function u and let $f = I_1F$ denote the inner-outer factorization of a nonzero function f in \mathcal{K}_u . Proposition 7.1 ensures that $g = Cf$ has the same outer factor as f whence we may write $g = I_2F$ for some inner function I_2 . Since $g = \overline{f}zu$ almost everywhere on \mathbb{T} it follows that $I_2F = \overline{I_1F}zu$ whence

$$I_1I_2 = \frac{\overline{F}zu}{F}. \quad (7.6)$$

The inner function $\mathcal{I}_F = I_1I_2$ is called the *associated inner function* of F with respect to u [46].¹⁰ In light of (7.6), we see that \mathcal{I}_F is uniquely determined by u and F .

On the other hand, if I_1 and I_2 are inner functions such that $I_1I_2 = \mathcal{I}_F$, then (7.6) implies that

$$I_1F = \overline{I_2F}zu$$

so that the functions $f = I_1F$ and $g = I_2F$ satisfy $f = \overline{g}zu$ almost everywhere on \mathbb{T} . By Proposition 3.3, it follows that f and g belong to \mathcal{K}_u and satisfy $Cf = g$. Putting this all together, we obtain the following result.

Proposition 7.7. *The set of all functions in \mathcal{K}_u having outer factor F is precisely*

$$\{\theta F : \theta \text{ inner and } \theta|_{\mathcal{I}_F}\}. \quad (7.8)$$

We define a partial ordering on (7.8) by declaring that $\theta_1F \leq \theta_2F$ if and only if $\theta_1|_{\mathcal{I}_F} \leq \theta_2|_{\mathcal{I}_F}$. With respect to this ordering, F and \mathcal{I}_FF are the unique minimal and maximal elements, respectively. Moreover, C restricts to an order-reversing bijection from (7.8) to itself.

7.3. Generators of \mathcal{K}_u . We say that a function f in \mathcal{K}_u *generates* \mathcal{K}_u if

$$\bigvee \{f, S^*f, S^{*2}f, \dots\} = \mathcal{K}_u. \quad (7.9)$$

Recall that Proposition 3.8 tells us that if $f = \overline{g}zu$ belongs to \mathcal{K}_u , then (7.9) holds if and only if u and the inner factor of g are relatively prime. Recognizing that $g = Cf$ in this setting immediately yields [46, Prop. 4.3].

¹⁰We should note that the influential article [31, Rem. 3.1.6] of Douglas, Shapiro, and Shields refers to u itself as the associated inner function of f .

Proposition 7.10. *If f belongs to \mathcal{K}_u and the inner factor of Cf is relatively prime to u , then f generates \mathcal{K}_u .*

Example 7.11. Each difference quotient $Ck_\lambda = (u(z) - u(\lambda))/(z - \lambda)$ generates \mathcal{K}_u since its conjugate $C(Ck_\lambda) = k_\lambda$ is outer. If u is a singular inner function (i.e., u is not divisible by any Blaschke product), then Frostman's Theorem [33, Ex. 8, Ch. 2], [38] asserts that k_λ generates \mathcal{K}_u for almost every (with respect to area measure) λ in \mathbb{D} . Indeed, Example 7.3 tells us that the inner factor of Ck_λ is precisely $b_{u(\lambda)}(u(z))/b_\lambda(z)$, which is a Blaschke product whenever $u(\lambda)$ is nonzero and does not lie in the exceptional set for u .

Corollary 7.12. *If f is an outer function in \mathcal{K}_u , then Cf generates \mathcal{K}_u . In particular, any self-conjugate outer function in \mathcal{K}_u generates \mathcal{K}_u .*

Example 7.13. Any boundary kernel k_ζ in \mathcal{K}_u generates \mathcal{K}_u . Indeed, Example 7.4 shows that any such function is a constant multiple of a self-conjugate outer function.

7.4. Quaternionic structure of 2×2 inner functions. It at least one instance, it turns out that conjugations on model spaces can yield structural insights into the nature of higher order inner functions. The following result can be found in [47], although it is stated in [40] without proof, along with several other related results.

Theorem 7.14. *If u is a nonconstant inner function, then the function $\Theta : \mathbb{D} \rightarrow M_2(\mathbb{C})$ defined by*

$$\Theta = \begin{bmatrix} a & -b \\ Cb & Ca \end{bmatrix} \tag{7.15}$$

is unitary almost everywhere on \mathbb{T} and satisfies $\det \Theta = u$ if and only if

- (i) a, b, c, d belong to \mathcal{K}_{zu} .
- (ii) $|a|^2 + |b|^2 = 1$ almost everywhere on \mathbb{T} .
- (iii) $Ca = d$ and $Cb = c$.

Here $C : \mathcal{K}_{zu} \rightarrow \mathcal{K}_{zu}$ is the conjugation $Cf = \overline{f}u$ on \mathcal{K}_{zu} .

We remark that (7.15) is entirely analogous to the representation of quaternions of unit modulus using 2×2 complex matrices. This representation was exploited in [24] to study the characteristic function of a complex symmetric contraction.

7.5. Cartesian decomposition. Each f in \mathcal{K}_u enjoys a *Cartesian decomposition* $f = a + ib$ where the functions

$$a = \frac{1}{2}(f + Cf), \quad b = \frac{1}{2i}(f - Cf),$$

are both fixed by C . With respect to this decomposition the conjugation C on \mathcal{K}_u assumes the form $Cf = a - ib$. To describe the functions that belong

to \mathcal{K}_u , it suffices to describe those functions that are *C-real* (i.e., such that $Cf = f$).

If $f = Cf$, then

$$f = \overline{fz}u.$$

Since u has unit modulus a.e. on \mathbb{T} , by replacing u with a suitable unimodular constant multiple of u (which does not change the model space \mathcal{K}_u), we may assume that $u(\zeta) = \zeta$ for some ζ in \mathbb{T} .

If u has an ADC at ζ , then Theorem 6.11 tells us that the boundary kernel k_ζ belongs to \mathcal{K}_u . In the event that k_ζ fails to belong to \mathcal{K}_u (i.e., k_ζ is not in H^2), we still can assert that k_ζ resides in the *Smirnov class* N^+ (quotients of bounded analytic functions with outer denominator, see Definition 2.18) [33, Sect. 2.5]. A little arithmetic tells us that

$$\begin{aligned} k_\zeta/\overline{k_\zeta} &= \frac{1 - \overline{\zeta}u}{1 - \overline{\zeta}z} \cdot \frac{1 - \zeta\overline{z}}{1 - \zeta\overline{u}} \\ &= \frac{1 - \overline{\zeta}u}{1 - \zeta\overline{u}} \cdot \frac{1 - \zeta\overline{z}}{1 - \overline{\zeta}z} \\ &= \overline{\zeta}u \left(\frac{\zeta\overline{u} - 1}{1 - \zeta\overline{u}} \right) \cdot \zeta\overline{z} \left(\frac{\zeta z - 1}{1 - \overline{\zeta}z} \right) \\ &= \zeta\overline{\zeta}\overline{z}u \\ &= \overline{z}u, \end{aligned}$$

which, when substituted into the equation $f = \overline{fz}u$, reveals that

$$f/k_\zeta = \overline{f/k_\zeta}.$$

Thus a function f in \mathcal{K}_u satisfies $f = Cf$ if and only if f/k_ζ belongs to N^+ and is real almost everywhere on \mathbb{T} .

Definition 7.16. A function f belonging to the Smirnov class N^+ is called a *real Smirnov function* if its boundary function is real valued almost everywhere on \mathbb{T} . Denote the set of all real Smirnov functions by \mathcal{R}^+ .

The following elegant theorem of Helson [55] provides an explicit formula for real Smirnov functions.

Theorem 7.17. *If $f \in \mathcal{R}^+$, then there exist inner functions ψ_1, ψ_2 such that*

$$f = i \frac{\psi_1 + \psi_2}{\psi_1 - \psi_2} \tag{7.18}$$

and $\psi_1 - \psi_2$ is outer.

Proof. The function

$$\tau(z) = i \frac{1+z}{1-z}$$

maps \mathbb{D} onto the upper half-plane. Thus $\tau^{-1} \circ f$ is of bounded characteristic (i.e., a quotient of H^∞ functions) and is unimodular almost everywhere on

\mathbb{T} . Thus $\tau^{-1} \circ f = \psi_1/\psi_2$ is a quotient of inner functions and hence f has the desired form. \square

Since each f in \mathcal{K}_u can be expressed in the form $(\alpha + i\beta)k_\zeta$ where α and β belong to \mathcal{R}^+ , it follows from Helson's Theorem that any function in \mathcal{K}_u can be written in a simple algebraic manner using only a handful of inner functions. In particular, pseudocontinuations arise directly through the mechanism of Schwarz reflection (i.e., through pseudocontinuations of inner functions).

Unfortunately, the Helson representation (7.18) can be difficult to compute with. For instance, it is a somewhat messy calculation to find a Helson representation for the *Köbe function*

$$k(z) = \frac{z}{(1-z)^2} = z + 2z^2 + 3z^3 + \dots,$$

which maps \mathbb{D} bijectively onto $\mathbb{C} \setminus (-\infty, -\frac{1}{4}]$. We therefore propose a more constructive description of the functions in \mathcal{R}^+ , which originates in [45,51]. First note that if $f = I_f F$ is the inner-outer factorization of a function \mathcal{R}^+ , then

$$I_f F = \left[\frac{-4I_f}{(1-I_f)^2} \right] \cdot \left[\frac{(1-I_f)^2 F}{-4} \right],$$

where the first term is in \mathcal{R}^+ and has the same inner factor as f and where the second is *outer* and in \mathcal{R}^+ . Thus to describe functions in \mathcal{R}^+ , it suffices to describe *real outer (RO) functions*. An infinite product expansion, in terms of Cayley-like transforms of inner functions, is obtained in [51]. We refer the reader there for further details.

8. ALEKSANDROV-CLARK MEASURES

The study of Aleksandrov-Clark measures dates back to the seminal work of Clark [29] and it continued with deep work of Aleksandrov [5,6,9] and Poltoratski [73]. Since then, Aleksandrov-Clark measures have appeared in the study of spectral theory, composition operators, completeness problems, and mathematical physics. Several sources for this important topic, including references to their applications and the connections mentioned above, are [26,72,77].

8.1. Clark measures. If μ is a probability measure on \mathbb{T} then the function

$$F_\mu(z) := \int_{\mathbb{T}} \frac{\zeta + z}{\zeta - z} d\mu(\zeta)$$

is analytic on \mathbb{D} and satisfies $F_\mu(0) = 1$. Furthermore, a simple computation shows that

$$\Re F_\mu(z) = \int_{\mathbb{T}} \frac{1 - |z|^2}{|\zeta - z|^2} d\mu(\zeta)$$

is positive on \mathbb{D} . The following classical result of Herglotz asserts that this process can be reversed [33, p. 10].

Theorem 8.1 (Herglotz). *If F is analytic on \mathbb{D} , $F(0) = 1$, and $\Re F > 0$ on \mathbb{D} , then there is a unique probability measure μ on \mathbb{T} such that $F = F_\mu$.*

If u is inner and $u(0) = 0$, then for each α in \mathbb{T} the function

$$\frac{1 + \bar{\alpha}u(z)}{1 - \bar{\alpha}u(z)}$$

satisfies the hypothesis of Herglotz's theorem. Indeed, this function is clearly analytic on \mathbb{D} , evaluates to 1 at $z = 0$, and satisfies

$$\Re \left(\frac{1 + \bar{\alpha}u(z)}{1 - \bar{\alpha}u(z)} \right) = \frac{1 - |u(z)|^2}{|\alpha - u(z)|^2} > 0.$$

By Herglotz' Theorem, there is a unique probability measure σ_α on \mathbb{T} such that

$$\frac{1 + \bar{\alpha}u(z)}{1 - \bar{\alpha}u(z)} = \int \frac{\zeta + z}{\zeta - z} d\sigma_\alpha(\zeta). \quad (8.2)$$

The resulting family of measures

$$\{\sigma_\alpha : \alpha \in \mathbb{T}\}$$

are called the *Clark measures corresponding to u* (see Subsection 8.3 for their generalization, the *Aleksandrov-Clark measures*). The following proposition summarizes some of their basic properties (see [26] for details).

Proposition 8.3. *For an inner function u satisfying $u(0) = 0$, the corresponding family of Clark measures $\{\sigma_\alpha : \alpha \in \mathbb{T}\}$ satisfy the following.*

- (i) $\sigma_\alpha \perp m$ for all α .
- (ii) $\sigma_\alpha \perp \sigma_\beta$ for $\alpha \neq \beta$.
- (iii) σ_α has a point mass at ζ if and only if $u(\zeta) = \alpha$ and $|u'(\zeta)| < \infty$.
Furthermore

$$\sigma_\alpha(\{\zeta\}) = \frac{1}{|u'(\zeta)|}.$$

- (iv) A carrier¹¹ for σ_α is the set

$$\left\{ \zeta \in \mathbb{T} : \lim_{r \rightarrow 1^-} u(r\zeta) = \alpha \right\}.$$

This process can also be reversed. Starting with a singular probability measure μ on \mathbb{T} (e.g., $\mu = \delta_1$) one forms the *Herglotz integral*

$$H_\mu(z) := \int_{\mathbb{T}} \frac{\zeta + z}{\zeta - z} d\mu(\zeta), \quad z \in \mathbb{D}.$$

¹¹By a *carrier* for σ_α we mean a Borel set $C \subset \mathbb{T}$ for which $\sigma_\alpha(A \cap C) = \sigma_\alpha(A)$ for all Borel subsets $A \subset \mathbb{T}$. Carriers are not unique and a carrier is often different from a support. For example, if a measure consists of a dense set of point masses on \mathbb{T} , then a carrier would consist of just these point masses, whereas the support would be \mathbb{T} .

This function has positive real part on \mathbb{D} and satisfies $H(0) = 1$. Thus

$$u_\mu(z) := \frac{H(z) - 1}{H(z) + 1} \quad (z \in \mathbb{D})$$

satisfies

$$\Re \left(\frac{1 + u_\mu(z)}{1 - u_\mu(z)} \right) = \frac{1 - |u_\mu(z)|^2}{|1 - u_\mu(z)|^2} = \int_{\mathbb{T}} P_z(\zeta) d\mu(\zeta).$$

Using the fact that

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} P_{rw}(\zeta) d\mu(\zeta) = 0$$

for m -almost every $w \in \mathbb{T}$ (since $\mu \perp m$ – see (2.11)) we see that u_μ is inner and μ is the Clark measure with $\alpha = 1$ corresponding to u_μ .

8.2. Clark unitary operator. For an inner function u with $u(0) = 0$, recall that the compressed shift is the operator $S_u : \mathcal{K}_u \rightarrow \mathcal{K}_u$ defined by $S_u f = P_u(zf)$. For each $\alpha \in \mathbb{T}$ define

$$U_\alpha : \mathcal{K}_u \rightarrow \mathcal{K}_u, \quad U_\alpha := S_u + \alpha \left(1 \otimes \frac{u}{z} \right), \quad (8.4)$$

where $1 \otimes \frac{u}{z}$ denotes the rank one operator satisfying $(1 \otimes \frac{u}{z})f = \langle f, \frac{u}{z} \rangle 1$ (note that since $u(0) = 0$ the constant function 1 belongs to \mathcal{K}_u). The main theorem here is the following (see [26] for details).

Theorem 8.5 (Clark [29]). *For an inner function u with $u(0) = 0$, the operator (8.4) is a cyclic unitary operator whose spectral measure is carried by the Borel set*

$$\left\{ \zeta \in \mathbb{T} : \lim_{r \rightarrow 1^-} u(r\zeta) = \alpha \right\}.$$

The eigenvalues of U_α are the points ζ in \mathbb{T} so that $u(\zeta) = \alpha$ and $|u'(\zeta)| < \infty$. The corresponding eigenvectors are the boundary kernels k_ζ .

Example 8.6. Consider the model space $\mathcal{K}_{z^3} = \vee\{1, z, z^2\}$. Letting $S_3 := S_{z^3}$, we see that

$$S_3 1 = z, \quad S_3 z = z^2, \quad S_3 z^2 = 0,$$

so that the matrix representation of S_3 with respect to the basis $\{1, z, z^2\}$ is

$$\begin{bmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

The Clark unitary operator in this case is

$$U_\alpha = S_3 + \alpha(1 \otimes z^2).$$

Similarly, since

$$U_\alpha 1 = z, \quad U_\alpha z = z^2, \quad U_\alpha z^2 = \alpha,$$

the matrix representation of U_α with respect to this basis is

$$\begin{bmatrix} 0 & 0 & \alpha \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

The unitarity of U_α follows easily from the fact that $|\alpha| = 1$. The eigenvalues of U_α are precisely the solutions to $u(z) = \alpha$ (i.e., the three cube roots of α) and corresponding eigenvectors are the boundary kernels

$$k_\beta(z) = \frac{1 - \bar{\beta}^3 z^3}{1 - \bar{\beta}z} = 1 + \bar{\beta}z + \bar{\beta}^2 z^2,$$

where $\beta^3 = \alpha$. Each eigenvector k_β has norm $\sqrt{3}$, which is precisely the square root of the modulus of the angular derivative (namely $3z^2$) of z^3 at the point β in \mathbb{T} . Moreover, the normalized eigenvectors of U_α form an orthonormal basis for \mathcal{K}_{z^3} (as expected since U_α is a unitary operator on a finite dimensional space).

The Clark theory also gives us a concrete spectral representation for U_α via the Clark measure σ_α .

Theorem 8.7 (Clark [29]). *For an inner function u with $u(0) = 0$, let σ_α be the unique finite positive Borel measure on \mathbb{T} satisfying*

$$\frac{1 + \bar{\alpha}u(z)}{1 - \bar{\alpha}u(z)} = \int \frac{\zeta + z}{\zeta - z} d\sigma_\alpha(\zeta).$$

The operator

$$(V_\alpha f)(z) = (1 - \bar{\alpha}u(z)) \int \frac{f(\zeta)}{1 - \bar{\zeta}z} d\sigma_\alpha(\zeta)$$

is a unitary operator from $L^2(\sigma_\alpha)$ onto \mathcal{K}_u . Furthermore, if

$$Z_\alpha : L^2(\sigma_\alpha) \rightarrow L^2(\sigma_\alpha), \quad (Z_\alpha f)(\zeta) = \zeta f(\zeta),$$

then $V_\alpha Z_\alpha = U_\alpha V_\alpha$.

Example 8.8. Returning to Example 8.6. Let $\zeta_1, \zeta_2, \zeta_3$ be the three solutions to $u(z) = \alpha$ (i.e., to $z^3 = \alpha$). Then the discrete measure

$$d\sigma_\alpha = \frac{1}{3}\delta_{\zeta_1} + \frac{1}{3}\delta_{\zeta_2} + \frac{1}{3}\delta_{\zeta_3}$$

satisfies

$$\frac{1 + \bar{\alpha}z^3}{1 - \bar{\alpha}z^3} = \int \frac{\zeta + z}{\zeta - z} d\sigma_\alpha(\zeta).$$

From the theorem above,

$$V_\alpha : L^2(\sigma_\alpha) \rightarrow \mathcal{K}_{z^3}, \quad (V_\alpha f)(z) = (1 - \bar{\alpha}z^3) \int \frac{f(\zeta)}{1 - \bar{\zeta}z} d\sigma_\alpha(\zeta).$$

To see this worked out, note that if

$$f = c_1\chi_{\zeta_1} + c_2\chi_{\zeta_2} + c_3\chi_{\zeta_3},$$

a typical element of $L^2(\sigma_\alpha)$, then

$$(Vf)(z) = (1 - \bar{\alpha}z^3) \left(c_1 \frac{1/3}{1 - \bar{\zeta}_1 z} + c_2 \frac{1/3}{1 - \bar{\zeta}_2 z} + c_3 \frac{1/3}{1 - \bar{\zeta}_3 z} \right).$$

Since $\zeta_j^3 = \alpha$, one can verify that the expression above is a polynomial of degree at most 2, which are precisely the elements of \mathcal{K}_{z^3} . Furthermore, $\{\sqrt{3}\chi_{\zeta_1}, \sqrt{3}\chi_{\zeta_2}, \sqrt{3}\chi_{\zeta_3}\}$ is an orthonormal basis for $L^2(\sigma_\alpha)$ and

$$(V_\alpha \sqrt{3}\chi_{\zeta_j})(z) = \frac{\sqrt{3}}{3} k_{\zeta_j}(z).$$

But since

$$\left\{ \frac{\sqrt{3}}{3} k_{\zeta_j}(z) : j = 1, 2, 3 \right\}$$

is an orthonormal basis for \mathcal{K}_{z^3} we see that V_α is unitary. From here one can easily see, by basic linear algebra, that V_α intertwines U_α with Z_α .

The adjoint of the unitary operator V_α is of particular interest. We know that $V_\alpha f$ belongs to \mathcal{K}_u for every f in $L^2(\sigma_\alpha)$ and, as such, $V_\alpha f$ has a radial limit almost everywhere with respect to Lebesgue measure m . The following theorem of Poltoratski [73] says much more.

Theorem 8.9 (Poltoratski [73]). *For $f \in L^2(\sigma_\alpha)$,*

$$\lim_{r \rightarrow 1^-} (V_\alpha f)(r\zeta) = f(\zeta)$$

for σ_α -almost every $\zeta \in \mathbb{T}$.

The significance of this result is that it says that functions in model spaces (i.e., the functions $V_\alpha f$ for $f \in L^2(\sigma_\alpha)$ in the previous theorem) have nontangential limits on finer sets than generic functions from H^2 . We have seen a result along these lines already in Theorem 6.11. The previous theorem says a bit more in that all functions from \mathcal{K}_u have non-tangential limits σ_α almost everywhere. This is especially notable since $\sigma_\alpha \perp m$ and a classical result [62,63] says that if E is any closed subset of \mathbb{T} with Lebesgue measure zero, then there exists an f in H^2 (which can be taken to be inner) whose radial limits do not exist on E . For example, if σ_α has an isolated point pass at ζ , then every function in \mathcal{K}_u has a nontangential limit at ζ . In this particular case, the result should not be surprising since, by Proposition 8.3, u will have a finite angular derivative at ζ and so by the Ahern-Clark result (Theorem 6.11) every function in \mathcal{K}_u has a nontangential limit at ζ .

Here is one more fascinating and useful gem about Clark measures due to Aleksandrov [5] (see also [26]).

Theorem 8.10 (Aleksandrov [5]). *Let u be an inner function with associated family of Clark measures $\{\sigma_\alpha : \alpha \in \mathbb{T}\}$. If $f \in C(\mathbb{T})$, then*

$$\int_{\mathbb{T}} \left(\int_{\mathbb{T}} f(\zeta) d\sigma_\alpha(\zeta) \right) dm(\alpha) = \int_{\mathbb{T}} f(\xi) dm(\xi).$$

Proof. For a fixed $z \in \mathbb{D}$ notice that if $P_z(\zeta)$ is the Poisson kernel, then taking real parts of both sides of (8.2) we get

$$\begin{aligned} \int_{\mathbb{T}} \left(\int_{\mathbb{T}} P_z(\zeta) d\sigma_\alpha(\zeta) \right) dm(\alpha) &= \int_{\mathbb{T}} \left(\frac{1 - |u(z)|^2}{|\alpha - u(z)|^2} \right) dm(\alpha) \\ &= \int_{\mathbb{T}} P_{u(z)}(\alpha) dm(\alpha) \\ &= 1 \\ &= \int_{\mathbb{T}} P_z(\zeta) dm(\zeta). \end{aligned}$$

Thus the theorem works for finite linear combinations of Poisson kernels. Using the fact that finite linear combinations of Poisson kernels are dense in $C(\mathbb{T})$ one can use a limiting argument to get the general result¹². \square

Remark 8.11. Aleksandrov [5] showed that $C(\mathbb{T})$ can be replaced by L^1 in the preceding disintegration theorem. There are a few technical issues to work out here. For example, the inner integrals

$$\int f(\zeta) d\sigma_\alpha(\zeta)$$

in the disintegration formula do not seem to be well defined for L^1 functions since indeed the measure σ_α can contain point masses on \mathbb{T} while L^1 functions are defined merely m -almost everywhere. However, amazingly, the function

$$\alpha \mapsto \int f(\zeta) d\sigma_\alpha(\zeta)$$

is defined for m -almost every α and is integrable. An argument with the monotone class theorem is used to prove this more general result. See also [26] for the details.

8.3. Aleksandrov-Clark measures. The alert reader will have noticed that the title of this section was Aleksandrov-Clark measures but we seem to have only discussed *Clark* measures. As it turns out, one can still examine the equation

$$\frac{1 + \bar{\alpha}u(z)}{1 - \bar{\alpha}u(z)} = \int \frac{\zeta + z}{\zeta - z} d\sigma_\alpha(\zeta)$$

but where u belongs to H^∞ and $\|u\|_\infty \leq 1$, but u is not necessarily inner. The measures σ_α will no longer be singular but they do form an interesting class of measures explored by Aleksandrov and others (see [26] for references). One can identify all the parts of these measures (absolutely continuous, singular continuous, point masses, etc.) as well as develop a decomposition theorem as before. Versions of these measures appear in mathematical physics through some papers of B. Simon and T. Wolff [90, 91].

¹²See [26, Ch. 9] for the details on the density of the linear span of the Poisson kernels in $C(\mathbb{T})$ as well as the completion of this limiting argument

9. COMPLETENESS PROBLEMS

In Subsection 5.3, we discussed circumstances under which a sequence $\{k_\lambda\}_{\lambda \in \Lambda}$ of kernel functions forms a Riesz basis for \mathcal{K}_u (see Theorem 5.16). More generally, one can ask the following question: If Λ is a sequence in \mathbb{D} , when does

$$K_\Lambda := \bigvee_{\lambda \in \Lambda} k_\lambda = \mathcal{K}_u?$$

Such problems are known as *completeness problems*.

By considering orthogonal complements and the reproducing property of k_λ , we see that $K_\Lambda = \mathcal{K}_u$ precisely when there are no nonzero functions in \mathcal{K}_u that vanish on Λ . We have already seen that $K_\Lambda = \mathcal{K}_u$ if Λ is not a Blaschke sequence or if Λ has an accumulation point in \mathbb{D} (Proposition 5.8). A little further thought shows that the same holds if Λ has an accumulation point in $\mathbb{T} \setminus \sigma(u)$ since otherwise there would exist a nonzero function in \mathcal{K}_u whose analytic continuation vanishes on a set having a limit point in its domain (Proposition 6.9). For general Blaschke sequences, the situation is more complicated. However, we can rephrase the completeness problem in terms of Toeplitz kernels¹³.

Proposition 9.1. *If Λ is a Blaschke sequence in \mathbb{D} and B is the Blaschke product corresponding to Λ , then for any inner function u we have*

$$K_\Lambda \neq \mathcal{K}_u \iff \ker T_{\bar{u}B} \neq \{0\}.$$

Proof. Note that if $K_\Lambda \neq \mathcal{K}_u$, then there is a nonzero function f in \mathcal{K}_u that vanishes on Λ . This happens precisely when $f = gB$ for some g in H^2 that is not identically zero. But since f belongs to \mathcal{K}_u we know that $\bar{u}f$ is contained in $\overline{zH^2}$. This means that $\bar{u}Bg$ belongs to $\overline{zH^2}$. Now apply the Riesz projection (of L^2 onto H^2) to $\bar{u}Bg$ to see that $T_{\bar{u}B}g = P(\bar{u}Bg) = 0$. For the other direction, simply reverse the argument. \square

Clark's approach in [29] to the completeness problem is based on the following related approximation problem of Paley-Wiener. Suppose that $\{x_n\}_{n \geq 1}$ is an orthonormal basis for a Hilbert space \mathcal{H} and $\{y_n\}_{n \geq 1}$ is a sequence in \mathcal{H} . If these sequences are *close* in some way, does $\{y_n\}_{n \geq 1}$ span \mathcal{H} ? There are various results which say this is often the case (see [74, Sec. 86] or [67] for a survey of these results) with various interpretations of the term *close*. We will see a specific example of this below.

Clark's approach to the completeness problem $K_\Lambda = \mathcal{K}_u$ was to find an orthonormal basis for \mathcal{K}_u consisting of (normalized) boundary kernel functions. One way to accomplish this is to find a unitary operator U on \mathcal{K}_u whose spectrum consists only of eigenvalues (pure point spectrum) and such that the eigenvectors of U are boundary kernel functions. But indeed we have already seen this can be done. From Clark's theorem (Theorem

¹³Kernels of Toeplitz operators have been studied before and we refer the reader to the papers [35, 46, 81] for further details.

8.5) we see that U_α is unitary and, under the right spectral circumstances (i.e., the corresponding Clark measure σ_α is purely atomic), its normalized eigenvectors

$$\left\{ \frac{k_\zeta}{\sqrt{|u'(\zeta)|}} : u(\zeta) = \alpha, |u'(\zeta)| < \infty \right\}$$

form an orthonormal basis for \mathcal{K}_u . From here one can apply a Paley-Wiener type result to obtain completeness results for the family $\{k_\lambda : \lambda \in \Lambda\}$. Let us give two illustrative examples from the original Clark paper [29].

Example 9.2. Suppose that u is an inner function and $\alpha \in \mathbb{T}$ such that the corresponding Clark measure σ_α is discrete, i.e.,

$$\sigma_\alpha = \sum_{n=1}^{\infty} \frac{1}{|u'(\zeta_n)|} \delta_{\zeta_n}.$$

Note the use of Proposition 8.3 here where $\{\zeta_n\}_{n \geq 1}$ are the solutions to $u = \alpha$. Since σ_α is a discrete measure and U_α is unitary, the normalized eigenvectors

$$h_{\zeta_n} := \frac{k_{\zeta_n}}{\sqrt{|u'(\zeta_n)|}}$$

are an orthonormal basis for \mathcal{K}_u . For our given sequence $\Lambda = \{\lambda_n\}_{n \geq 1} \subset \mathbb{D}$, let

$$h_{\lambda_n} := \frac{k_{\lambda_n}}{\|k_{\lambda_n}\|}$$

be the normalized kernel functions. If the sequence $\{\lambda_n\}_{n \geq 1}$ satisfies

$$\sum_{n=1}^{\infty} \|h_{\zeta_n} - h_{\lambda_n}\|^2 < 1,$$

then a generalization of a theorem of Paley and Wiener (see [74, Sec. 86]) says that $\{h_{\lambda_n}\}_{n \geq 1}$ forms an unconditional basis for \mathcal{K}_u , meaning that every f in \mathcal{K}_u has an expansion $f = \sum_n a_n h_{\lambda_n}$.

This next example brings in some earlier work of Sarason [78].

Example 9.3. Consider the model space \mathcal{K}_u with inner function

$$u(z) := \exp\left(\frac{z+1}{z-1}\right).$$

The Clark measure σ_1 corresponding to u has the set $\{u = 1\}$ as its carrier, which turns out to be the discrete set

$$\zeta_n := \frac{1 + 2\pi in}{1 - 2\pi in}, \quad (n \in \mathbb{Z})$$

In [78] it is shown that the operator

$$[Wf](z) := \frac{\sqrt{2}}{1-z} \int_0^1 f(t) u^t(z) dt$$

defines unitary operator which maps \mathcal{K}_u onto $L^2[0, 1]$. In our current setting, the real usefulness of this operator comes from the formula

$$W(e^{i\gamma t}) = qk_\lambda, \quad \gamma := i\frac{1 + \bar{\lambda}}{1 - \bar{\lambda}},$$

where $|\lambda| \leq 1$ and q is a constant. In particular, this means that W maps each orthonormal basis element $e^{2\pi int}$ for $L^2[0, 1]$ to a constant times k_{ζ_n} . So, for a given a sequence $\{\lambda_n\}_{n \in \mathbb{Z}} \subset \mathbb{D}$ satisfying

$$\max_{n \in \mathbb{Z}} |\gamma_n - 2\pi n| < \frac{\log 2}{\pi},$$

a theorem of Paley and Wiener (see [74, Sec. 86]) says that $\{e^{i\gamma_n t}\}_{n \in \mathbb{Z}}$ forms an unconditional basis for $L^2[0, 1]$. Under this criterion, we get that the normalized kernels $\{k_{\lambda_n}/\|k_{\lambda_n}\| : n \in \mathbb{Z}\}$ form an unconditional basis for \mathcal{K}_u .

We will consider other unitary operators that transfer orthonormal bases of L^2 to kernel functions in the next section.

We also point out a recent result of Baranov and Dyakonov [13] that also discusses when a perturbation of Clark bases form Riesz bases.

10. MODEL SPACES FOR THE UPPER-HALF PLANE

10.1. **The Hardy space again.** Let

$$\mathbb{C}_+ := \{z \in \mathbb{C} : \Im z > 0\}$$

denote the upper half plane and let \mathcal{H}^2 denote the *Hardy space of the upper half plane*. These are the analytic functions f on \mathbb{C}_+ for which

$$\sup_{y>0} \int_{-\infty}^{\infty} |f(x + iy)|^2 dx < \infty.$$

The reader will immediately notice the analogue of the “bounded integral mean” condition (2.1) from the Hardy space H^2 on the unit disk. As it turns out, the majority of the theory from H^2 (on the disk) transfers over *mutatis mutandis* to \mathcal{H}^2 . For example, every f in \mathcal{H}^2 has an almost everywhere well-defined “radial” boundary function

$$f(x) := \lim_{y \rightarrow 0^+} f(x + iy)$$

and this function belongs to $L^2(dx) := L^2(\mathbb{R}, dx)$. Moreover,

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \sup_{y>0} \int_{-\infty}^{\infty} |f(x + iy)|^2 dx.$$

This allows us to endow \mathcal{H}^2 with an inner product

$$\langle f, g \rangle := \int_{-\infty}^{\infty} f(x)\overline{g(x)} dx,$$

where, in a manner that is analogous to the H^2 case, $f(x)$ and $g(x)$ are the almost everywhere defined boundary functions of f, g in \mathcal{H}^2 . Still further, equating H^2 with the functions in $L^2(\mathbb{T}, m)$ whose negatively indexed Fourier coefficients vanish, we have the corresponding characterization

$$\mathcal{H}^2 := \{\widehat{f} : f \in L^2(0, \infty)\},$$

where

$$\widehat{f}(t) = \int_{-\infty}^{\infty} f(x)e^{-2\pi ixt} dx$$

is the *Fourier transform* of f .

The reason that most of the Hardy space theory on the disk can be imported into \mathcal{H}^2 is because of the fact that the operator

$$\mathcal{U} : L^2(m) \rightarrow L^2(dx), \quad (\mathcal{U}f)(x) := \frac{1}{\sqrt{\pi}(x+i)}f(w(x)),$$

where

$$w(z) := \frac{z-i}{z+i}$$

is a conformal map from \mathbb{C}_+ onto \mathbb{D} , is a unitary map from $L^2(\mathbb{T}, m)$ onto $L^2(\mathbb{R}, dx)$. In particular, \mathcal{U} maps H^2 unitarily onto \mathcal{H}^2 and maps $L^\infty(\mathbb{T}, m)$ isometrically onto $L^\infty(\mathbb{R}, dx)$.

10.2. Inner functions. We say that an analytic function Θ on \mathbb{C}_+ is *inner* if $|\Theta(z)| \leq 1$ for all $z \in \mathbb{C}_+$ and if the almost everywhere defined boundary function $\Theta(x)$ is unimodular almost everywhere on \mathbb{R} . The two most basic types of inner functions on \mathbb{C}_+ are

$$S^c(z) := e^{icz} \quad (c > 0)$$

and

$$b_\lambda(z) := \frac{z-\lambda}{z-\bar{\lambda}}. \quad (\lambda \in \mathbb{C}_+)$$

The first class of examples are the most basic type of *singular inner function* on \mathbb{C}_+ and the second type is the most basic type of *Blaschke product*.

Further examples of singular inner functions are given by

$$S_{\mu,c}(z) := e^{icz} \exp\left(-\frac{1}{\pi i} \int_{-\infty}^{\infty} \left(\frac{1}{x-z} - \frac{x}{1+x^2}\right) d\mu(x)\right),$$

where $c > 0$ and μ is a positive measure on \mathbb{R} that is singular with respect to Lebesgue measure and is *Poisson finite*, in the sense that

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} d\mu(x) < \infty.$$

The general Blaschke product is formed by specifying its zeros

$$\Lambda = \{\lambda_n\}_{n \geq 1} \subset \mathbb{C}_+ \setminus \{i\}$$

and forming the product

$$B_\Lambda(z) = b_i(z)^m \prod_{n \geq 1} \epsilon_n b_{\lambda_n}(z),$$

where the constants

$$\epsilon_n := \frac{|\lambda_n^2 + 1|}{\lambda_n^2 + 1}$$

are chosen so that $\epsilon_n b_{\lambda_n}(i) > 0$. It is well known [52] that the product above converges uniformly on compact subsets of \mathbb{C}_+ to an inner function if and only if the zeros $\{\lambda_n\}_{n \geq 1}$ satisfy the *Blaschke condition*

$$\sum_{n \geq 1} \frac{\Im \lambda_n}{1 + |\lambda_n|^2} < \infty.$$

Every inner function Θ can be factored uniquely as

$$\Theta = e^{i\gamma} B_\Lambda S_{\mu,c} \quad (\gamma \in [0, 2\pi))$$

10.3. Model spaces. For an inner function Θ define a *model space* \mathcal{K}_Θ on the upper-half plane as

$$\mathcal{K}_\Theta := \mathcal{H}^2 \ominus \Theta \mathcal{H}^2.$$

Much of the theory for the model spaces \mathcal{K}_u on the disk carries over to the upper-half plane. For instance, via Proposition 3.3, we can regard \mathcal{K}_Θ as a space of boundary functions on \mathbb{R} by noting that

$$\mathcal{K}_\Theta := \mathcal{H}^2 \cap \overline{\Theta \mathcal{H}^2}. \quad (10.1)$$

An inner function Θ has an analytic continuation across $\mathbb{R} \setminus \sigma(\Theta)$, where

$$\sigma(\Theta) := \left\{ z \in \mathbb{C}_+^- : \liminf_{\lambda \rightarrow z} |\Theta(\lambda)| = 0 \right\}$$

is the *spectrum* of Θ , which turns to be the union of the closure the zeros of Θ along with the support of the singular measure for Θ . Every f in \mathcal{K}_Θ has an analytic continuation across $\mathbb{R} \setminus \sigma(\Theta)$. Using the identity (10.1), we see that for each f in \mathcal{K}_Θ there is a corresponding g in \mathcal{H}^2 for which $f = \overline{g}\Theta$ almost everywhere on \mathbb{R} . As was done with (6.7) this allows us to obtain a formula

$$\tilde{f}(z) := \frac{\overline{g(\overline{z})}}{\overline{\Theta(\overline{z})}} \quad (z \in \mathbb{C}_-)$$

for a pseudocontinuation of f (to the lower half-plane \mathbb{C}_-) of bounded type.

Of particular importance here are the reproducing kernel functions

$$K_\lambda(z) = \frac{i}{2\pi} \frac{1 - \overline{\Theta(\lambda)}\Theta(z)}{z - \overline{\lambda}} \quad (\lambda, z \in \mathbb{C}_+)$$

These belong to \mathcal{K}_Θ and satisfy

$$f(\lambda) = \langle f, K_\lambda \rangle \quad (\lambda \in \mathbb{C}_+)$$

10.4. Clark theory again. As to be expected, there is a Clark theory for \mathcal{K}_Θ . Indeed, for an inner function Θ , the function

$$m = i \frac{1 + \Theta}{1 - \Theta}$$

is analytic on \mathbb{C}_+ and satisfies $\Im m \geq 0$ there (i.e., m is a *Herglotz function*). In this setting, the Herglotz representation theorem guarantees the existence of parameters $b \geq 0, c \in \mathbb{R}$, and a positive Poisson finite measure μ_Θ on \mathbb{R} so that

$$m(\lambda) = bz + c + \frac{1}{\pi} \int \left(\frac{1}{x - \lambda} - \frac{x}{1 + x^2} \right) d\mu_\Theta(x).$$

Using Poltoratski's result (Theorem 8.9) we can define the operator

$$Q_\Theta : \mathcal{K}_\Theta \rightarrow L^2(\mu_\Theta), \quad Qf = f|_{C_\Theta}, \quad (10.2)$$

where C_Θ is a carrier for Θ . This operator turns out to be unitary.

As a final remark, the alert reader is probably getting *déjà vu* when looking at the function m and its associated inner function Θ . The current discussion is probably reminding the reader of our earlier treatment of Clark measures. Indeed, if one begins with an inner Θ and looks at the family of inner functions

$$\{\bar{\alpha}\Theta : \alpha \in \mathbb{T}\},$$

there is an associated family of Poisson finite measures

$$\{\mu_{\bar{\alpha}\Theta} : \alpha \in \mathbb{T}\},$$

the Clark measures associated with Θ . Many of the properties of Clark measures that hold for inner u on \mathbb{D} have direct analogs for inner functions Θ on \mathbb{C}_+ .

10.5. Weyl-Titchmarsh inner functions. In this section we point out a relationship between Schrödinger operators and model spaces. In order not to get too deep into technical details, we will be a little vague about proper definitions. The reader looking for precision should consult [64, 95].

For a real potential q in $L^2(a, b)$, define the *Schrödinger operator*

$$u \mapsto u'' + qu.$$

If one imposes the boundary condition

$$u(b) \cos \varphi + u'(b) \sin \varphi = 0,$$

this operator becomes a densely defined, self-adjoint operator on $L^2(a, b)$.

For each λ in \mathbb{C} the differential equation $u'' + qu = \lambda u$ has a non-trivial solution u_λ and one can form the *Weyl-Titchmarsh function* (at a) by

$$m(\lambda) := \frac{u'_\lambda(a)}{u_\lambda(a)}.$$

One can show that the function Θ defined on \mathbb{C}_+ by

$$\Theta := \frac{m - i}{m + i}$$

is inner.

For each $\lambda \in \mathbb{C}_+$ define

$$w_\lambda(x) := \frac{u_\lambda(x)}{u'_\lambda(a) + iu_\lambda(a)}$$

and notice that w_λ is also a solution to the Schrödinger equation $u'' + qu = \lambda u$, the so called *normalized solution*. Now define the following transform W on $L^2(a, b)$ (sometimes called the *modified Fourier transform*) by

$$[Wf](\lambda) := \int_a^b f(x)w_\lambda(x)dx \quad (\lambda \in \mathbb{C})$$

Schrödinger operators and model spaces are related by means of the following theorem from [64].

Theorem 10.3 (Makarov-Poltoratski). *The modified Fourier transform W is, up to a factor of $\sqrt{\pi}$, a unitary operator from $L^2(a, b)$ onto \mathcal{H}_Θ . Furthermore, $W\overline{w_\lambda} = \pi K_\lambda$.*

With the preceding theorem one can recast the completeness problems for solutions of the Schrödinger equation in $L^2(a, b)$ in terms of the completeness problem for reproducing kernels in \mathcal{H}_Θ . Secondly, the unitary operator $WQ_\Theta : L^2(a, b) \rightarrow L^2(\mu_\Theta)$ yields a spectral representation of the Schrödinger operator $u \mapsto u'' + qu$ in that it is unitarily equivalent to the densely defined operator multiplication by the independent variable on $L^2(\mu_\Theta)$.

11. GENERALIZATIONS OF MODEL SPACES

11.1. Model spaces in H^p . One can also consider model subspaces of H^p for $p \neq 2$. Recall that if $0 < p < \infty$, then H^p is the space of analytic functions f on \mathbb{D} for which

$$\|f\|_p := \lim_{r \rightarrow 1^-} \left(\int_{\mathbb{T}} |f(r\zeta)|^2 dm(\zeta) \right)^{1/2}$$

is finite. If $p \geq 1$, then H^p is a Banach space while if $0 < p < 1$, then H^p is a topological vector space. Almost all of the basic function-theoretic results that hold for H^2 functions carry over to H^p (e.g., each H^p has an almost everywhere defined nontangential boundary function that belongs to L^p).

For $p \neq 2$, the spaces H^p are not Hilbert spaces, although they do have readily identifiable duals:

- For $1 < p < \infty$, the dual space $(H^p)^*$ can be identified with H^q , where q is the Hölder conjugate index to p (i.e., $\frac{1}{p} + \frac{1}{q} = 1$).
- For $p = 1$, the dual space $(H^1)^*$ can be identified with the space of all analytic functions of bounded mean oscillation (BMOA),
- For $0 < p < 1$, $(H^p)^*$ can be identified with certain classes of smooth functions on \mathbb{D}^- (Lipschitz or Zygmund classes).

In each of these cases, the dual pairing is given by the *Cauchy pairing*

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} f(r\zeta) \overline{g(r\zeta)} dm(\zeta) \quad (11.1)$$

where $f \in H^p$ and $g \in (H^p)^*$. Standard references are [33, 52].

The unilateral shift $S : H^p \rightarrow H^p$ defined as usual by $Sf = zf$ is continuous on H^p and, for $1 \leq p < \infty$, the invariant subspaces of S are still the Beurling-type invariant subspaces uH^p for some inner u . The invariant subspaces of the backward shift¹⁴

$$Bf = \frac{f - f(0)}{z}$$

on H^p are known to be

$$\mathcal{K}_u^p := H^p \cap \overline{uzH^p} \quad (1 \leq p < \infty)$$

For $1 < p < \infty$, the proof relies on using the duality relationship between $(H^p)^*$ and H^q via the integral pairing (11.1) to compute the annihilator of uH^q . For $p = 1$, the backward shift invariant subspaces are still the same (i.e., $\mathcal{K}_u^1 = H^1 \cap \overline{uzH^1}$), except that the proof has to deal with some complications from the more difficult dual space of H^1 . When $0 < p < 1$, the backward shift invariant subspaces are too complicated to describe in this survey. A full accounting all of these results are found in [27]. Most of the function-theoretic results of this survey for model spaces \mathcal{K}_u in H^2 can be restated appropriately for \mathcal{K}_u^p spaces – usually with nearly the same proof.

11.2. deBranges-Rovnyak spaces. There is an important generalization of model spaces, the *deBranges-Rovnyak spaces*, that play an increasingly important role in analysis. Unlike the model spaces \mathcal{K}_u , which are parameterized by inner functions u , these spaces are parameterized by functions u in H_1^∞ , the unit ball in H^∞ (i.e., $u \in H^\infty, \|u\|_\infty \leq 1$). These spaces often have similar properties as model spaces but they also have many differences. We will not go into great detail here but just point out the existence of these types of spaces and their very basic properties. The standard reference for this subject is Sarason's book [82] from 1994, although a new book by Fricain and Mashreghi should appear soon.

For $u \in H_1^\infty$ (not necessarily inner!), define the kernel function

$$k_\lambda(z) := \frac{1 - \overline{u(\lambda)}u(z)}{1 - \overline{\lambda}z}. \quad (\lambda, z \in \mathbb{D})$$

The author will recognize this kernel, when u is an inner function, as the reproducing kernel for the model space \mathcal{K}_u , but here u is not necessarily inner. We construct a reproducing kernel Hilbert space $\mathcal{H}(u)$ of analytic

¹⁴Much of the literature here abuses notation here and often uses the notation $S^*f = (f - f(0))/z$ for the backward shift on H^p even though, technically speaking, the adjoint S^* is defined on $(H^p)^*$ and not H^p .

functions on \mathbb{D} from this kernel in the following way. First we notice that this kernel is a positive in the sense that

$$\sum_{1 \leq l, j \leq n} c_j \bar{c}_l k_{\lambda_j}(\lambda_l) \geq 0$$

for every set of constants c_1, \dots, c_n and points $\lambda_1, \dots, \lambda_n$ in \mathbb{D} . We initially populate our space $\mathcal{H}(u)$ with finite linear combinations of kernel functions and define a norm on these functions in a way that makes the k_λ the reproducing kernels for $\mathcal{H}(u)$. We do this by defining

$$\left\| \sum_{j=1}^n c_j k_{\lambda_j} \right\|^2 = \left\langle \sum_{j=1}^n c_j k_{\lambda_j}, \sum_{l=1}^n c_l k_{\lambda_l} \right\rangle := \sum_{1 \leq l, j \leq n} c_j \bar{c}_l k_{\lambda_j}(\lambda_l).$$

One can check that this is a norm and that the corresponding inner product on the vector space of finite linear combinations of kernel functions, make this vector space a pre-Hilbert space. The deBranges-Rovnyak space $\mathcal{H}(u)$ is the closure of this pre-Hilbert space under this norm. A more standard way of defining $\mathcal{H}(u)$ in terms of Toeplitz operators is described in [82].

Using the positivity of the kernel, one can prove that $\mathcal{H}(u)$ is reproducing kernel Hilbert space that is contractively contained in H^2 . When $\|u\|_\infty < 1$, then $\mathcal{H}(u) = H^2$ with an equivalent norm. On the other hand when u is inner, then $\mathcal{H}(u) = \mathcal{K}_u$ with equality of norms.

Recall that the compressed shift S_u on \mathcal{K}_u serves as a model for certain types of contraction operators. It turns out that $\mathcal{H}(u)$ spaces can be used to model other types of contractions where the condition (i) of Theorem 4.1 is somewhat relaxed.

12. TRUNCATED TOEPLITZ OPERATORS

We do not attempt to give a complete overview of the rapidly developing theory of truncated Toeplitz operators. The recent survey article [50] and the seminal article [83] of Sarason should be consulted for a more thorough treatment. We content ourselves here with a brief summary of the principal definitions and basic results.

Recall from Subsection 3.1 that the Toeplitz operator $T_\varphi : H^2 \rightarrow H^2$ with symbol φ in L^∞ is defined by $P(\varphi f)$, where P is the orthogonal projection from L^2 onto H^2 . One can show that T_φ is a bounded operator on H^2 that satisfies $T_\varphi^* = T_{\bar{\varphi}}$ and $\|T_\varphi\| = \|\varphi\|_\infty$. The (j, k) entry of the matrix representation of T_φ with respect to the orthonormal normal basis $\{1, z, z^2, \dots\}$ of H^2 is $\hat{\varphi}(k - j)$, which yields an infinite Toeplitz matrix (constant along the diagonals).

An old result of Brown and Halmos [19] gives a convenient algebraic characterization of Toeplitz operators: a bounded operator T on H^2 is a Toeplitz operator if and only

$$T = STS^*, \tag{12.1}$$

where S is the unilateral shift on H^2 . Many other things are known about Toeplitz operators: their spectrum, essential spectrum, commutator ideals, algebras generated by certain collections of Toeplitz operators, etc. These topics are all thoroughly discussed in [18, 32, 66].

It is well-known that the commutant

$$\{S\}' := \{A \in \mathcal{B}(H^2) : SA = AS\}$$

of the unilateral shift S is equal to the set

$$\{T_\varphi : \varphi \in H^\infty\}$$

of all *analytic* Toeplitz operators. This naturally leads one to consider the commutant $\{S_u\}'$ of the compressed shift S_u . The answer, due to Sarason [79], prompts us to consider the following.

Definition 12.2. If u is inner and $\varphi \in L^\infty$, then the *truncated Toeplitz operator* (TTO) with symbol φ is the operator $A_\varphi^u : \mathcal{K}_u \rightarrow \mathcal{K}_u$ defined by $A_\varphi^u f = P_u(\varphi f)$, where P_u is the orthogonal projection (5.5) of L^2 onto \mathcal{K}_u .

Theorem 12.3 (Sarason). *For inner u ,*

$$\{S_u\}' = \{A_\varphi^u : \varphi \in H^\infty\}.$$

This theorem, which initiated the general study of *commutant lifting theorems*, gives an initial impetus for the study of truncated Toeplitz operators. However, to obtain the full class of truncated Toeplitz operators, one needs to consider symbols in L^2 , as opposed to L^∞ . For φ in L^2 , one can define an operator A_φ^u on $\mathcal{K}_u \cap H^\infty$ (which is dense in \mathcal{K}_u via Proposition 5.11) by

$$A_\varphi^u f := P_u(\varphi f).$$

If A_φ^u can be extended to a bounded operator on all of \mathcal{K}_u , then we say A_φ^u is a bounded truncated Toeplitz operator. When there is no chance of confusion, we write A_φ instead of A_φ^u . We let \mathcal{T}_u denote the set of all bounded truncated Toeplitz operators on \mathcal{K}_u , remarking that \mathcal{T}_u is a weakly closed linear space.

There are some superficial similarities between truncated Toeplitz operators and Toeplitz operators. For instance, one has the adjoint formula $A_\varphi^* = A_{\overline{\varphi}}$ and an analogue of the Brown-Halmos result (12.1): a bounded operator A on \mathcal{K}_u is a truncated Toeplitz operator if and only if

$$A_z A A_z^* = A + R,$$

where R is a certain operator on \mathcal{K}_u of rank ≤ 2 . However, the differences are greater than the similarities. For instance, the symbol of a Toeplitz operator is unique, yet for truncated Toeplitz operators $A_\varphi = A_\psi$ if and only if

$$\varphi - \psi \in uH^2 + \overline{uH^2}.$$

In particular, the inequality $\|A_\varphi\| \leq \|\varphi\|_\infty$ is frequently strict. While there are no compact Toeplitz operators, many compact TTOs exist (e.g., any TTO

on \mathcal{K}_u if $\dim \mathcal{K}_u < \infty$). Finally, not every bounded truncated Toeplitz operator can be represented with a bounded symbol.

Recall from Section 7 that $Cf = \overline{f}zu$ defines a conjugation on \mathcal{K}_u . It turns out that each truncated Toeplitz operator is *complex symmetric*, in the sense that $A_\varphi = CA_\varphi^*C$. More generally, we say that a bounded operator T on a complex Hilbert space \mathcal{H} is a complex symmetric operator (CSO) if there exists a conjugation C on \mathcal{H} so that $T = CT^*C$ [39–43]. The class of complex symmetric operators is surprisingly large and, somewhat surprisingly, many CSOs can be shown to be unitarily equivalent to truncated Toeplitz operators [28, 48, 50, 92]. Clarifying the precise relationship between CSOs and TTOs is an ongoing effort, chronicled to some extent in [50].

Finally, we mention the following beautiful connection between Clark’s unitary operators (Section 8) and truncated Toeplitz operators.

Theorem 12.4. *If $\varphi \in L^\infty$ and u is inner, then*

$$A_\varphi^u = \int \varphi(U_\alpha) dm(\alpha),$$

in the sense that

$$\langle A_\varphi^u f, g \rangle = \int \langle \varphi(U_\alpha) f, g \rangle dm(\alpha),$$

for f, g in \mathcal{K}_u .

13. THINGS WE DID NOT MENTION

In this final section we point the reader in the direction of other aspects of model spaces that we do not have the time to fully explore.

13.1. Carleson measures. Carleson in [21, 22] (see also [52]) characterized those measures μ on \mathbb{D} for which the inclusion operator from H^2 to $L^2(\mu)$ is continuous, i.e.,

$$\int_{\mathbb{D}} |f|^2 d\mu \leq C \|f\|^2 \quad (f \in H^2)$$

where $C > 0$ is a positive number independent of f . This result has been generalized to many other Hilbert spaces of analytic functions. What are the Carleson measures for the model spaces? A discussion of these measures along with historical references can be found in [17].

13.2. Vector-valued model spaces. One can form the vector-valued Hardy space H^2 of \mathbb{C}^n -valued analytic functions on \mathbb{D} for which the integral means

$$\int_{\mathbb{T}} \|f(r\zeta)\|_{\mathbb{C}^n}^2 dm(\zeta)$$

are bounded as $r \rightarrow 1^-$. For a matrix valued inner function U on \mathbb{D} (i.e., a matrix-valued analytic function $U(z)$ on \mathbb{D} that satisfies $\|U\| \leq 1$ on \mathbb{D} and which is unitary valued a.e. on \mathbb{T}), we can define the \mathbb{C}^n -valued model space $H^2 \ominus UH^2$. The compression of the shift to these model spaces are the model operators for contractions via a similar Sz.-Nagy-Foiaş theory

for higher defect indices. Some well-known sources for all this operator theory are [14, 67, 69, 75]. There is even a Clark theory for these vector-valued model spaces including a version of the Aleksandrov disintegration theorem (Theorem 8.10) [36, 65].

13.3. Hankel operators. There are strong connections between model spaces and Hankel operators. The standard texts on the subject are [68, 69, 71].

13.4. Extremal problems. There is a nice connection between model spaces, truncated Toeplitz operators, and classical extremal problems. For a rational function ψ whose poles are in \mathbb{D} , a classical problem in complex analysis is to compute the quantity

$$\sup_{f \in H_1^1} \left| \frac{1}{2\pi i} \oint_{\mathbb{T}} \psi(\zeta) f(\zeta) d\zeta \right|, \quad (13.1)$$

along with the functions f for which the above supremum is achieved. In the above definition, H_1^1 denotes the unit ball in the Hardy space H^1 . These types of problems and extensions of them, where ψ belongs to L^∞ and is not assumed to be rational, have been studied since the early twentieth century. We refer the reader to [49] for a survey of these classical results. In that same paper is the following result:

Theorem 13.2. *If ψ is a rational function whose poles are contained in \mathbb{D} then*

$$\sup_{f \in H_1^1} \left| \frac{1}{2\pi i} \oint_{\mathbb{T}} \psi(\zeta) f(\zeta) d\zeta \right| = \sup_{g \in H_2^2} \left| \frac{1}{2\pi i} \oint_{\mathbb{T}} \psi(\zeta) g^2(\zeta) d\zeta \right|. \quad (13.3)$$

Furthermore, if u is the finite Blaschke product whose zeros are precisely those poles with corresponding multiplicities and if $\varphi := u\psi$, then

$$\sup_{f \in H_1^1} \left| \frac{1}{2\pi i} \oint_{\mathbb{T}} \psi(\zeta) f(\zeta) d\zeta \right| = \|A_\varphi^u\|,$$

where A_φ^u is the analytic truncated Toeplitz operator on \mathcal{K}_u with symbol φ .

In fact, the linear and quadratic supremums in (13.3) are the same for general ψ in $L^\infty(\mathbb{T})$ [23].

REFERENCES

1. Evgeny V. Abakumov, *Cyclicity and approximation by lacunary power series*, Michigan Math. J. **42** (1995), no. 2, 277–299. MR 1342491 (96f:47056)
2. J. Agler and J. E. McCarthy, *Pick interpolation and Hilbert function spaces*, Graduate Studies in Mathematics, vol. 44, American Mathematical Society, Providence, RI, 2002. MR 1882259 (2003b:47001)
3. P. R. Ahern and D. N. Clark, *On functions orthogonal to invariant subspaces*, Acta Math. **124** (1970), 191–204. MR 0264385 (41 #8981a)
4. ———, *Radial limits and invariant subspaces*, Amer. J. Math. **92** (1970), 332–342. MR 0262511 (41 #7117)
5. A. B. Aleksandrov, *Multiplicity of boundary values of inner functions*, Izv. Akad. Nauk Armyan. SSR Ser. Mat. **22** (1987), no. 5, 490–503, 515. MR 931885 (89e:30058)

6. ———, *Inner functions and related spaces of pseudocontinuable functions*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **170** (1989), no. Issled. Linein. Oper. Teorii Funktsii. 17, 7–33, 321. MR 1039571 (91c:30063)
7. ———, *On the existence of angular boundary values of pseudocontinuable functions*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **222** (1995), no. Issled. po Linein. Oper. i Teor. Funktsii. 23, 5–17, 307. MR 1359992 (97a:30046)
8. ———, *Gap series and pseudocontinuations. An arithmetic approach*, Algebra i Analiz **9** (1997), no. 1, 3–31. MR 1458416 (98d:30006)
9. ———, *Embedding theorems for coinvariant subspaces of the shift operator. II*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **262** (1999), no. Issled. po Linein. Oper. i Teor. Funkts. 27, 5–48, 231. MR 1734326 (2001g:46047)
10. Alexandru Aleman, Nathan S. Feldman, and William T. Ross, *The Hardy space of a slit domain*, Frontiers in Mathematics, Birkhäuser Verlag, Basel, 2009. MR 2548414 (2011m:30095)
11. Alexandru Aleman and Stefan Richter, *Simply invariant subspaces of H^2 of some multiply connected regions*, Integral Equations Operator Theory **24** (1996), no. 2, 127–155. MR 1371943 (99b:47010a)
12. N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337–404. MR 0051437 (14,479c)
13. Anton Baranov and Konstantin Dyakonov, *The Feichtinger conjecture for reproducing kernels in model subspaces*, J. Geom. Anal. **21** (2011), no. 2, 276–287. MR 2772073 (2012a:30130)
14. H. Bercovici, *Operator theory and arithmetic in H^∞* , Mathematical Surveys and Monographs, vol. 26, American Mathematical Society, Providence, RI, 1988. MR 954383 (90e:47001)
15. Robert D. Berman and William S. Cohn, *Tangential limits of Blaschke products and functions of bounded mean oscillation*, Illinois J. Math. **31** (1987), no. 2, 218–239. MR 882111 (88g:30036)
16. Arne Beurling, *On two problems concerning linear transformations in Hilbert space*, Acta Math. **81** (1948), 17. MR 0027954 (10,381e)
17. A. Blandigneres, E. Fricain, F. Gaunard, A. Hartmann, and W. T. Ross, *Reverse Carleson embeddings for model spaces*, To appear, J. London. Math. Soc.
18. Albrecht Böttcher and Bernd Silbermann, *Analysis of Toeplitz operators*, second ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2006, Prepared jointly with Alexei Karlovich. MR 2223704 (2007k:47001)
19. A. Brown and P. R. Halmos, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. **213** (1963/1964), 89–102. MR 0160136 (28 #3350)
20. G. T. Cargo, *Angular and tangential limits of Blaschke products and their successive derivatives*, Canad. J. Math. **14** (1962), 334–348. MR 0136743 (25 #204)
21. Lennart Carleson, *An interpolation problem for bounded analytic functions*, Amer. J. Math. **80** (1958), 921–930. MR 0117349 (22 #8129)
22. ———, *Interpolations by bounded analytic functions and the corona problem*, Ann. of Math. (2) **76** (1962), 547–559. MR 0141789 (25 #5186)
23. I. Chalendar, E. Fricain, and D. Timotin, *On an extremal problem of Garcia and Ross*, Oper. Matrices **3** (2009), no. 4, 541–546. MR 2597679 (2011b:30130)
24. N. Chevrot, E. Fricain, and D. Timotin, *The characteristic function of a complex symmetric contraction*, Proc. Amer. Math. Soc. **135** (2007), no. 9, 2877–2886 (electronic). MR 2317964 (2008c:47025)
25. Ole Christensen, *An introduction to frames and Riesz bases*, Applied and Numerical Harmonic Analysis, Birkhäuser Boston Inc., Boston, MA, 2003. MR 1946982 (2003k:42001)
26. J. A. Cima, A. L. Matheson, and W. T. Ross, *The Cauchy transform*, Mathematical Surveys and Monographs, vol. 125, American Mathematical Society, Providence, RI, 2006. MR 2215991 (2006m:30003)

27. J. A. Cima and W. T. Ross, *The backward shift on the Hardy space*, Mathematical Surveys and Monographs, vol. 79, American Mathematical Society, Providence, RI, 2000. MR 1761913 (2002f:47068)
28. Joseph A. Cima, Stephan Ramon Garcia, William T. Ross, and Warren R. Wogen, *Truncated Toeplitz operators: spatial isomorphism, unitary equivalence, and similarity*, Indiana Univ. Math. J. **59** (2010), no. 2, 595–620. MR 2648079 (2011i:47035)
29. D. N. Clark, *One dimensional perturbations of restricted shifts*, J. Analyse Math. **25** (1972), 169–191.
30. R. B. Crofoot, *Multipliers between invariant subspaces of the backward shift*, Pacific J. Math. **166** (1994), no. 2, 225–246. MR 1313454 (95k:47009)
31. R. G. Douglas, H. S. Shapiro, and A. L. Shields, *Cyclic vectors and invariant subspaces for the backward shift operator.*, Ann. Inst. Fourier (Grenoble) **20** (1970), no. fasc. 1, 37–76. MR 0270196 (42 #5088)
32. Ronald G. Douglas, *Banach algebra techniques in operator theory*, Academic Press, New York, 1972, Pure and Applied Mathematics, Vol. 49. MR 0361893 (50 #14335)
33. P. L. Duren, *Theory of H^p spaces*, Academic Press, New York, 1970.
34. K. Dyakonov and D. Khavinson, *Smooth functions in star-invariant subspaces*, Recent advances in operator-related function theory, Contemp. Math., vol. 393, Amer. Math. Soc., Providence, RI, 2006, pp. 59–66. MR 2198372 (2007b:30069)
35. Konstantin M. Dyakonov, *Kernels of Toeplitz operators via Bourgain’s factorization theorem*, J. Funct. Anal. **170** (2000), no. 1, 93–106. MR 1736197 (2000m:47036)
36. Sam Elliott, *A matrix-valued Aleksandrov disintegration theorem*, Complex Anal. Oper. Theory **4** (2010), no. 2, 145–157. MR 2652521 (2011g:46076)
37. E. Fabry, *Sur les séries de Taylor qui ont une infinité de points singuliers*, Acta Math. **22** (1898–1899), 65–88.
38. O. Frostman, *Sur les produits de Blaschke*, Kungl. Fysiografiska Sällskapetets i Lund Föreläsningar [Proc. Roy. Physiog. Soc. Lund] **12** (1942), no. 15, 169–182. MR 0012127 (6,262e)
39. S. R. Garcia, *Conjugation and Clark operators*, Recent advances in operator-related function theory, Contemp. Math., vol. 393, Amer. Math. Soc., Providence, RI, 2006, pp. 67–111. MR 2198373 (2007b:47073)
40. S. R. Garcia and M. Putinar, *Complex symmetric operators and applications*, Trans. Amer. Math. Soc. **358** (2006), no. 3, 1285–1315 (electronic). MR 2187654 (2006j:47036)
41. ———, *Complex symmetric operators and applications. II*, Trans. Amer. Math. Soc. **359** (2007), no. 8, 3913–3931 (electronic). MR 2302518 (2008b:47005)
42. S. R. Garcia and W. R. Wogen, *Complex symmetric partial isometries*, J. Funct. Anal. **257** (2009), no. 4, 1251–1260. MR 2535469
43. ———, *Some new classes of complex symmetric operators*, Trans. Amer. Math. Soc. **362** (2010), no. 11, 6065–6077. MR 2661508
44. Stephan R. Garcia and Mihai Putinar, *Interpolation and complex symmetry*, Tohoku Math. J. (2) **60** (2008), no. 3, 423–440. MR 2453732 (2009k:47048)
45. Stephan Ramon Garcia, *A $*$ -closed subalgebra of the Smirnov class*, Proc. Amer. Math. Soc. **133** (2005), no. 7, 2051–2059 (electronic). MR 2137871 (2005m:30040)
46. ———, *Conjugation, the backward shift, and Toeplitz kernels*, J. Operator Theory **54** (2005), no. 2, 239–250. MR 2186351 (2006g:30055)
47. ———, *Inner matrices and Darlington synthesis*, Methods Funct. Anal. Topology **11** (2005), no. 1, 37–47. MR 2130935 (2005m:30039)
48. Stephan Ramon Garcia, Daniel E. Poore, and William T. Ross, *Unitary equivalence to a truncated Toeplitz operator: analytic symbols*, Proc. Amer. Math. Soc. **140** (2012), no. 4, 1281–1295. MR 2869112 (2012m:47004)
49. Stephan Ramon Garcia and William T. Ross, *A non-linear extremal problem on the Hardy space*, Comput. Methods Funct. Theory **9** (2009), no. 2, 485–524. MR 2572653 (2010k:47062)

50. ———, *Recent progress on truncated Toeplitz operators, Blaschke products and their applications*, Fields Inst. Commun., vol. 65, Springer, New York, 2013, pp. 275–319. MR 3052299
51. Stephan Ramon Garcia and Donald Sarason, *Real outer functions*, Indiana Univ. Math. J. **52** (2003), no. 6, 1397–1412. MR 2021044 (2004k:30129)
52. J. Garnett, *Bounded analytic functions*, first ed., Graduate Texts in Mathematics, vol. 236, Springer, New York, 2007. MR 2261424 (2007e:30049)
53. J. Hadamard, *Essai sur l'étude des fonctions données par leur développement de Taylor*, J. Math. **8** (1892), 101–186.
54. Andreas Hartmann and William T. Ross, *Bad boundary behavior in star invariant subspaces II*, Ann. Acad. Sci. Fenn. Math. **37** (2012), no. 2, 467–478. MR 2987079
55. Henry Helson, *Large analytic functions. II*, Analysis and partial differential equations, Lecture Notes in Pure and Appl. Math., vol. 122, Dekker, New York, 1990, pp. 217–220. MR 1044789 (92c:30039)
56. D. Hitt, *Invariant subspaces of H^2 of an annulus*, Pacific J. Math. **134** (1988), no. 1, 101–120. MR 953502 (90a:46059)
57. K. Hoffman, *Banach spaces of analytic functions*, Prentice-Hall Series in Modern Analysis, Prentice-Hall Inc., Englewood Cliffs, N. J., 1962. MR 0133008 (24 #A2844)
58. S. V. Hruščëv, N. K. Nikol'skiĭ, and B. S. Pavlov, *Unconditional bases of exponentials and of reproducing kernels*, Complex analysis and spectral theory (Leningrad, 1979/1980), Lecture Notes in Math., vol. 864, Springer, Berlin, 1981, pp. 214–335. MR 643384 (84k:46019)
59. P. Koosis, *Introduction to H_p spaces*, second ed., Cambridge Tracts in Mathematics, vol. 115, Cambridge University Press, Cambridge, 1998, With two appendices by V. P. Havin [Viktor Petrovich Khavin]. MR 1669574 (2000b:30052)
60. T. L. Kriete, III, *On the Fourier coefficients of outer functions*, Indiana Univ. Math. J. **20** (1970/71), 147–155. MR 0412429 (54 #555)
61. ———, *A generalized Paley-Wiener theorem*, J. Math. Anal. Appl. **36** (1971), 529–555. MR 0288275 (44 #5473)
62. A. J. Lohwater and G. Piranian, *The boundary behavior of functions analytic in a disk*, Ann. Acad. Sci. Fenn. Ser. A. I. **1957** (1957), no. 239, 17. MR 0091342 (19,950c)
63. N. Lusin and J. Priwaloff, *Sur l'unicité et la multiplicité des fonctions analytiques*, Ann. Sci. École Norm. Sup. (3) **42** (1925), 143–191. MR 1509265
64. N. Makarov and A. Poltoratski, *Meromorphic inner functions, Toeplitz kernels and the uncertainty principle*, Perspectives in analysis, Math. Phys. Stud., vol. 27, Springer, Berlin, 2005, pp. 185–252.
65. R. T. W. Martin, *Unitary perturbations of compressed n -dimensional shifts*, Complex Anal. Oper. Theory **7** (2013), no. 4, 765–799. MR 3079829
66. Rubén A. Martínez-Avenidaño and Peter Rosenthal, *An introduction to operators on the Hardy-Hilbert space*, Graduate Texts in Mathematics, vol. 237, Springer, New York, 2007. MR 2270722 (2007k:47051)
67. N. Nikolski, *Treatise on the shift operator*, Springer-Verlag, Berlin, 1986.
68. ———, *Operators, functions, and systems: an easy reading. Vol. 1*, Mathematical Surveys and Monographs, vol. 92, American Mathematical Society, Providence, RI, 2002, Hardy, Hankel, and Toeplitz, Translated from the French by Andreas Hartmann. MR 1864396 (2003i:47001a)
69. ———, *Operators, functions, and systems: an easy reading. Vol. 2*, Mathematical Surveys and Monographs, vol. 93, American Mathematical Society, Providence, RI, 2002, Model operators and systems, Translated from the French by Andreas Hartmann and revised by the author. MR 1892647 (2003i:47001b)
70. V. Paulsen, *An introduction to the theory of reproducing kernel Hilbert spaces*, www.math.uh.edu/~vern/rkhs.pdf, 2009.
71. V. V. Peller, *Hankel operators and their applications*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003. MR 1949210 (2004e:47040)

72. A. Poltoratski and D. Sarason, *Aleksandrov-Clark measures*, Recent advances in operator-related function theory, Contemp. Math., vol. 393, Amer. Math. Soc., Providence, RI, 2006, pp. 1–14. MR 2198367 (2006i:30048)
73. A. G. Poltoratskiĭ, *Boundary behavior of pseudocontinuable functions*, Algebra i Analiz **5** (1993), no. 2, 189–210. MR 1223178 (94k:30090)
74. Frigyes Riesz and Béla Sz.-Nagy, *Functional analysis*, Frederick Ungar Publishing Co., New York, 1955, Translated by Leo F. Boron. MR 0071727 (17,175i)
75. M. Rosenblum and J. Rovnyak, *Hardy classes and operator theory*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1985, Oxford Science Publications. MR 822228 (87e:47001)
76. W. T. Ross and H. S. Shapiro, *Generalized analytic continuation*, University Lecture Series, vol. 25, American Mathematical Society, Providence, RI, 2002. MR 1895624 (2003h:30003)
77. Eero Saksman, *An elementary introduction to Clark measures*, Topics in complex analysis and operator theory, Univ. Málaga, Málaga, 2007, pp. 85–136. MR 2394657 (2009g:47041)
78. D. Sarason, *A remark on the Volterra operator*, J. Math. Anal. Appl. **12** (1965), 244–246. MR 0192355 (33 #580)
79. ———, *Generalized interpolation in H^∞* , Trans. Amer. Math. Soc. **127** (1967), 179–203. MR 0208383 (34 #8193)
80. ———, *Nearly invariant subspaces of the backward shift*, Contributions to operator theory and its applications (Mesa, AZ, 1987), Oper. Theory Adv. Appl., vol. 35, Birkhäuser, Basel, 1988, pp. 481–493. MR 1017680 (90m:47012)
81. ———, *Kernels of Toeplitz operators*, Toeplitz operators and related topics (Santa Cruz, CA, 1992), Oper. Theory Adv. Appl., vol. 71, Birkhäuser, Basel, 1994, pp. 153–164. MR 1300218 (95k:47039)
82. ———, *Sub-Hardy Hilbert spaces in the unit disk*, University of Arkansas Lecture Notes in the Mathematical Sciences, 10, John Wiley & Sons Inc., New York, 1994, A Wiley-Interscience Publication. MR 1289670 (96k:46039)
83. ———, *Algebraic properties of truncated Toeplitz operators*, Oper. Matrices **1** (2007), no. 4, 491–526. MR 2363975 (2008i:47060)
84. Harold S. Shapiro, *Weakly invertible elements in certain function spaces, and generators in ℓ_1* , Michigan Math. J. **11** (1964), 161–165. MR 0166343 (29 #3620)
85. ———, *Generalized analytic continuation*, Symposia on Theoretical Physics and Mathematics, Vol. 8 (Symposium, Madras, 1967), Plenum, New York, 1968, pp. 151–163. MR 0241614 (39 #2953)
86. ———, *Functions nowhere continuable in a generalized sense*, Publ. Ramanujan Inst. No. **1** (1968/1969), 179–182. MR 0267080 (42 #1982)
87. Joel H. Shapiro, *Composition operators and classical function theory*, Universitext: Tracts in Mathematics, Springer-Verlag, New York, 1993. MR 1237406 (94k:47049)
88. N. A. Shirokov, *Ideals and factorization in algebras of analytic functions that are smooth up to the boundary*, Trudy Mat. Inst. Steklov. **130** (1978), 196–222, 223, Spectral theory of functions and operators. MR 505686 (80j:46085)
89. ———, *Division and multiplication by inner functions in spaces of analytic functions smooth up to the boundary*, Complex analysis and spectral theory (Leningrad, 1979/1980), Lecture Notes in Math., vol. 864, Springer, Berlin, 1981, pp. 413–439. MR 643387 (83c:30028)
90. B. Simon, *Spectral analysis of rank one perturbations and applications*, Mathematical quantum theory. II. Schrödinger operators (Vancouver, BC, 1993), CRM Proc. Lecture Notes, vol. 8, Amer. Math. Soc., Providence, RI, 1995, pp. 109–149. MR 1332038 (97c:47008)
91. B. Simon and T. Wolff, *Singular continuous spectrum under rank one perturbations and localization for random Hamiltonians*, Comm. Pure Appl. Math. **39** (1986), no. 1, 75–90. MR 820340 (87k:47032)
92. E. Strouse, D. Timotin, and M. Zarrabi, *Unitary equivalence to truncated Toeplitz operators*, Indiana Univ. Math. J. **61** (2012), no. 2, 525–538. MR 3043586

93. Béla Sz.-Nagy, Ciprian Foias, Hari Bercovici, and László Kérchy, *Harmonic analysis of operators on Hilbert space*, second ed., Universitext, Springer, New York, 2010. MR 2760647 (2012b:47001)
94. S. Takenaka, *On the orthonormal functions and a new formula of interpolation*, Jap. J. Math. **2** (1925), 129 – 145.
95. Gerald Teschl, *Mathematical methods in quantum mechanics*, Graduate Studies in Mathematics, vol. 99, American Mathematical Society, Providence, RI, 2009, With applications to Schrödinger operators. MR 2499016 (2010h:81002)
96. G. C. Tumarkin, *Description of a class of functions admitting an approximation by fractions with preassigned poles*, Izv. Akad. Nauk Armjan. SSR Ser. Mat. **1** (1966), no. 2, 89–105. MR 34 #6123

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, CLAREMONT, CALIFORNIA, 91711,
USA

E-mail address: Stephan.Garcia@pomona.edu

URL: <http://pages.pomona.edu/~sg064747>

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF RICHMOND,
RICHMOND, VIRGINIA, 23173, USA

E-mail address: wross@richmond.edu

URL: <http://facultystaff.richmond.edu/~wross>