

2003

UreR, the Transcriptional Activator of the *Proteus mirabilis* Urease Gene Cluster, Is Required for Urease Activity and Virulence in Experimental Urinary Tract Infections

Jonathan D. Dattelbaum
University of Richmond, jdattalb@richmond.edu

C. Virginia Locketell

David E. Johnson

Harry L.T. Mobley

Follow this and additional works at: <http://scholarship.richmond.edu/chemistry-faculty-publications>

 Part of the [Biochemistry Commons](#), and the [Chemistry Commons](#)

Recommended Citation

Dattelbaum, Jonathan D., C. Virginia Locketell, David E. Johnson, and Harry L. T. Mobley. "UreR, the Transcriptional Activator of the *Proteus Mirabilis* Urease Gene Cluster, Is Required for Urease Activity and Virulence in Experimental Urinary Tract Infections." *Infection and Immunity* 71, no. 2 (2003): 1026-030. doi:10.1128/IAI.71.2.1026-1030.2003.

This Article is brought to you for free and open access by the Chemistry at UR Scholarship Repository. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact scholarshiprepository@richmond.edu.

UreR, the Transcriptional Activator of the *Proteus mirabilis* Urease Gene Cluster, Is Required for Urease Activity and Virulence in Experimental Urinary Tract Infections

Jonathan D. Dattelbaum,¹ C. Virginia Locketell,² David E. Johnson,^{2,3}
and Harry L. T. Mobley^{1*}

Department of Microbiology and Immunology¹ and Division of Infectious Disease, Department of Medicine,²
University of Maryland School of Medicine, and Veterans Affairs Medical Center,³
Baltimore, Maryland 21201

Received 24 September 2002/Returned for modification 22 October 2002/Accepted 31 October 2002

***Proteus mirabilis*, a cause of complicated urinary tract infection, produces urease, an essential virulence factor for this species. UreR, a member of the AraC/XylS family of transcriptional regulators, positively activates expression of the *ure* gene cluster in the presence of urea. To specifically evaluate the contribution of UreR to urease activity and virulence in the urinary tract, a *ureR* mutation was introduced into *P. mirabilis* HI4320 by homologous recombination. The isogenic *ureR::aphA* mutant, deficient in UreR production, lacked measurable urease activity. Expression was not detected in the UreR-deficient strain by Western blotting with monoclonal antibodies raised against UreD. Urease activity and UreD expression were restored by complementation of the mutant strain with *ureR* expressed from a low-copy-number plasmid. Virulence was assessed by transurethral cochallenge of CBA mice with wild-type and mutant strains. The isogenic *ureR::aphA* mutant of HI4320 was outcompeted in the urine ($P = 0.004$), bladder ($P = 0.016$), and kidneys ($P \leq 0.001$) 7 days after inoculation. Thus, UreR is required for basal urease activity in the absence of urea, for induction of urease by urea, and for virulence of *P. mirabilis* in the urinary tract.**

Proteus mirabilis commonly infects the urinary tracts of individuals with structurally abnormal urinary tracts or long-term indwelling catheters (20). Many virulence factors have been identified from this organism, including several types of fimbriae (1, 17, 18), hemolysin (28), flagella (21), and immunoglobulin A (IgA) protease (26, 30). Importantly, *P. mirabilis* produces a urea-inducible urease, which hydrolyzes urea into ammonia and ultimately carbon dioxide. As a result of ammonia production, an increase in local pH causes precipitation of normally soluble calcium and magnesium ions. These salt crystals can grow to remarkable size to produce bladder and kidney stones (16), which are a hallmark of infections with *Proteus* spp. (6).

We have shown previously that urease is an essential virulence factor in *P. mirabilis* urinary tract infections (UTIs) (11, 12). A urease-negative mutant was constructed by using homologous recombination to disrupt *ureC*, the main urease structural subunit. The urinary tracts of mice infected with the *P. mirabilis* isogenic urease mutant were colonized with significantly fewer numbers than those of mice infected with the parental strain (12). This mutant also demonstrated a reduced ability to persist after a 1- or 2-week infection and did not promote bladder and kidney stone formation (11).

The *P. mirabilis* urease gene cluster consists of *ureABC*, which encode the apoenzyme structural subunits, and *ureDEFG*, which encode proteins that facilitate insertion of the essential nickel ions into the catalytic site (22). Urease operon

expression is positively activated by UreR, a member of the AraC/XylS family of transcriptional regulators (3, 9, 23). UreR, a dimer of identical 293-amino-acid polypeptides (24), binds urea (5), causing the protein to bind avidly to both the *ureR* and *ureD* promoters within the 491-bp *ureR-ureD* intergenic region (27). RNA polymerase is recruited by UreR, thus activating transcription (27). To date, all of the work elucidating the functional role of UreR activation of the urease gene cluster has focused on UreR in vitro by using reporter constructs expressed in *Escherichia coli*. Since UreR controls expression of the entire gene cluster, we investigated how UreR function contributes to basal urease production and virulence of *P. mirabilis* in a mouse model of ascending UTI.

P. mirabilis HI4320, originally isolated from an elderly (>65 years old) woman with urinary catheter-associated bacteriuria, possesses urea-inducible urease activity (13, 29). *E. coli* DH5 α was used as the host strain for transformation of plasmids other than pGP704 (19) and derivatives which require *E. coli* DH5 α *pir*. Plasmid pMID1010 carries the entire *P. mirabilis* gene cluster cloned into pBR322 (14). Strains were maintained on Luria broth supplemented with ampicillin (100 μ g/ml) or kanamycin (50 μ g/ml) when necessary. Nonswarming agar (per liter, 10 g of tryptone, 5 g of yeast extract, 0.5 g of NaCl, and 15 g of agar, with 0.5% glycerol) was used to prevent *P. mirabilis* swarming when cultured on plates (2).

Construction of the *ureR::aphA* mutation in *P. mirabilis* HI4320. To determine the contribution of *ureR* to urease expression in vivo, we constructed a *P. mirabilis* strain deficient in UreR production according to the scheme shown in Fig. 1. The *ureR* open reading frame was subcloned and insertionally inactivated with a kanamycin resistance cassette. The fragment containing disrupted *ureR* was cloned into a pGP704-based

* Corresponding author. Mailing address: Department of Microbiology and Immunology, University of Maryland School of Medicine, 655 W. Baltimore St., Baltimore, MD 21201. Phone: (410) 706-0466. Fax: (410) 706-6751. E-mail: hmobley@umaryland.edu.




FIG. 1. Construction of a *P. mirabilis* *ureR* mutant. (A) A 2.535-kb *BsmI* fragment from pMID1010 encompassing most of *ureR*, *ureD*, *ureA* and the first 51 nucleotides of *ureB* was treated with T4 DNA polymerase to fill in overhanging ends and render the fragment blunt and cloned into the *SmaI* site of pBluescript SK+ treated with shrimp alkaline phosphatase to create pJD10. (B) After digestion of pJD10 with *NruI* (corresponding to codon 25 of *ureR*) and shrimp alkaline phosphatase treatment, a blunt-ended kanamycin resistance cassette (*aphA*), excised from pUC-4 κ by digestion with *HincII*, was inserted into pJD10. Kanamycin- and ampicillin-resistant versions of DH5 α harboring the new construct, pJD11, were isolated. Arrows represent the direction of transcription. (C) A 3.8-kb *XbaI-KpnI* fragment containing the entire insert region of pJD11 was excised from an agarose gel (*XmnI* digestion was also used to remove the pBluescript vector from the desired fragment) and subcloned into pGP704 (19) to create the suicide construct pJD12. The suicide construct pJD12 was electroporated into *P. mirabilis* HI4320, and kanamycin-resistant transformants were selected on nonswarming agar.

suicide vector as described in the legend to Fig. 1. The resulting suicide construct, pJD12, was electroporated into *P. mirabilis* HI4320. To verify that a double crossover event had occurred, Kan^r colonies were screened for ampicillin sensitivity; 4 of 525 colonies were identified as having appropriate characteristics (Fig. 1C). Homologous recombination of the wild-type *ureR* with *ureR::aphA* was confirmed by PCR by using chromosomal DNA and primers specific for *ureR*. A band of approximately 0.9 kb was amplified in the parent strain, while a band of approximately 2.2 kb (*ureR* plus Kan^r cassette) was observed in the Kan^r Amp^s mutant strain (data not shown). This mutant, designated *P. mirabilis* HI4320 *ureR::aphA*, was found to swarm normally and had a growth rate identical to that of the wild type (data not shown).

Urease activity in the *ureR::aphA* mutant. Since UreR functions as the transcriptional activator of the urease gene cluster, urease activity was quantified in vitro for *P. mirabilis* HI4320 and the isogenic *ureR::aphA* mutant (Fig. 2). The wild-type strain displayed a 5.7-fold increase in urease activity after culture in the presence of 100 mM urea. Conversely, although the level of detection for the urease assay is quite sensitive (approximately 10 nmol of ammonia), urease activity was undetectable for the mutant strain under identical experimental conditions.

Complementation of the *ureR* mutation. To determine whether urease activity could be restored to the *P. mirabilis* *ureR* mutant by complementation, *ureR* was cloned into the *EcoRV-BamHI* site of pACYC184 and was expressed from the tetracycline promoter. Following electroporation of the construct into the *ureR* mutant strain, urease activity was measured and the production of UreD was assayed by Western blotting (Fig. 2). Complementation by *ureR* in *trans* resulted in wild-type levels of basal urease activity as well as a fourfold induction (~60% of that of the wild type) after culture in the presence of 100 mM urea. While *ureR* provided in multicopy restored urea inducibility, the level of induction was not statistically equivalent to that observed in the wild-type strain.

The data to date suggest that the genes encoding the urease structural subunits and accessory proteins (except *ureG*) are transcribed on a single polycistronic mRNA (4). Because UreR is a transcriptional activator, it is likely that only a few copies of the gene product are required for activation. It is therefore not surprising that UreR was undetectable on Western blots of either wild-type *P. mirabilis* or the isogenic *ureR* mutant (data not shown). To determine if expression of the urease operon occurred, we measured UreD production by using Western blotting for the parent, mutant, and complemented mutant (Fig. 2). As predicted from the presence of basal urease activity, *P. mirabilis* produced a small but detectable amount of UreD in the absence of urea. The amount of UreD increased significantly upon addition of 100 mM urea, which correlated with the fold increase observed for urease activity. Under identical experimental conditions, UreD expression was not detected in the *P. mirabilis* *ureR::aphA* mutant regardless of urea levels, whereas UreD expression was restored by complementation with cloned *ureR* expressed in *trans* from a plasmid. These data demonstrated that a fully functional *ure* gene cluster was present and that insertional inactivation of *ureR* did not exert polar effects on other *ure* genes. Importantly, it was revealed that UreR is responsible for the low basal levels of urease activity in uninduced *P. mirabilis*.

The *P. mirabilis* *ureR::aphA* mutant is outcompeted by the parent strain in a murine cochallenge model of UTI. The virulence of *P. mirabilis* parent and mutant strains was assessed in a murine cochallenge model (15) by using a modification (10) of the procedure developed by Hagberg et al. (7). For cochallenge experiments, 10 6- to 8-week-old female CBA mice (Harlan Sprague Dawley, Indianapolis, Ind.) were infected transurethrally with an approximate 1:1 ratio (5×10^6 CFU of each strain) of the *P. mirabilis* parent and *ureR::aphA* mutant strains. Seven days after inoculation, urine was collected and mice were sacrificed. Quantitative counts from the urine, bladder, and kidneys were obtained by plating on nonswarming agar (2) to allow growth of both parent and mutant strains and on nonswarming agar with kanamycin (50 μ g/ml) to detect the mutant strain alone. The limit of detection for determination of viable bacteria was 10^2 CFU per milliliter of urine or per gram of tissue. In these studies, the mutant was unable to colonize the bladder or kidneys in 8 of 10 mice (median value, 10^2 CFU, which is below the limit of detection) (Fig. 3). As a result of the inability to produce urease, the *ureR::aphA* mutant strain was dramatically outcompeted by the wild-type strain in the urine ($P = 0.004$), bladder ($P = 0.016$)




FIG. 2. *P. mirabilis* urease activity in wild-type HI4320, *ureR::aphA* mutant, and complemented strains. Urease activity (upper panel) was determined by using the phenol-hypochlorite reaction (31). Overnight cultures (5 ml) of the *P. mirabilis* wild-type strain HI4320, the *ureR* mutant (*ureR::aphA*), and the complemented strain (Comp.) were used to inoculate fresh medium (1:100) with or without 100 mM urea containing the appropriate antibiotics when necessary. Bacteria from mid-exponential-phase cultures (optical density at 600 nm, 0.4) (3 ml) were resuspended in 1 ml of 50 mM HEPES, pH 7.5, and lysed by passage through a French pressure cell at 20,000 lb/in². The protein concentration of crude extract was determined by the bicinchoninic acid assay (Pierce, Rockford, Ill.). Total protein (10 μg) was diluted into 1 ml of assay buffer (50 mM HEPES, pH 7.5, and 25 mM urea) and incubated for 20 min at 37°C. Ammonia production was measured by using the reaction of phenyl nitroprusside and alkali hypochlorite (31). Optical density was measured at 625 nm by using ammonium chloride as a standard. For Western blotting (lower panel), monoclonal anti-UreD serum was prepared from a murine hybridoma cell line as previously described (8). Crude extract prepared for urease assays was assessed for production of UreD. Protein was subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred overnight onto Immobilon-P membrane (Millipore, Inc.). Anti-UreD serum (1:100) and anti-mouse IgG conjugated with horseradish peroxidase (1:2,000) were used as primary and secondary antibodies, respectively. Protein bands were visualized on X-ray film by using chemiluminescence reagents from Amersham Biosciences (Piscataway, N.J.).

and kidneys ($P < 0.001$) as well as the urinary tract overall ($P < 0.001$).

UreR is required for basal urease activity in the absence of urea, for induction of urease by urea, and for virulence in the

urinary tract. Urease is an essential virulence factor for *P. mirabilis* colonization and persistence in UTI (11, 12). UreR, an AraC/XylS family transcriptional regulator, has been shown in vitro to positively activate *ure* gene cluster expression in a




FIG. 3. Cochallenge of *P. mirabilis* parent and *ureR::aphA* isogenic mutant strains in the CBA mouse model of ascending UTI. Mice were transurethrally infected with 10^7 CFU containing a 1:1 ratio of *P. mirabilis* HI4320 and *ureR* mutant strains. After a 7-day infection, urine, bladder, and kidney specimens were collected and the CFU per milliliter or per gram of tissue of parent and mutant strains was determined for each mouse. Each unique symbol refers to CFU for the wild type (WT) and the *ureR* mutant from one cochallenged mouse. The horizontal dashed line represents the lower limit of detection of viable bacteria. For ease of comparison, the median values (black bars) are connected by dotted lines. Statistical differences between the median number of log₁₀ CFU per milliliter or per gram of tissue for *P. mirabilis* HI4320 and its *ureR::aphA* mutant were determined by using the Wilcoxon rank sum test.

urea-dependent manner. To properly assess the role of UreR regulation activity in *P. mirabilis* in vivo, homologous recombination was used to construct a *P. mirabilis* mutant that was incapable of producing UreR. A kanamycin resistance gene, *aphA*, was inserted at the *ureR* locus in the *P. mirabilis* chromosome by homologous recombination with a pGP704-based suicide construct (Fig. 1). This mutant did not produce urease in the absence or in the presence of urea as measured by using the phenol-hypochlorite assay on mid-exponential-phase cultures (Fig. 2). We have known for many years that *P. mirabilis* produced basal levels of urease activity in the absence of exogenously added urea (14). Since there is a complete loss of basal urease activity in the *ureR* mutant, we conclude that in *P. mirabilis*, low levels of UreR may account for basal urease expression in the absence of urea. Urease activity was restored upon complementation with *ureR* expressed on a low-copy-number plasmid (Fig. 2). Interestingly, full wild-type levels of induction were not reached in the complemented mutant. It is possible that translational coupling plays a role in which newly translated UreR must bind immediately to the adjacent *ureR-ureD* intergenic region. UreR synthesized from the complementing plasmid would have to diffuse to the chromosomal target, an event that may be less efficient than the wild-type mechanism.

The isogenic *ureR::aphA* mutant of strain HI4320 was outcompeted by the parental strain in 10 of 10 CBA mice following transurethral infection (Fig. 3); indeed, in only one mouse were the numbers of the *ureR* mutant and parent strain similar. In our experience, as well as that of others (25), coinfection with parental and mutant strains within the same host represents a more sensitive method to test for attenuation of virulence. To our knowledge, this is the first report of a cochallenge experiment with *P. mirabilis* and a urease-negative mutant. We infected 10 CBA mice with a 1:1 mixture ($\sim 5 \times 10^6$ CFU of each strain) of the HI4320 parent and *ureR::aphA* mutant strains. The parental strain outcompeted the mutant strain in all mice tested (Fig. 3). The low fitness and complete attenuation of the mutant strain confirms the role of urease as an essential virulence factor for *P. mirabilis* and demonstrates that the urease of the wild-type strain is not sufficient to sustain the urease-negative mutant in the urinary tract. Interestingly, in a previously published study (12), the wild type and the urease-negative (*ureC*) mutant of *P. mirabilis* were independently inoculated into the bladders of mice. While the mutant colonized the urine, bladder, and kidneys in ~ 100 -fold fewer CFU per milliliter or per gram, this mutant was nevertheless able to colonize the urinary tract to some degree in the absence of a wild-type strain. The inability of the wild-type strain to support the cocolonization of the *ureR* mutant suggests that the alteration of urine pH by urease is not the sole action that provides advantage to *P. mirabilis*. It is likely that urease, a cytoplasmic protein, provides additional advantages, such as availability of ammonia for glutamine production and protein synthesis. These latter functions could not be provided to the mutant by the wild-type strain.

Cochallenge of CBA mice with both parent and mutant strains represents a powerful method to directly compare the requirement of urease production to *P. mirabilis* virulence in a single mouse. High levels (10^7 to 10^8 CFU/g of tissue) of the *P. mirabilis* wild type were recovered from a moribund mouse

(Fig. 3, black square) due to the formation of an obstructive bladder stone. Interestingly, the highest mutant counts were also observed for this mouse. Our lab has previously visualized *P. mirabilis* attached to the surfaces of bladder and kidney epithelium and within crevices of bladder and kidney stones (16). We hypothesize that residing close to the stone may help *P. mirabilis* evade the host immune response. This observation suggests that stone formation may not only contribute to *P. mirabilis* persistence in the urinary tract (11) but may also provide an advantage to nonureolytic pathogens as well.

To our knowledge, UreR has not been shown to affect the expression of any other operon besides the *ure* gene cluster. Since our data demonstrate that the absence of the UreR regulatory protein leads to inhibition of urease expression, this protein may serve as a possible target for treatment or eradication of *P. mirabilis* from the urinary tract. Further experimentation is required to determine if other *ureR*-positive *Enterobacteriaceae* family members will display a similar loss of urease activity following perturbation of *ureR*.

This work was supported by Public Health Service grants AI23328 and DK47920 from the National Institutes of Health. J.D.D. was funded by a National Research Service Award (F32 AI51078) from the NIH.

REFERENCES

- Bahrani, F. K., G. Massad, C. V. Lockett, D. E. Johnson, R. G. Russell, J. W. Warren, and H. L. Mobley. 1994. Construction of an MR/P fimbrial mutant of *Proteus mirabilis*: role in virulence in a mouse model of ascending urinary tract infection. *Infect. Immun.* **62**:3363–3371.
- Belas, R., D. Erskine, and D. Flaherty. 1991. Transposon mutagenesis in *Proteus mirabilis*. *J. Bacteriol.* **173**:6289–6293.
- D'Orazio, S. E. F., and C. M. Collins. 1993. The plasmid-encoded urease gene cluster of the family *Enterobacteriaceae* is positively regulated by UreR, a member of the AraC family of transcriptional activators. *J. Bacteriol.* **175**:3459–3467.
- D'Orazio, S. E. F., and C. M. Collins. 1995. UreR activates transcription at multiple promoters within the plasmid-encoded urease locus of the *Enterobacteriaceae*. *Mol. Microbiol.* **16**:145–155.
- Gendlina, I., D. M. Gutman, V. Thomas, and C. M. Collins. 2002. Urea-dependent signal transduction by the virulence regulator UreR. *J. Biol. Chem.* **277**:37349–37358.
- Griffith, D. P., D. M. Musher, and C. Itin. 1976. Urease: the primary cause of infection-induced urinary stones. *Investig. Urol.* **13**:346–350.
- Hagberg, L., I. Engberg, R. Freter, J. Lam, S. Olling, and C. Svanborg-Eden. 1983. Ascending unobstructed urinary tract infection in mice caused by pyelonephritogenic *Escherichia coli* of human origin. *Infect. Immun.* **40**:273–283.
- Heimer, S. R., and H. L. T. Mobley. 2001. Interaction of *Proteus mirabilis* urease apoenzyme and accessory proteins identified with yeast two-hybrid technology. *J. Bacteriol.* **183**:1423–1433.
- Island, M. D., and H. L. T. Mobley. 1995. *Proteus mirabilis* urease: operon fusion and linker insertion analysis of *ure* gene organization, regulation, and function. *J. Bacteriol.* **177**:5653–5660.
- Johnson, D. E., C. V. Lockett, M. Hall-Craigs, H. L. T. Mobley, and J. W. Warren. 1987. Uropathogenicity in rats and mice of *Providencia stuartii* from long-term catheterized patients. *J. Urol.* **138**:632–635.
- Johnson, D. E., R. G. Russell, C. V. Lockett, J. C. Zulty, J. W. Warren, and H. L. T. Mobley. 1993. Contribution of *Proteus mirabilis* urease to persistence, urolithiasis, and acute pyelonephritis in a mouse model of ascending urinary tract infection. *Infect. Immun.* **61**:2748–2754.
- Jones, B. D., C. V. Lockett, D. E. Johnson, J. W. Warren, and H. L. T. Mobley. 1990. Construction of a urease-negative mutant of *Proteus mirabilis*: analysis of virulence in a mouse model of ascending urinary tract infection. *Infect. Immun.* **58**:1120–1123.
- Jones, B. D., and H. L. T. Mobley. 1987. Genetic and biochemical diversity of ureases of *Proteus*, *Providencia*, and *Morganella* species isolated from urinary tract infection. *Infect. Immun.* **55**:2198–2203.
- Jones, B. D., and H. L. T. Mobley. 1988. *Proteus mirabilis* urease: genetic organization, regulation, and expression of structural genes. *J. Bacteriol.* **170**:3342–3349.
- Li, X., D. E. Johnson, and H. L. T. Mobley. 1999. Requirement of MrpH for mannose-resistant *Proteus*-like fimbria-mediated hemagglutination by *Proteus mirabilis*. *Infect. Immun.* **67**:2822–2833.

16. Li, X., H. Zhao, C. V. Lockett, C. B. Drachenberg, D. E. Johnson, and H. L. Mobley. 2002. Visualization of *Proteus mirabilis* within the matrix of urease-induced bladder stones during experimental urinary tract infection. *Infect. Immun.* **70**:389–394.
17. Massad, G., F. K. Bahrani, and H. L. Mobley. 1994. *Proteus mirabilis* fimbriae: identification, isolation, and characterization of a new ambient-temperature fimbria. *Infect. Immun.* **62**:1989–1994.
18. Massad, G., C. V. Lockett, D. E. Johnson, and H. L. Mobley. 1994. *Proteus mirabilis* fimbriae: construction of an isogenic *pmfA* mutant and analysis of virulence in a CBA mouse model of ascending urinary tract infection. *Infect. Immun.* **62**:536–542.
19. Miller, V. L., and J. J. Mekalanos. 1988. A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *J. Bacteriol.* **170**:2575–2583.
20. Mobley, H. L. 1996. Virulence of *Proteus mirabilis*, p. 245–269. In H. L. Mobley and J. W. Warren (ed.), *Urinary tract infections: molecular pathogenesis and clinical management*. ASM Press, Washington, D.C.
21. Mobley, H. L., R. Belas, V. Lockett, G. Chippendale, A. L. Trifillis, D. E. Johnson, and J. W. Warren. 1996. Construction of a flagellum-negative mutant of *Proteus mirabilis*: effect on internalization by human renal epithelial cells and virulence in a mouse model of ascending urinary tract infection. *Infect. Immun.* **64**:5332–5340.
22. Mobley, H. L., M. D. Island, and R. P. Hausinger. 1995. Molecular biology of microbial ureases. *Microbiol. Rev.* **59**:451–480.
23. Nicholson, E. B., E. A. Concaugh, P. A. Foxall, M. D. Island, and H. L. Mobley. 1993. *Proteus mirabilis* urease: transcription and regulation by UreR. *J. Bacteriol.* **175**:465–473.
24. Poor, C. A., C. Coker, J. D. Dattelbaum, and H. L. T. Mobley. 2001. Identification of domains of UreR, an AraC-like transcriptional regulator of the urease gene cluster in *Proteus mirabilis*. *J. Bacteriol.* **183**:4526–4535.
25. Salama, N. R., G. Otto, L. Tompkins, and S. Falkow. 2001. Vacuolating cytotoxin of *Helicobacter pylori* plays a role during colonization in a mouse model of infection. *Infect. Immun.* **69**:730–736.
26. Senior, B. W., M. Albrechtsen, and M. A. Kerr. 1987. *Proteus mirabilis* strains of diverse type have IgA protease activity. *J. Med. Microbiol.* **24**:175–180.
27. Thomas, V. J., and C. M. Collins. 1999. Identification of UreR binding sites in the *Enterobacteriaceae* plasmid-encoded and *Proteus mirabilis* urease gene operons. *Mol. Microbiol.* **31**:1417–1428.
28. Toth, V., and L. Emody. 2000. Proteus virulence: involvement of the pore forming alpha-hemolysin (a short review). *Acta Microbiol. Immunol. Hung.* **47**:457–470.
29. Warren, J. W., J. H. Tenney, H. L. Hoopes, H. L. Muncie, and W. C. Anthony. 1982. A prospective microbiologic study of bacteriuria in patients with chronic indwelling urethral catheters. *J. Infect. Dis.* **146**: 719–723.
30. Wassif, C., D. Cheek, and R. Belas. 1995. Molecular analysis of a metalloprotease from *Proteus mirabilis*. *J. Bacteriol.* **177**:5790–5798.
31. Weatherburn, M. W. 1967. Phenol-hypochlorite reaction for determination of ammonia. *Anal. Chem.* **39**:971–974.

Editor: D. L. Burns