

2015

## One-pot silyl ketene acetal-formation-Mukaiyama–Mannich additions to imines mediated by trimethylsilyl trifluoromethanesulfonate

C. Wade Downey  
*University of Richmond, wdowney@richmond.edu*

Jared A. Ingersoll

Hadleigh M. Glist

Carolyn M. Dombrowski

Adam T. Barnett

Follow this and additional works at: <https://scholarship.richmond.edu/chemistry-faculty-publications>

 Part of the [Organic Chemistry Commons](#)

---

### Recommended Citation

C. Wade Downey, Jared A. Ingersoll,\* Hadleigh M. Glist,\* Carolyn M. Dombrowski,\* Adam T. Barnett.\* One-pot silyl ketene acetal-formation-Mukaiyama–Mannich additions to imines mediated by trimethylsilyl trifluoromethanesulfonate. *European Journal of Organic Chemistry* 2015, 7287-7291.

This Article is brought to you for free and open access by the Chemistry at UR Scholarship Repository. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact [scholarshiprepository@richmond.edu](mailto:scholarshiprepository@richmond.edu).

# One-pot silyl ketene acetal-formation-Mukaiyama–Mannich additions to imines mediated by trimethylsilyl trifluoromethanesulfonate

C. Wade Downey,<sup>\*[a]</sup> Jared A. Ingersoll,<sup>[a]</sup> Hadleigh M. Glist,<sup>[a]</sup> Carolyn M. Dombrowski,<sup>[a]</sup> and Adam T. Barnett<sup>[a]</sup>

Dedication ((optional))


---

[a] C. W. Downey, J. A. Ingersoll, H. M. Glist, C. M. Dombrowski, A. T. Barnett  
Department of Chemistry  
University of Richmond  
28 Westhampton Way  
Richmond, VA, USA 23173  
[wdowney@richmond.edu](mailto:wdowney@richmond.edu)  
<http://chemistry.richmond.edu/faculty/wdowney/>

Supporting information for this article is given via a link at the end of the document. ((Please delete this text if not appropriate))

---

**Abstract:** In the presence of trimethylsilyl trifluoromethanesulfonate and trialkylamine base, thioesters are readily converted to silyl ketene acetals in situ and undergo Mukaiyama–Mannich addition to *N*-phenylimines in one pot. The silyl triflates appears to play two roles, activating both the thioester and the imine. This process also works well when thioesters are replaced with amides, esters, or ketones. Products are isolated as desilylated anilines without the necessity of a deprotection step. Yields range from 65–99%.



## Introduction

The Mukaiyama–Mannich reaction occupies an honored position in the field of organic synthesis, providing a convergent route to complex  $\alpha$ -amino carbonyl compounds through a carbon-carbon bond-forming process.<sup>[1,2]</sup> Significant advances have been made in this field, particularly with respect to the development of enantioselective variants,<sup>[3,4]</sup> three-component coupling reactions,<sup>[5,6]</sup> and proton- or hydrogen bond-catalyzed reactions.<sup>[7]</sup> Each of these examples, however, relies upon the independent generation and purification of the silyl ketene acetal reaction partner, a mild nucleophile that usually requires activation of the imine acceptor with a Lewis acid in order to achieve the desired reactivity.

Previously, we reported the one-pot silyl ketene acetal-formation-Mukaiyama–Mannich additions of thioesters and ketones to nitrones in the presence of trimethylsilyl trifluoromethanesulfonate (TMSOTf) and an amine base (eq 1).<sup>[8,9]</sup> Because of the known affinity of silicon for Lewis basic oxygen species, nitrones were a logical target for our trialkylamine/TMSOTf system.<sup>[10]</sup> Activation at nitrogen, however, appeared to be a considerably greater challenge, because the nitrogen-silicon bond generated over the course of the reaction (~85 kcal/mol) is considerably weaker than a silicon-oxygen bond (~108 kcal/mol).<sup>[11]</sup> Nonetheless, the analogy to our previous work with *N*-phenylnitrones drew us to *N*-phenylaldimines as potential electrophiles. Furthermore, the *N*-monoalkylated aniline products share structural properties with a number of cardiovascular drugs such as methylclothiazide,<sup>[12]</sup> furosemide,<sup>[13]</sup> bumetanide,<sup>[14]</sup> and dabigatran etexilate<sup>[15]</sup> (Figure 1). We now describe the ability of TMSOTf to mediate the addition of thioesters, amides, esters, and ketones to *N*-phenylimines through a two-step, one-pot process.






Figure 1. Some aniline-containing pharmaceuticals



## Results and Discussion

Initial discovery and optimization experiments were performed with acetophenone and the *N*-phenylimine derived from benzaldehyde ((*E*)-*N*-benzylideneaniline), but generally inconsistent results (vide infra) prompted us to examine *S*-phenyl thioacetate as a more convenient enolate precursor. When the thioacetate was treated with TMSOTf, Et<sub>3</sub>N, and the imine in CH<sub>2</sub>Cl<sub>2</sub> at room temperature, high conversion (<90%) to the Mannich product was observed (eq 2). Interestingly, desilylation of the product appeared to proceed spontaneously under the reaction conditions. When CH<sub>2</sub>Cl<sub>2</sub> was substituted with other solvents (Et<sub>2</sub>O, toluene, THF, cyclopentyl methyl ether) much lower conversion was observed.



A plausible mechanism is presented in Figure 2. Enolization of the thioester occurs conveniently by treatment with TMSOTf and Et<sub>3</sub>N, a base that gave consistently better results than *i*-Pr<sub>2</sub>NEt in terms of reproducibility and overall conversion for thioester substrates.<sup>[16]</sup> Activation of the imine appears to be accomplished by residual TMSOTf (vide infra), accelerating the Mukaiyama–Mannich addition and presumably yielding an *N*-silylated aniline as the product. Net silyl transfer from oxygen to

another molecule of imine was facile as expected, but we were pleased to discover that spontaneous desilylation of the aniline nitrogen also occurs under the reaction conditions. This result stands in some contrast to literature precedent, which includes reports that *N*-alkylated anilines are readily silylated in the presence of TMSCl and Et<sub>3</sub>N (eq 3),<sup>[17]</sup> conditions quite analogous to those employed here. The presence of a carbonyl conveniently positioned to hydrogen bond with the free aniline proton may provide a thermodynamic driving force for the exchange of the silicon group with the acidic proton on the nascent ammonium salt.





Figure 2. Proposed Mechanistic Scheme



Given this apparent ability of the ammonium salt to act as a weak acid under the conditions, the possibility that it might be the activator of the imine warranted investigation. Brønsted acids have been shown to catalyze similar reactions through the protonation of nitrogen-based electrophiles, a motif recently demonstrated in aziridine openings<sup>[18]</sup> as well as Mukaiyama–Mannich additions<sup>[19]</sup> In order to investigate this possibility, a number of control experiments were performed (Table 1). When the independently prepared the silyl ketene acetal of *S*-phenyl thioacetate was treated with imine in methylene chloride, only a trace of product was observed after 2 h (entry 1). Addition of 1.0 equiv Et<sub>3</sub>N•HCl to the reaction mixture did accelerate the reaction, but only 16% conversion was observed after 2 h (entry 2), suggesting that ammonium salts are not likely to be the


active catalyst under our optimized conditions. In contrast, when the silyl ketene acetal and imine were treated with 1.0 equiv TMSOTf and stirred for 1 h, 89% conversion to the Mannich product was observed (entry 3). These results suggest that the most likely catalyst is a cationic silicon species, and that desilylation of a preliminary *N*-silylated product occurs spontaneously under the reaction conditions.

Table 1. Identification of Lewis acid catalyst

entry	catalyst	conv (%) <sup>[b]</sup>
1	none	trace
2	Et <sub>3</sub> N (1.0 equiv)	0
3	Et <sub>3</sub> N•HCl (1.0 equiv)	16
4	TMSOTf (1.0 equiv)	89

[a] Reaction conditions: 0.2 mmol silyl ketene acetal, 0.24 mmol imine, 0.2 mmol catalyst, 1.0 mL CH<sub>2</sub>Cl<sub>2</sub>, rt, 2 h. [b] Determined by <sup>1</sup>H NMR spectroscopy of the unpurified reaction mixture


A variety of imines was tested under the optimized reaction conditions, including *N*-tosyl, *N*-phosphoryl, *N*-alkyl, and *N*-(4-methoxy)phenyl imines, but only *N*-phenyl imines showed consistently high reactivity (Table 2). Other imines typically reacted with 0–50% conversion under optimized conditions. A wide range of aryl imines, including both electron-poor and electron-rich substrates, reacted in high yield with *S*-phenyl thioacetate (entries 1–6), except for the *p*-nitrobenzaldehyde derivative (entry 4). Initial conversion to product **1d** did appear to be quite high, affording >95% conversion as determined by <sup>1</sup>H NMR spectroscopy of the unpurified reaction mixture. The product was not stable to silica gel chromatography, however, and decomposed completely during attempted purification. Treatment of the silica gel with Et<sub>3</sub>N or Et<sub>2</sub>NH did not prevent decomposition, which appears to proceed primarily through retro-Mannich reaction as evidenced by recovery of the *S*-phenyl thioacetate. Nonetheless, other aryl aldimines were quite successful, including heteroaryl derivatives and the sterically challenging 2-naphthyl substrate (entries 7–9).

**Table 2.** Mannich reactions of *S*-phenyl thioacetate


entry	R	product	yield (%) <sup>[b]</sup>
1	Ph	<b>1a</b>	81
2	4-MePh	<b>1b</b>	81
3	4-MeOPh	<b>1c</b>	75
4	4-NO <sub>2</sub> Ph	<b>1d</b>	0 <sup>[c]</sup>
5	4-FPh	<b>1e</b>	87
6	4-BrPh	<b>1f</b>	83
7	2-naphthyl	<b>1g</b>	84
8	2-furyl	<b>1h</b>	93
9	2-thiophenyl	<b>1i</b>	87


[a] Reaction conditions: 1.0 mmol thioester, 1.2 mmol imine, 1.3 mmol TMSOTf, 1.2 mmol Et<sub>3</sub>N, 2.5 mL CH<sub>2</sub>Cl<sub>2</sub>, rt, 2 h. [b] Isolated yield after chromatography. [c] Product decomposed during purification on silica gel.

Extension to other thioesters was rapidly achieved (Table 3). Substitution on the aryl position of the thioester was easily tolerated (entries 1-3), including the sterically encumbered 2-naphthalenethiol derivative. Replacement of the aryl thiol-derived moiety with a benzyl mercaptyl group led to smooth reactivity, yielding the *S*-benzyl product **2d**. When the acetyl residue was replaced with a propionyl group, the typical high level of reactivity was maintained, providing product **3** in 98% yield (eq 4). The reaction rate did slow significantly, however, requiring overnight stirring to achieve optimal conversion. Diastereoselectivity was moderate but significant (61:39 anti:syn<sup>[20,21]</sup> as determined by <sup>1</sup>H NMR spectroscopy of the unpurified reaction mixture).


**Table 3.** Mannich reactions of various thioesters


entry	R	product	yield (%) <sup>[b]</sup>
1	4-FPh	<b>2a</b>	99
2	4-BrPh	<b>2b</b>	81
3	2-naphthyl	<b>2c</b>	74
4	benzyl	<b>2d</b>	84

[a] Reaction conditions: 1.0 mmol thioester, 1.2 mmol imine, 1.3 mmol TMSOTf, 1.2 mmol Et<sub>3</sub>N, 2.5 mL CH<sub>2</sub>Cl<sub>2</sub>, rt, 2 h. [b] Isolated yield after chromatography.




Amides are also excellent substrates.<sup>[22]</sup> Treatment of *N*-phenyl tertiary amides<sup>[23]</sup> with the benzaldehyde-derived *N*-phenylimine provided β-amino amides in very high yield (eq 4, >95% yield in both cases). These remarkably positive and consistent results suggest that *N*-aryl amides are powerful substrates in these reactions, and bear further study. It should be noted, however that when *N*-acetylmorpholine or *N*-methoxy-*N*-methylacetamide was employed as the enolate precursor, only very low yield was observed (6% and 36% yield, respectively).



Mukaiyama–Mannich reactions of thioester and acyclic amide substrates<sup>[24]</sup> have been relatively unexplored and therefore constituted the primary focus of this study. In contrast, methyl ester derivatives have been widely used in Mukaiyama–Mannich reactions, albeit after independent synthesis and isolation of their silyl ketene acetal derivatives.<sup>[21,25]</sup> Although reliable methods already exist for the two-step version of this reaction for ester substrates, we were still eager to test our conditions. Somewhat to our surprise, application of the conditions optimized for thioesters to ester substrates was quite disappointing, generally providing <20% conversion even with extended reaction times. Ultimately, however, suitable yields could be achieved by simply increasing the stoichiometry of the TMSOTf from 1.3 to 1.7 equiv. As illustrated in Table 4, a range of acetate esters acted as effective substrates under these

conditions, including the industrially friendly isopropyl acetate and the easily substituted phenyl acetate.

**Table 4.** Mannich reactions of various esters




entry	R	product	yield (%) <sup>[b]</sup>
1	Me	<b>5a</b>	73
2	Et	<b>5b</b>	80
3	<i>i</i> -Pr	<b>5c</b>	78
4	Ph	<b>5d</b>	69

[a] Reaction conditions: 1.0 mmol thioester, 1.2 mmol imine, 1.7 mmol TMSOTf, 1.2 mmol Et<sub>3</sub>N, 2.5 mL CH<sub>2</sub>Cl<sub>2</sub>, rt, overnight. [b] Isolated yield after chromatography.


Upon evaluation of our success with thioester, amide, and ester enolate precursors, a reexamination of ketones was appropriate. Historically, aryl ketones have been the highest-performing substrates in our one-pot enol silane formation-carbon-carbon bond-forming reactions,<sup>[26]</sup> and in the present study they again provided acceptable yields for a range of aryl aldimines (Table 5). Nonetheless, these reactions were complicated by the consistent generation of a competing product. It was ultimately confirmed that the byproducts resulted from Friedel–Crafts attack of the *N*-phenyl group on the carbonyl, a phenomenon preceded in the literature<sup>[27]</sup> but not observed with other substrates in our study. The identity of this byproduct was confirmed through treatment of adduct **6a** with TMSOTf and *i*-Pr<sub>2</sub>NEt, which resulted in about 50% conversion to cyclic byproduct **7** as determined by <sup>1</sup>H NMR spectroscopy (eq 5). Further optimization attempts failed to eliminate the generation of this class of byproducts under the reaction conditions. No evidence of analogous byproduct formation was observed for non-ketone substrates.

**Table 5.** Mannich reactions of acetophenone



entry	R	product	yield (%) <sup>[b]</sup>
1	Ph	<b>5a</b>	82
2	4-MeOPh	<b>5b</b>	78
3	4-NO <sub>2</sub> Ph	<b>5c</b>	68
4	4-FPh	<b>5d</b>	76
5	4-BrPh	<b>5e</b>	65

[a] Reaction conditions: 1.0 mmol thioester, 1.2 mmol imine, 1.7 mmol TMSOTf, 1.2 mmol Et<sub>3</sub>N, 2.5 mL CH<sub>2</sub>Cl<sub>2</sub>, rt, overnight. [b] Isolated yield after chromatography.



## Conclusions

We have successfully developed the one-pot silyl ketene acetal-formation-Mukaiyama–Mannich addition of thioesters to *N*-phenylimines. This methodology has been extended to include amide, ester, and ketone precursors, all of which provide products in deprotected form and in high yield. The TMSOTf appears to play two roles: first as a silylating agent that generates the requisite silyl ketene acetal intermediate, and second as a Lewis acid that activates the imine toward nucleophilic attack. Further study will be directed toward silyl triflate activation of other nitrogen-based electrophiles.

## Experimental Section

**General.** Reactions were carried out under an atmosphere of nitrogen with a septum cap in oven-dried glassware with magnetic stirring. Methylene chloride was purified by passage through a bed of activated alumina.<sup>[28]</sup> Trimethylsilyl trifluoromethanesulfonate (TMSOTf) was stored in a Schlenk flask under inert atmosphere. Hunig's base (*i*-Pr<sub>2</sub>NEt) was distilled from calcium hydride and stored in a Schlenk flask under inert atmosphere. All other chemicals were used as received or prepared according to literature precedent. Purification of reaction products was carried out by flash chromatography using silica gel (230–400 mesh). Analytical thin layer chromatography was performed on silica gel plates. Visualization was accomplished with UV light and phosphomolybdic acid stain, followed by heating. Infrared spectra were recorded on an FT-IR spectrometer. <sup>1</sup>H NMR spectra were recorded on a 500 MHz spectrometer or 300 MHz spectrometer, and are reported in ppm using solvent as an internal standard (CDCl<sub>3</sub> at 7.28 ppm). Data are

reported as (ap = apparent, s = singlet, d = doublet, t = triplet, q = quartet, sx = sextet, sp=septet, m = multiplet, b = broad; coupling constant(s) in Hz; integration). Proton-decoupled  $^{13}\text{C}$  NMR spectra were recorded on a 125 MHz spectrometer or 75 MHz spectrometer, and are reported in ppm using solvent as an internal standard ( $\text{CDCl}_3$  at 77.0 ppm). High-resolution mass spectra were obtained by electrospray ionization unless otherwise indicated. Melting points were determined using a capillary melting point apparatus.

#### General Procedure A. Mukaiyama–Mannich reactions of thioesters

To an oven-dried round-bottomed flask under  $\text{N}_2$  atmosphere was added methylene chloride (2.5 mL), thioester (1.00 mmol), and  $\text{Et}_3\text{N}$  (167  $\mu\text{L}$ , 121 mg, 1.20 mmol). Trimethylsilyl trifluoromethanesulfonate (235  $\mu\text{L}$ , 289 mg, 1.30 mmol) was added dropwise. After 5 min, imine (1.20 mmol) was added and the reaction was stirred for 2 h. The reaction mixture was passed through a column of silica (2 cm x 1 cm) with methylene chloride. The solvent was removed in vacuo. Column chromatography of the residue (0 – 15% EtOAc/hexanes) provided the product.

#### General Procedure B. Mukaiyama–Mannich reactions of amides

To an oven-dried round-bottomed flask under  $\text{N}_2$  atmosphere was added amide (1.00 mmol), methylene chloride (2.5 mL), and  $i\text{-Pr}_2\text{NEt}$  (209  $\mu\text{L}$ , 155 mg, 1.20 mmol). Trimethylsilyl trifluoromethanesulfonate (235  $\mu\text{L}$ , 289 mg, 1.30 mmol) was added dropwise. After 5 min,  $N$ -benzylideneaniline (217 mg, 1.20 mmol) was added and the reaction was stirred for 2 h. The reaction mixture was passed through a column of silica (2 cm x 1 cm) with methylene chloride. The solvent was removed in vacuo. Column chromatography of the residue (0 – 20% EtOAc/hexanes with 1% diethylamine) provided the product.

#### General Procedure C. Mukaiyama–Mannich reactions of esters

To an oven-dried round-bottomed flask under  $\text{N}_2$  atmosphere was added  $N$ -benzylideneaniline (217 mg, 1.20 mmol), methylene chloride (1.0 mL), ester (1.00 mmol), and  $i\text{-Pr}_2\text{NEt}$  (209  $\mu\text{L}$ , 155 mg, 1.20 mmol). Trimethylsilyl trifluoromethanesulfonate (308  $\mu\text{L}$ , 378 mg, 1.70 mmol) was added dropwise, and the reaction was stirred for overnight. The reaction mixture was passed through a column of silica (2 cm x 1 cm) with methylene chloride. The solvent was removed in vacuo. Column chromatography of the residue (0 – 20% EtOAc/hexanes) provided the product.

#### General Procedure D. Mukaiyama–Mannich reactions of acetophenone

To an oven-dried round-bottomed flask under  $\text{N}_2$  atmosphere was added methylene chloride (2.5 mL), acetophenone (115  $\mu\text{L}$ , 120 mg 1.00 mmol), and  $i\text{-Pr}_2\text{NEt}$  (209  $\mu\text{L}$ , 155 mg, 1.20 mmol). Trimethylsilyl trifluoromethanesulfonate (235  $\mu\text{L}$ , 289 mg, 1.30 mmol) was added dropwise. After 5 min, imine (1.20 mmol) was added and the reaction was stirred for 2 h. The reaction mixture was passed through a column of silica (2 cm x 1 cm) with methylene chloride. The solvent was removed in vacuo. Column chromatography of the residue (0 – 5% EtOAc/hexanes) provided the product.

## Acknowledgements

We thank the National Science Foundation RUI program (CHE-1057591) and the Camille and Henry Dreyfus Foundation for

funding. We are indebted to the National Science Foundation (CHE-0541848) and the University of California-Riverside for mass spectral data.

**Keywords:** Mukaiyama–Mannich • aniline synthesis • silyl ketene acetal • silyl triflate • one pot

- [1] For recent reviews, see: a) Merino, P., *Sci. Synth.*, P. W. N. M. Van Leeuwen, Ed., **2014**, 2, 311-331. b) T. Akiyama, *Comp. Org. Synth.*, 2<sup>nd</sup> Edition, P. Knochel, G. A. Molander, Eds., **2014**, 2, 629-681. c) G. Roman, *Eur. J. Med. Chem.* **2015**, 89, 743-816.
- [2] For recent examples, see: a) S. Matsukawa, K. Fukazawa, J. Kimura, *RSC Adv.* **2014**, 4, 27780-27786. b) E. Lee, T. Ryu, Y. Park, S. Park, P. H. Lee, *Adv. Synth. Catal.* **2013**, 355, 1585-1596. c) T. Kano, T. Yurino, D. Asakawa, K. Maruoka, *Angew. Chem. Int. Ed.* **2013**, 52, 5532-5534. d) B. Ranieri, A. Sartori, C. Curti, L. Battistini, G. Rasso, G. Pelosi, G. Casiraghi, F. Zanardi, *Org. Lett.* **2014**, 16, 932-935.
- [3] For recent reviews, see: a) N. Mase, C. F. Barbas, III, *Comprehensive Enantioselective Organocatalysis*, P. I. Dalko, Ed., **2013**, 3, 793-840. b) N. H. Nguyen, A. B. Hughes, B. E. Sleebs, *Curr. Org. Chem.* **2014**, 18, 260-289. c) B. Karimi, D. Enders, E. Jafari, *Synthesis* **2013**, 45, 2769-2812.
- [4] For a recent example, see: a) C. Curti, L. Battistini, B. Ranieri, G. Pelosi, G. Rasso, G. Casiraghi, F. Zanardi, *J. Org. Chem.* **2011**, 76, 2248-2252.
- [5] For recent reviews, see: a) Q. Zhang, X. Liu, X. Feng, *Curr. Org. Synth.* **2013**, 10, 764-785. b) C. Schneider, M. Sickert, In *Chiral Amine Synth.*, T. C. Nugent, Ed., Wiley: New York, **2010**; 157-177.
- [6] For recent examples, see: a) A. Sartori, L. Dell'Amico, L. Battistini, C. Curti, S. Rivara, D. Pala, P. S. Kerry, G. Pelosi, G. Casiraghi, G. Rasso, *Org. Biomolec. Chem.* **2014**, 12, 1561-1569. b) A. Sartori, L. Dell'Amico, C. Curti, L. Battistini, G. Pelosi, G. Rasso, G. Casiraghi, F. Zanardi, *Adv. Synth. Catal.* **2011**, 353, 3278-3284. c) B. M. Ruff, S. Zhong, M. Nieger, M. Sickert, C. Schneider, S. Brase *Eur. J. Org. Chem.* **2011**, 6558-6566. d) S. Dhanasekaran, A. Kayet, A. Suneja, V. Bisai, V. K. Singh, *Org. Lett.* **2015**, 17, 2780-2783.
- [7] For recent examples, see: a) C. R. Jones, G. D. Pantos, A. J. Morrison, M. D. Smith, *Angew. Chem. Int. Ed.* **2009**, 48, 7391-7394. b) Q. Wang, M. van Gemmeren, B. List, *Angew. Chem. Int. Ed.* **2014**, 53, 13592-13595. c) D. S. Giera, M. Sickert, C. Schneider, *Org. Lett.*, **2008**, 10, 4259-4262. d) F. Abels, C. Lindemann, C. Schneider, *Chem. Eur. J.* **2014**, 20, 1964-1979.
- [8] C. W. Downey, C. M. Dombrowski, E. N. Maxwell, C. L. Safran, O. A. Akomah, *Eur. J. Org. Chem.* **2013**, 5716-5720.
- [9] For Mukaiyama–Mannich additions to nitrones using ester-derived silyl ketene acetals, see: a) C. Qian, L. Wang, *Tetrahedron* **2000**, 56, 7193-7197. (b) S.-I. Murahashi, Y. Imada, T. Kawakami, K. Harada, T. Yonemushi, N. Tomita, *J. Am. Chem. Soc.* **2002**, 124, 2888-2889. b) Merino, P.; Jimenez, P.; Tejero, T. *J. Org. Chem.* **2006**, 71, 4685-4688. c) Diez-Martinez, A.; Tehero, T.; Merino, P. *Tetrahedron: Asymmetry* **2010**, 21, 2934-2943. For computational studies of the mechanism of this reaction type, see: d) A. Milet, Y. Gimbert, A. E. Greene, *J. Comput. Chem.* **2005**, 27, 157-162. e) L. R. Domingo, M. Arnó, P. Merino, T. Tejero, *Eur. J. Org. Chem.* **2006**, 3464-3472.
- [10] For representative examples, see: a) C. W. Downey, M. W. Johnson, D. H. Lawrence, A. S. Fleisher, K. J. Tracy, *J. Org. Chem.* **2010**, 75, 5351-5354. b) C. W. Downey, A. S. Fleisher, J. R. Rague, C. L. Safran, M. E. Venable, R. D. Pike, *Tetrahedron Lett.* **2011**, 52, 4756-4759. c) C. Wade Downey, S. E. Covington, D. C. Obenshain, E. Halliday, J. T. Rague, D. N. Confair *Tetrahedron Lett.* **2014**, 55, 5213-5215. d) C. W. Downey, E. N. Maxwell, D. N. Confair *Tetrahedron Lett.* **2014**, 55, 4959-4961.

- [11] J. E. Huheey, E. A. Keiter, R. L. Keiter, *Inorganic Chemistry: Principles of Structure and Reactivity*, 4th ed; HarperCollins: New York, 1993.
- [12] S. Katsuki, M. Ito *Lancet* **1966**, 1966-II, 530-2.
- [13] F. C. Reubi *Ann. New York Acad. of Sci.* **1966**, 139, 433-42.
- [14] E. H. Ostergaard, M. P. Magnussen, C. K. Nielsen, E. Eilertsen, H. H. Frey *Arzneimittel-Forschung* **1972**, 22, 66-72.
- [15] N. H. Huel, H. Nar, H. Priepke, U. Ries, J. M. Stassen, W. Wienen; Nar; Priepke; Ries; Stassen; Wienen *J. Med. Chem.* **2002**, 45, 1757-66.
- [16] The superiority of Et<sub>3</sub>N vs. *i*-Pr<sub>2</sub>NEt for the deprotonation of thioesters has been observed in a previous study with nitron acceptors. See reference 8.
- [17] For example, *N*-methylaniline can be silylated with TMSCl and Et<sub>3</sub>N in good yield: a) C. J. Smith, M. W. S. Tsang, A. B. Holmes, R. L. Danheiser, J. W. Tester, *Org. Biomolec. Chem.* **2005**, 3, 3767-3781. b) C. J. Smith, T. R. Early, A. B. Holmes, R. E. Shute, *Chem. Commun.* **2004**, 1976-1977.
- [18] For a recent example, see: E. B. Rowland, G. B. Rowland, E. Rivera-Otero, J. C. Antilla, *J. Am. Chem. Soc.* **2007**, 129, 12084-12085.
- [19] For a recent example, see reference 7c.
- [20] Identity of major isomer tentatively assigned based on H-H coupling constants (major = 6.7 Hz, minor = 5.3 Hz).
- [21] For a related and highly diastereoselective Ti-mediated aldol reaction between unactivated esters and *O*-methyloximes, see: T. Funatomi, S. Nakazawa, K. Matsumoto, R. Nagase, Y. Tanabe *Chem Commun.* **2008**, 771-773.
- [22] For similar system employing catalytic TIPSOTf, see: S. Kobayashi, H. Kiyohara, M. Yamaguchi *J. Am. Chem. Soc.* **2011**, 133, 708-711.
- [23] Secondary amides form relatively unreactive silyl imidates under the reaction conditions. See reference 10b.
- [24] For an example of an acyclic amide-derived silyl ketene acetal employed in the Mukaiyama-Mannich reaction, see reference 7c.
- [25] For examples, see references 2a, 6c, 7a, 7b, and 7d. Ethyl ester Mannich products are frequently prepared under Reformatsky conditions. For recent examples, see: a) S. Sengmany, E. Le Gall, M. Troupel, *Synlett* **2008**, 1031-1035. b) X. Chen, C. Zhang, H. Wu, X. Yu, W. Su, J. Cheng, *Synthesis* **2007**, 3233-3239.
- [26] a) C. W. Downey, M. W. Johnson, *Tetrahedron Lett.* **2007**, 48, 3559-3562. b) C. W. Downey, M. W. Johnson, K. J. Tracy, *J. Org. Chem.* **2008**, 73, 3299-3302. See also references 8 and 10c.
- [27] a) H. Laurent-Robert, B. Garrigues, J. Dubac, *Synlett* **2000**, 1160-1162. b) Y. Makioka, T. Shindo, Y. Taniguchi, K. Takaki, Y. Fujiwara, *Synthesis* **1995**, 27, 801-804. c) S. Kobayashi, H. Ishitani, S. Nagayama, *Synthesis* **1995**, 27, 1195-1202.
- [28] Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, 15, 1518-1520.

WILEY-VCH


---

## Entry for the Table of Contents (Please choose one layout)

Layout 1:

## FULL PAPER

Text for Table of Contents

**Key Topic\***

Author(s), Corresponding Author(s)\*


Page No. – Page No.

Title

\*one or two words that highlight the emphasis of the paper or the field of the study

Layout 2:

## FULL PAPER



Thioesters, amides, esters, and ketones undergo conversion to the respective silyl ketene acetals and addition to imines in one pot when treated with trimethylsilyl trifluoromethanesulfonate and trialkylamine.

\*one or two words that highlight the emphasis of the paper or the field of the study

**Mukaiyama–Mannich**

C. Wade Downey,\* Jared A. Ingersoll,  
Hadleigh M. Glist, Carolyn M.  
Dombrowski, Adam T. Barnett

Page No. – Page No.

**One-pot silyl ketene acetal-formation-  
Mukaiyama–Mannich additions to  
imines mediated by trimethylsilyl  
trifluoromethanesulfonate**