

9-9-2010

# The Identity of the Crackling, Luminescent Frog of Suriname (*Rana typhonia* Linnaeus, 1758) (Amphibia, Anura)

E. O. Lavilla

José A. Langone

José M. Padial

Rafael O. de Sá

University of Richmond, rdesa@richmond.edu

Follow this and additional works at: <http://scholarship.richmond.edu/biology-faculty-publications>Part of the [Biology Commons](#), [Population Biology Commons](#), [Terrestrial and Aquatic Ecology Commons](#), and the [Zoology Commons](#)

## Recommended Citation

Lavilla, E. O., J. A. Langone, J. M. Padial, and Rafael O. de Sá. "The Identity of the Crackling, Luminescent Frog of Suriname (*Rana typhonia* Linnaeus, 1758) (Amphibia, Anura)." *Zootaxa* 2671 (November 9, 2010): 17-30.

This Article is brought to you for free and open access by the Biology at UR Scholarship Repository. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact [scholarshiprepository@richmond.edu](mailto:scholarshiprepository@richmond.edu).



## The identity of the crackling, luminescent frog of Suriname (*Rana typhonia* Linnaeus, 1758) (Amphibia, Anura)

E.O. LAVILLA<sup>1</sup>, J.A. LANGONE<sup>2</sup>, J.M. PADIAL<sup>3,4</sup> & R.O. DE SÁ<sup>5</sup>

<sup>1</sup>Fundación Miguel Lillo, Miguel Lillo 251, 4000 Tucumán, Argentina, E-mail: eolavilla@gmail.com

<sup>2</sup>Departamento de Herpetología, Museo Nacional de Historia Natural, Casilla de Correo 399, 11.000, Montevideo, Uruguay.  
E-mail: pplangone@fcien.edu.uy

<sup>3</sup>Department of Evolution, Genomics and Systematics, Evolutionary Biology Centre (EBC), Uppsala University, Norbyvägen 18D, 75236 Uppsala, Sweden; E-mail: jose.m.padial@ebc.uu.se

<sup>4</sup>Current address: Division of Vertebrate Zoology (Herpetology), American Museum of Natural History, Central Park West at 79<sup>th</sup> street, New York, NY 10024.

<sup>5</sup>Department of Biology, University of Richmond, Richmond, VA, 23173, USA. E-mail: rdesa@richmond.edu

<sup>6</sup>Corresponding author

### Abstract

Review of the literature and recently available field notes from the collector of the type allows a reconsideration of the identity of the Linnaean name *Rana typhonia*. We provide evidence to demonstrate that the Linnaean species is neither a bufonid nor an Asiatic ranid, but a Neotropical hylid. Subsequently, we consider *Rana typhonia* as an older synonym of *Rana venulosa* Laurenti, 1768, redescribing its holotype under the new combination, *Trachycephalus typhonius* (Linnaeus, 1758).

**Key words:** Nomenclature, *Trachycephalus venulosus*, *Rana typhonia*, Holotype

### Introduction

Although more than 250 years of Linnaean taxonomy have elapsed since the publication of Linnaeus's (1758) *Systema Naturae*, longstanding mistakes related to some of the very first species described under the Linnaean era (i.a., Dubois and Ohler, 1996; Lavilla *et al.* 2010) still persist. Here, we analyze the identity of *Rana typhonia* Linnaeus, 1758, and come to a radically different view on the identity of that frog. The tenth edition of *Systema Naturae* (Linnaeus 1758) marks the beginning of Linnaean taxonomy in zoology. In this work, Linnaeus included seventeen species of anurans, fifteen of which come from his previous contributions or those by diverse other authors. The remaining two, the only ones originally described by Linnaeus in that work, were the Palearctic *Rana variegata* (today in the genus *Bombina*) and the Neotropical *Rana typhonia*. During the last 250 years the name *Rana typhonia* was associated with a group of Neotropical toads with dorsolaterally expanded supratympanic crests (the *Bufo typhonius* or *margaritifer* phenetic group, *sensu* Duellman and Schulte 1992; Frost 2010). *Rana typhonia* was considered a junior synonym of the Asiatic *Hoplobatrachus tigerinus* (Daudin, An XI [1801]) by Hoogmoed (1989), based on Andersson (1900), who first proposed the synonymy. Although the priority of *Rana typhonia* Linnaeus, 1758 over *Rana tigerina* Daudin, An XI is evident, no formal taxonomic action was done. However, the identity of *Rana typhonia* is still an open question for several other reasons.

First, the Asian provenance of *Rana typhonia* is challenged by the fact that Linnaeus (1758) explicitly expressed that its origin was "America". Furthermore, he also explicitly quoted the name of the collector, Rolander (an exception among the amphibians included in the tenth edition of the *Systema Naturae*), referring to one of his disciples who traveled to Surinam. Second, based on: (a) Boie (1827), who analyzed a fragment

of Daniel Rolander's diary (the *Diarium Surinamicum, quod sub itinere exotico conscripsit Daniel Rolander, tomus I and II*), which remained unpublished in the Danish Botanical Library for more than 250 years and (b) on the integral translation done by a team led by James Dobreff (with Clarck Dahlman, Joseph Morgan and Joseph Tipton) and edited by Hansen (2008), it is now possible to demonstrate without a doubt that the Linnaean *Rana typhonia* is neither a bufonid nor an Asian ranid, but a Neotropical hylid.

Thus, this contribution focuses on: (1) the clarification of the attribution of the Linnaean name *Rana typhonia*, (2) establish a type locality for this neotropical taxon, and (3) the re-description of the holotype of the Linnaean species, based on its identification among the specimens of the *Donatio Alströmer* (Thunberg 1787, Lönnberg 1896; Wallin 2001).

## Material and methods

The first part of this contribution is a bibliographic inquest and, as such, the analyzed materials were [almost] all the texts that, starting with Linnaeus (1758) and ending with Hansen (2008), dealt with the life and work of Daniel Rolander and his collections. Only those contributions considered relevant are included in the text.

The second part consists of a re-description of the holotype of *Rana typhonia* under the new combination *Trachycephalus typhoni* (Linnaeus 1758), as a senior synonym of *Trachycephalus venulosus* (Laurenti 1768), based on the individual number 134 at the Evolution Museum of Uppsala University (UUZM).

Measurements were taken with digital calliper to the nearest 0.1 mm; abbreviations (in mm) are: SVL (snout–vent length); HL (head length); HW (head width); IND (internarial distance); END (eye to nostril distance); ED (eye diameter); UEW (upper eyelid width); IOD (interorbital distance); TD (tympanum diameter); HAL (hand length); THL (thigh length); TL (tibia length); FL (foot length, including tarsus). Webbing formulae follow Savage and Heyer (1997).

Following ICZN Art. 8.3 (International Commission on Zoological Nomenclature, ICZN 1999) the name *Rana crepitans* Rolander in Boie (1827) (*nomen oblitum*), made available by Hansen (2008), is not considered here a scientific binomen for nomenclatorial purposes, and the name *Rana typhonica* used by Hansen (2008) is considered a misspelling of *Rana typhonia* Linnaeus 1758.

## Results

### 1. Comments on the collector of *Rana typhonia* and his collections

*Rana typhonia* is unique among the anurans considered by Linnaeus (1758) in having an explicitly named collector, quoted as “Rolander”, one of Linnaeus's disciples who traveled to Surinam between 1755 and 1756. The “official history” says that after the trip “...Rolander brought home a considerable collection of subjects of natural history, but was ungrateful enough not to present his kind patron with any of them...” (Pulteney 1805: 132); furthermore, Andersson (1900) quoted “...I do not know whether Rolander, who traveled in Surinam and West India (sic), brought home from these places any collections, among which Linnaeus discovered the type specimen of his *Rana typhonia*...”

These, and several other more or less offensive comments (including a recent one by Speake 2003, who stated that Rolander lost his mind in Guyana and returned a helpless cripple) are unjustified considering that Linnaeus gave Rolander's name to at least two species [*Cimex (Oblongus) Rolandri* and *Phalaena (Tortrix) Rolandriana*] and in the “*Collectanea*” specified in the *Systema Naturae*, Linnaeus (1758) included as one of his sources “*D. Rolandri in Surinamum and Eustatium. 1755.*” Furthermore, the tenth edition of the *Systema Naturae* assigned Rolander as the collector of about one hundred Neotropical taxa, including 85 species of insects (listed in Dobreff, 2009b), at least one bat (*Vespertilio spectrum*), seven birds (*Falco sufflator*, *F. cachinnans*, *Gracula foetida*, *G. barita*, *Ardea striata*, *A. aequinoctialis*, and *Loxia minuta*), one amphibian (*Rana typhonia*), one lizard (*Lacerta angulata*), and one snake (*Boa canina*) (names follow Linnaeus 1758).

## 2. *Rana typhonia* Linnaeus, 1758, the crackling and luminescent frog of Rolander

Searching for the identity of *Rana typhonia* and following ICZN Art. 72.4.1.1, we analyzed Daniel Rolander's *Diarium Surinamicum, quod sub itinere exotico conscripsit Daniel Rolander, tomus I and II. 1754-1756*. As noted above, until the integral English translation published by Hansen (2008), the only clue we had about Rolander's diary was an excerpt published by Boie (1827) (that of Duméril and Bibron 1841 is a French translation of Boie's paper). Hansen (2008) work gives us a completely new prospect on the identity of *Rana typhonia*. Also, thanks to the kindness of Prof. Dr. James Dobreff, leading translator of Rolander's Diary (Hansen 2008) and Editor of Daniel Rolander *Diarium Surinamicum* (in press), we had a deep and first hand analysis of the conflictive first paragraph of the description, published as "...*Rana crepitans, quae Rana typhonia\* dicitur...*" by Boie (1827) (the asterisk refers in Boie's work to "*Lin. system. nat. Seba thes I pag. 114 t. 71 f. 3. 4.*").

According to Dobreff (2009a) the *Diarium*, drafted during the trip, was exhaustively reviewed and updated in Copenhagen between 1763 and 1765, resulting in the inclusion, as notes, of several Linnaean names. Rolander originally diary says "... *Rana crepitans, quae Rana Surinamensis dicitur...*", with a footnote referring to "Seba, Thes.I. pag. 114, t. 71, f. 3. 4." Then, some time after the publication of the tenth edition of the *Systema Naturae* (Linnaeus 1758), Rolander crossed out "*Surinamensis*" and added just above the deletion "*typhonia*", forgetting or likely not bothering to remove the footnote referring to Seba's *Rana, Surinamensis*. Lastly, the Danish naturalist Martin Vahl got the original manuscript of *Diarium Surinamicum* and put an asterisk after "*typhonia*", referring to his own note in the margin "*Lin. Sys.*" (Dobreff *in litt.*, April 10<sup>th</sup>, 2010).

In Rolander's diary, the species was characterized as:

"...*Rana crepitans, quae Rana ~~Surinamensis~~ typhonia dicitur magnitudine Ranae aquaticae est, supra fusca, subtus flava; puncta elevata, convexa, inaequalia per corpus sparsa conspiciuntur: eadem haec puncta instructa sunt emissariis, e quibus excernere solent humorem illum lacteum, quo totum corpus obducit, ut quae modo fusca erat, intra inomentum albissima appareat. Palmae tetradactylae fissae; plantae pentadactylae, subpalmatae; digitorum apices rotundati, planiusculi. Indis americanis cibo sunt...*".

Thus, Rolander's description shows that *Rana crepitans*, which was named *Rana ~~Surinamensis~~ typhonia* has a size comparable to that of *Rana aquaticae* (probably a *Rana temporaria* or a *Rana esculenta sensu* Linnaeus 1758), is dorsally dark and yellowish ventrally, and has elevated, convex, and unequal warts scattered across the body. From these warts, a milky fluid is produced that covers the entire body, so the dark dorsal areas suddenly appear white. The hands have four fingers, separated, and the feet five toes, semipalmated, being the tip of the digits flattened and rounded. The American Indians eat them.

In addition, Boie (1827) included a seemingly textual transcription of Rolander's diary on the events on November, 19<sup>th</sup> [1755], in order to discuss a case of luminescence in the frog collected by Rolander. In that section, Rolander commented his vicissitudes trying to identify the source of the sounds that bewildered him at night, until he realized that the responsables were a group of frogs hidden in the roof of a barn, and the fact that on that night he was awakened by a noise similar to that of a crow, describing the situation as "... *somnum mihi ademit crepitatio cornicans, ingrattissima...*".

In summary, Rolander's corrections on his own manuscript suggest that Linnaeus eventually described and named his *Rana crepitans* as *Rana typhonia*. If the explicit note added by Rolander to his manuscript is still considered unconvincing, the reinterpretation of Linnaeus's (1758) diagnosis in light of this new information should help to support the identity proposed herein. *Rana typhonia* was characterized as:

*R[ana] auricularibus lobis ovatis. Habitat in America, clamitans nocte fono cornicis tetro lucente.*  
Rolander.

The following translation and analysis of Linnaeus's description further supports our interpretation:

1. *R[ana] auricularibus lobis ovatis*: The “ovate ear lobes” had two different interpretations. Dozens of authors had followed Schneider (1799), who re-defined this character state as “...*Margo capitis aucta membrana altra supra oculos et aures eminente*” (i.e., “...Margins of the head with a noticeable high membrane on the eyes and ears...”); this was the basis for the application of the name *Rana typhonia* to toads now placed in the *Rhinella typhonia* or *margaritifera* group. On the other hand, other authors (i.a., Andersson 1900; Hoogmoed 1989) considered them as “the large blown out vocal sacs” shown by the individual labeled by Quensel in the Stockholm Museum catalogue as *Rana typhonia*. Furthermore, this character state was later the basis for Andersson's (1900) synonymization of *Rana typhonia* with *Hoplobatrachus tigerinus*.
2. *Habitat in America*: Although the origin is imprecise by modern standards, it fits the Linnaean style [note that Linnaeus 1758, described anurans from *America* (5), *Europa(e)* (5), *Exteris regionibus* (2), *Indiis* (2), *Surinami* (2), and *Virginia* (1)].
3. *clamitans nocte fono cornicis tetro lucente*  
Kitchell and Dundee (1994) translated the later sentence as a single phrase, meaning “calling by night with the foul sound of the crow as it grows light”. Under the new evidence, we consider the statement to encompass two characters:
  - 3.1 *Clamitans nocte fono cornicis*: This characterizes the strident call of the species, similar to that of a crow, and has a noticeable similitude with the reference to the “...*crepitatio cornicans, ingratisissima*...” mentioned by Rolander for his *Rana crepitans*, and
  - 3.2 *tetro lucente*: Refers to the ability to produce a dim light or glow. Rolander noted that the barn where the frogs were calling seemed as if illuminated by a yellow foxfire (“...*horreum idem forte intravi, quod crepitantibus Ranis, quasi igne flavescente fatuo, illustrabatur*...”), and the glow was purportedly produced by the frogs “...as they alternately open and close their mouths to croak...” (Boie 1827; Hansen 2008).

Considering that Linnaeus was never in Surinam, and almost certainly never heard the mating call of *Rana typhonia*, nor did he observe the alleged bioluminescence, it is highly probable that the origin of these two key characters in Linnaeus's description stem from Rolander's notes.

The diagnosis previously analyzed was followed by a brief description, applicable to various species of treefrogs (Linnaeus 1758):

*Dorsum rugis quatuor longitudinalibus, punctis elevatis, maculisque nigris. Pedes mutici. Palmae tetradactylae fissae; Plantae pentadactylae palmatae; digitis angustis; secundo longissimo; absque unguibus orbiculatis.*

Kitchell and Dundee (1994) translated the above paragraph as:

“The back has four longitudinal wrinkles, raised dots, and black spots. The feet are stubby, front feet four-toed and split, rear feet five-toed and palmate. The toes are narrow, the second being the longest, but lacking rounded claws”.

The last character state (“but lacking rounded claws”) describes any frog except a treefrog; however, the key is in the latin word *absque*. It can be translated as a negative, as Kitchell and Dundee (1994) did, but *absque* also means “with the exception of” (Various Authors 1968). Thus, we translate the sentence “...*digitis angustis; secundo longissimo; absque unguibus orbiculatis*...” as “... toes narrow, except for the rounded tip, the second being the longest...”.

Consequently, and also based on Rolander (*in* Boie 1827 and Hansen 2008) and Linnaeus (1758), we have no doubts that *Rana typhonia* is the name given by Linnaeus to Rolander's *Rana crepitans*.

### 3. The identity of *Rana typhonia* Linnaeus, 1758

Based on Linnaeus (1758) (L) and Rolander (*in* Boie 1827) (R), it is possible to re-define *Rana typhonia* as a tree frog (R) with a size comparable to a European water frog (R) [about 10 cm SVL], in which males have paired vocal sacs (L). The body is covered by elevated, convex, and unequal warts that produce an abundant milky secretion (R). The hands have four separated fingers (R, L) and the feet five semipalmated (R) or palmated toes (L), with expanded tips (R, L), the second being the longest (L). The mating call is similar to the call of a crow (R, L), and that supposedly produced a gloomy light (R, L). The species was collected by Rolander (L) in America (L).

Aside from an incongruent character-state (i.e., the relative length of the second toe, explainable by considering it as the second digit from the foot's outer edge), the frog's habits, size, number of vocal sacs, skin texture, quality and amount of secretions, characteristics of the mating calls, and geographical origin, lead us to think that *Rana typhonia* is a species comparable to those contained either in the genera *Trachycephalus* Tschudi, 1838 or *Osteocephalus* Steindachner, 1862.

Regarding the alluded luminescence, Rolander noted that when the frogs croak, "... their wide-opened mouth glitters with a yellow colour. This is why, as they alternately open and close their mouths to croak, they seem to be emitting a sort of yellow fire. Thus, if you enter late in the evening a building filled with their croaking, you will notice a sort of yellowish flame, which vanishes during the pauses in their song..." (Hansen 2008). This peculiar trait has never been reported again in a Surinamese frog, and it is unlikely that this would represent any kind of amphibian bioluminescence. A plausible explanation, following Rolander's observations on the persistence of the light produced by an insect that he called *Cantahris pyrallis* (see entrance of June, 17th), is that the "yellowish will o' the wisp" observed glowing from the frog's mouth was due to the persistence of active remains of luciferin-luciferase complex after eating fireflies.

The key question is: Did Linnaeus have an individual of any of these genera in his hands? A series of circumstantial evidence suggest that in fact he did.

Despite the legend, Linnaeus included Rolander as one of the sources of the *Systema Naturae* X, described about one hundred species collected by his disciple in Surinam and St. Eustatius, and, based on the field observations referenced in the case of *Rana typhonia*, probably had access not only to a part of Rolander's material, but likely to his notes as well.

Another clue comes from Thunberg (1787), who listed the type of *Rana typhonia* among the material donated by Alströmer to Linnaeus starting in 1749 and extending over a period of several years. The specimen was reported by Lönnberg (1896) as "8) «*Rana typhonia*» Thunberg non Linnaeus is *Hyla venulosa* (Laurenti)", and by Wallin (2001), who identified the specimen as UUZM 134, and considered it a misidentification by Lönnberg (1896) due to an error in Thunberg's label. Lönnberg's and Wallin's confusion and efforts to clarify the specimens' identification are understandable because since the end of the 18th century (probably starting with Schneider 1799) up to the end of the 20<sup>th</sup> century, the name *Rana typhonia* has been associated with a toad, not with a treefrog. However, it is critical to note that Lönnberg (1896) considered that the Alströmer Donation was a Class B collection, defined as those "...which have been known by Linnaeus, and have been under his care so that the determination of these specimens, when they agree with the descriptions by the same name in «*Systema Naturae*», have a certain authority. In fact, in such cases when Linnaeus in «*Systema Naturae*» does not refer to any other specimens, figures or descriptions, it can almost be supposed that some of these animals have been the types. The identification of these specimens thus has a certain value in dubious cases, as it is always sure that Linnaeus has seen them, and known their characteristics..."

Lönnberg's (1896) statement strongly supports the idea that the individual identified by Thunberg (1787) as *Rana typhonia* was the one on which Linnaeus based his description, a point of view also supported by the presence of the individual hitherto considered the holotype of *Gymnoderus foetidus* (Linnaeus 1758) (= *Gracula foetida*), also collected by Rolander during the same field collection of *Rana typhonia* (Thunberg 1787; Lönnberg 1896).

In summary, we consider that the individual UUZM 134, today identifiable as *Trachycephalus venulosus* (Laurenti, 1768) as the holotype of *Rana typhonia* Linnaeus, 1758, following ICZN Arts. 72.4.1.1 and 73.1.2 (ICZN 1999) and based on the following circumstantial evidence:

1. Linnaeus (1758) cited “Rolander” as the collector of his *Rana typhonia*, and stated “America” as the locality of the specimen.
2. Rolander explicitly stated the synonymy of his *Rana crepitans* with Linnaeus (1758) *Rana typhonia*.
3. The morphological and ethological characteristics of Rolander’s *Rana crepitans* match with some of those of *Trachycephalus*.
4. The occurrence of Rolander specimens described as new species by Linnaeus (1758) scattered among diverse collections [i.e., De Greer’s, for almost all insects (Dobref 2009 a, b); *Adolphi-Friederici* for *Boa canina* (= *Corallus caninus*) (Thunberg 1787; Lönnberg 1896), and *Jonae Alströmer* for *Gracula foetida* (= *Gymnoderus foetidus*) (Thunberg 1787; Lönnberg 1896)] supports the idea that the *Rana typhonia* specimen in the Alströmer Donation (Thunberg 1787) corresponds to the one collected by Rolander.
5. Based on the comments under Type B collections in Lönnberg (1896), it is highly probable that the specimen cited by Thunberg (1787) as *Rana typhonia*, and later identified as *Hyla venulosa* by Lönnberg (1896) and Wallin (2001), was the one studied by Linnaeus.

This challenges the current conception of the identification of the holotype of *Rana typhonia*. Furthermore, the sources of the uncertainty on its existence and whereabouts are twofold:

- (a) The mistaken characterization of *Rana typhonia* as a toad by Schneider (1799), a fact that likely led Lönnberg (1896) and Wallin (2001) to overlook the specimen deposited at Uppsala Museum under the name *Rana typhonia* because it was clearly a tree-frog, and
- (b) The erroneous belief that the holotype was housed at the Stockholm Museum, a fact that led Quensel (followed by Andersson 1900) to identify the individual NRM 142 (a male of *Hoplobatrachus tigerinus*), under the name *Rana typhonia*, with the known subsequent story<sup>1</sup>.

The holotype of *Rana typhonia* (UUZM 134) is preserved in good condition at the Evolution Museum of Uppsala University. Our examination of this specimen reveals that *Rana typhonia* is in fact a member of the genus *Trachycephalus*, and more specifically it corresponds to what is currently known as *Trachycephalus venulosus*. Thus, *Trachycephalus venulosus* (Laurenti, 1768) is considered a junior synonym of *Trachycephalus typhoni* (Linnaeus, 1758), nov. comb.

#### 4. Characterization of the Holotype of *Trachycephalus typhoni* (Linnaeus, 1758) (Figure 1)

Adult male, UUZM 134 (Rolander and Linnaeus’s characters in brackets). Robust build; head wider than long, HL 92% of HW, HL 29% of SVL, HW 32% of SVL. Snout truncated in dorsal view, rounded in profile; *canthus rostralis* rounded; loreal region slightly convex. Nostrils closer to tip of snout than to eyes; internarial distance longer than eye to nostril distance and 60% of eye diameter. Eye to nostril distance shorter than eye diameter, than upper eyelid width and than interorbital distance, but longer than tympanum diameter. Tympanum ovate, annulus distinct; tympanum largely separated from eye, its diameter shorter than eye diameter, TD 49% of ED. Dorsal skin coarsely warty, warts mostly irregular in shape, low, and flat, with a few scattered ones enlarged and conical, most common on the posterior half of the dorsum (R: *puncta elevata, convexa, inaequalia per corpus sparsa conspiciuntur*; L: *punctis elevatis, maculisque nigris*); skin on head and eyelids slightly granular; flanks warty, with warts larger than those on dorsum; dorsal folds absent (L: *Dorsum rugis quatuor longitudinalibus*; not coincident); skin on belly and groin coarsely areolate; anal region

1. In the electronic herpetological database of the Swedish Museum of Natural History, under NRM 142 - *Rana typhonia* is the following remark: “Systematic position and type status to be evaluated. If Rolander’s specimen (the type) was donated to the academy and later catalogued by Hornstedt, that specimen probably was discarded by Sparrman or mixed by Quensel. Evaluation includes finding and examining all specimens included in the catalogue 1802 /EIA, 2004”

not modified; dorsal and ventral surfaces of arms smooth with scarce scattered low and soft granules; dorsal surface of thighs and tibiae smooth. Vocal sacs double, lateral, well developed (L: *auricularibus lobis ovatis*), nuptial pads on thumb, single, rugose, dark brown. Dentigerous process of vomers prominent, almost contacting medially, laying between and at the same level of the choanae, bearing a row of vomerine teeth. Tongue large, rounded, almost completely attached. Hand with robust fingers, webbed, tips expanded, rounded; webbing formulae I - II<sub>1/2</sub> - 2III1 - 1IV; finger lengths 3 > 4 > 2 > 1; subarticular tubercles conical, prominent; few small, rounded and low supernumerary tubercles present; outer metacarpal tubercle divided into two elongated, unequal in length, and low tubercles; inner metacarpal tubercle elongated, equal in length to the larger inner tubercle, but more prominent (R and L: *Palmae tetradactylae fissae*). Legs slender, tibia length equals thigh length; SVL 94% of the combined tibia and thigh lengths. Foot length shorter than tibia and thigh lengths, 38% of SVL. Toes robust, webbed (R: *plantae pentadactylae, subpalmatae*, L: *Plantae pentadactylae palmatae*), webbing formula I1 - 2II1 - 2III1 - 1IV1 - 1V; toe lengths 4 > 3 > 5 > 2 > 1; toe tips expanded, disk complete, rounded (R: *digitorum apices rotundati, planiusculi*; L: *digitis angustis ... absque unguibus orbiculatis*); subarticular tubercles large, rounded, supernumerary tubercles not evident; outer metatarsal tubercle round, low, inconspicuous; inner metatarsal tubercle ovate, slightly elevated; sole of tarsus with small inconspicuous granules; tarsal fold absent.

Measurements (in mm): SVL 72.6 (R: *magnitudine Ranae aquaticae est*); HL 21.1; HW 23.0; IND 5.6; END 5.6; ED 9.6; UEW 5.9; IOD 7.2; TD 4.7; HAL 20.0; THL 33.9; TL 34.0; FL 27.35.

Color in preservative: overall dorsal background reddish cream with irregular reddish brown blotches; the most conspicuous mark is an irregular, inverted Y-shaped mark that runs from occipital to sacral regions (R: *supra fusca*); small irregular blotches in interocular region; flanks cream with a reddish brown, irregular, and diffuse stripe, and a large diffuse blotch behind the arm; arms and legs with transverse reddish brown stripes. Tympanic membrane cream, but areas immediately posterior to eyes, around the tympanic membrane, and around the vocal sac, reddish brown. Vocal sacs brown. *Canthus rostralis* and loreal regions cream. Ventral region reddish cream, darker on gular region (R: *subtus flava*). Plantar surfaces faded reddish brown. Fingers and toes cream, barred with reddish brown stripes.

## 5. Comments on the type locality

Museum records associated with specimen UUZM 134 provide no clues regarding the provenance of the holotype and the only published data mentions “America” in the original description (Linnaeus 1758) rendering inapplicable sections 1 and 2 of ICZN Recommendation 76A.1.

A first restriction can be made to Surinam, the country visited by the collector and, reading his Journal, it is possible to establish that Rolander caught at least one individual of *Trachycephalus typhonius* on the night of November 19<sup>th</sup>, 1755, in a barn. Although no locality is mentioned, it was probably located at Brouwershaven, one of Carl Gustav Dahlberg wife’s Plantations, on the left bank of the Wajamo River, a tributary of the Pirica River, at about nine parasangs from Paramaribo (see entrances of July 21<sup>st</sup> and 22<sup>nd</sup> in Hansen 2008). Alternatively, based on the Journal (Rec. 76A.1.3), it is possible to establish that Rolander heard frog calls and was aware of their eggs on July, 9<sup>th</sup>, 1755 in Paramaribo, in one of the funniest passages of his Diary (“...everyday I come across their eggs immersed in a slimy substance and floating on the water's surface. By the cemetery that is next to the church of the Reformed there has been constructed an opera house, where twice a week operas are performed late in the evening. However, the frogs almost always break into the sonorous arrangement and harmony of the musical instruments with their extremely loud, and just as unwelcome, noise so as to disrupt the charm of the music and make for the audience a very unpleasant croaking...”).

Faced with our second goal of determining the type locality for the species, we chose the most conservative of the two options, without prejudice to other clarification (Rec. 76A.1.4). The cemetery alluded by Rolander and called “De Oranjetuin” (the Orange Garden), was sited on today’s Kerkplein (the Church’s square), while the Reformed Church was in the upper floor of a large building placed at the site of the current Royal Bank of Trinidad and Tobago (RBTT Bank), at number 1, Kerkplein Street, Paramaribo, Suriname ([http://www.surinameembassyjakarta.org/index2.php?main=visit\\_historicwalk.php](http://www.surinameembassyjakarta.org/index2.php?main=visit_historicwalk.php)).



**FIGURE 1.** Holotype of *Trachycephalus typhoni* (Linnaeus, 1758), adult male, UUZM 134, SVL 72.6 mm.

#### **6. Comments on some forgotten names related to *Trachycephalus typhoni* (Linnaeus, 1758)**

##### ***Rana vesicaria* Fermin, 1765.**

The species is included among the *Anura incertae sedis* (Frost 2010), but the presence of paired vocal sacs (“... *Cette Grenouille a à chaque côté de la mâchoire inférieure, une Vessie, qui dans les grandes chaleurs sont remplies d’air...*”) and its origin (“Hollande Equinoxiale”, i.e., Surinam), leads us to consider it as a synonym of *Trachycephalus typhoni*.

Although Fermin’s book was included in the Official Index of Rejected and Invalid Works by Opinion 660 (International Commission on Zoological Nomenclature 1963), only the names of seven species of turtles

were formally included in the list of rejected taxa. The rejection of the book suppress all the contained species, and our comment on *Rana vesicaria* is only intended to tie a loose end in the nomenclature of Neotropical amphibians.

### ***Hyla micans* Oken, 1836.**

At the end of the French translation of Boie's (1827) paper on Rolander's luminescent frog ("*Rana crepitans*, quae *Rana typhonia* dicitur"), Duméril and Bibron (1841) noted that no subsequent authors (including themselves in the systematic section of the eighth volume of the *Érpetologie Générale*) cited the species. This statement is partially true; although never mentioned as *Rana crepitans*, Rolander's species, under a different name, was occasionally cited in the literature.

In fact, in 1836 Oken translated into German Boie's (1827) transcription of Rolander's *Diarium Surinamicum*, naming the species as *H[yla] micans*, noting that the author considered this frog as a toad ("*...Bufo typhoni*us, *margaritif*er..."), despite all the evidence contained in the translated paper. *Hyla micans* was overlooked by the great majority of herpetologists and the only references we found hitherto, aside from the original, are those of Clark (1953), Hartwig (1863; 1871), Holder and Holder (1885), Kappler (1887), Korn and Smith (1959), Schoedler (1853), Schumacher (1844), and Shufeldt (1896). All these contributions contain only a couple of lines with the anecdote of the luminescence, in some cases noting (erroneously) that it was due to corporal secretions, but without any new information.

### ***Discodactylus* Wagler, 1833.**

The name *Discodactylus* was coined for the individuals depicted in figures 1 to 5, plate 71, of Seba (1734), the same plate and figures mentioned by Rolander (see above). The figures represent two different species, named by Wagler *Discodactylus tibicen* (figs. 1–2, males; 3 female) (a replacement name for *Hyla tibiatrix* Laurenti, 1768 and *Calamita tibicen* Merrem, 1820), and *Discodactylus marmoratus* (figures 4 and 5) (a replacement name for *Hyla marmorata* Daudin, An XI). The name *Discodactylus* was coined by Wagler to replace the names *Hyla* and *Calamita* due to their apparent inadequacy.

Although *Discodactylus* Wagler 1833 antedates *Trachycephalus* Tschudi, 1838, Wagler's name fits the considerations of Art. 23.9 of the Code (IZCN 1999) and, consequently, it is possible to apply the reversal of precedence. *Discodactylus* and its contained species (*Discodactylus dumerili*, *D. marmoratus*, *D. phacophorus*, *D. pulcher*, *D. ruber*, *D. tibicen*, and *D. tuberculatus*) were included in Sherborn (1932) *Index Animalium*, but it was never used again in reference to an amphibian, being ignored to the point that Fitzinger (1843: 18; 95) used it to name a genus of geckos, fulfilling the requirements of Arts. 23.9.1.1 and 23.9.6. In turn, *Trachycephalus* has been used extensively, e.g., just in the last five years, by at least Abrunhosa *et al.* (2006); Acosta-Galvis *et al.* (2006); Alvarez *et al.* (2009); Barrio-Amoros and Brewer-Carias (2008); Bastazini *et al.* (2007); Bernarde (2007); Borges and Freitas (2007); Borges-Martins *et al.* (2007); Brandão and Araújo (2008); Cabagna Zenklusen *et al.* (2009); Cintra *et al.* (2009); Cisneros-Heredia (2006; 2007); De Freitas and Oliveira Lima (2009); Faivovich *et al.* (2005); Frost *et al.* (2006); Jungfer (2010); Kok and Kalamandeen (2008); Kwet and Sole (2008); Pimenta and Canedo (2007); Ramirez Valverde *et al.* (2009); Rigolo *et al.* (2008); Rodrigues *et al.* (2005); Rodrigues (2008); Wilms *et al.* (2005); Wogel *et al.* (2006); and Ziegler (2008), fulfilling the requirements of Art. 23.9.1.2.

In conclusion, we consider *Trachycephalus* Tschudi, 1838 as *nomen protectum* and *Discodactylus* Wagler, 1833 as *nomen oblitum*.

### **7. Additional remarks.**

As currently understood, *Trachycephalus typhoni*us is a species distributed from southern Mexico to the North of Argentina (Frost 2010; IUCN 2010). This extensive geographic range, together with its morphological variability (i.a., Duellman 2001; Eterovick and Sazima 2004; McDiarmid 1968), suggest that this taxon may represent a species complex. This situation sets up the scenario for a complex nomenclatorial history, whose resolution will be possible only after of a comprehensive study, including morphology, calls, ecology and genetic data of this taxon throughout its entire distribution.

The following is an abbreviated “state of the art” synonymy, based primarily on Frost (2010). It includes only the original morphonims (in the sense of Dubois 2000) and the various generic combinations of different specific epithets, mentioning in each case only the first author of the *nomina* and only the first page where it appears.

***Trachycephalus typhonius* (Linnaeus, 1758)**

- Rana typhonia* Linnaeus, 1758: 211.  
*Trachycephalus typhonius* — this contribution
- Rana venulosa* Laurenti, 1768: 31.  
*Hyla venulosa* — Daudin 1800: 17.  
*Hypsiboas venulosa* — Wagler 1830: 201.  
*Phrynohyas (Acrodytes) venulosa* — Fitzinger 1843: 30.  
*Scytotis venulosus* — Cope 1866: 85.
- Hyla tibiatrix* Laurenti, 1768: 34.  
*Auletris tibiatrix* — Wagler 1830: 201.  
*Discodactylus tibiatrix* — Wagler 1833: 888.  
*Dendrohyas tibiatrix* — Tschudi 1838: 34
- [?] *Hyla viridi-fusca* Laurenti, 1768: 34.
- [x] *Rana reticularis* Lacépède, 1788: 296, 458. Rejected.
- Hyla intermixta* Daudin, *In* Sonnini and Latreille, An. X: 182.
- Hyla variegata* Daudin, An XI: 20.
- Rana meriana* Shaw, 1802: 133.
- Calamita tibicen* Merrem, 1820: 172.  
*Discodactylus tibicen* — Wagler 1833: 888.
- [x] *Hyla zonata* Spix, 1824: 41. Rejected.  
*Hypsiboas zonata* — Wagler 1830: 200.  
*Phrynohyas zonata* — Fitzinger 1843: 30.
- Hyla bufonia* Spix, 1824: 42.  
*Hypsiboas bufonia* — Wagler 1830: 200.
- [x] *H[yla]. micans* Oken, 1836: 471. *Nomen oblitum*.  
*R[ana]. micans* Schoedler 1853: 587.
- Hyla vermiculata* Duméril and Bibron, 1841: 563.
- Hyla lichenosa* Günther, 1858: 327.
- Scytotis hebes* Cope, 1862: 354.  
*Phrynohyas hebes* — Duellman 1956: 16.  
*Trachycephalus venulosus hebes* — Kwet and Sole 2008: 55
- Hyla spilomma* Cope, 1877: 86.  
*Acrodytes spilomma* — Taylor 1944: 64.  
*Phrynohyas spilomma* — Duellman 1956: 28.
- Hyla paenulata* Brocchi, 1879: 21.
- Hyla nigropunctata* Boulenger, 1882: 366.
- Hyla palpebrogranulata* Andersson, 1906: 14.
- Hyla wettsteini* Ahl, 1933: 25.
- Hyla zernyi* Ahl, 1933: 27.
- Acrodytes inflata* Taylor, 1944: 63.  
*Phrynohyas inflata* — Duellman 1956: 22.
- Acrodytes modesta* Taylor and Smith, 1945: 594.  
*Hyla modesta* — Mertens 1952: 30.

*Phrynohyas modesta* — Duellman 1956: 25.  
*Hyla macrotis* Andersson, 1945: 70.  
*Phrynohyas ingens* Duellman, 1956: 22.  
*Phrynohyas latifasciata* Duellman, 1956: 24.  
*Phrynohyas corasterias* Shannon and Humphrey, 1957: 15.  
*Hyla adenoderma* Lutz, 1968: 3.  
*Phrynohyas adenoderma* — Vanzolini 1986: 31.  
*Argenteohyla altamazonica* Henle, 1981: 134.

## Acknowledgements

We would like to thank the following colleagues for their assistance with some of the older references used in this work: Dione Seripieri, library of the Museu de Zoologia, Universidade de São Paulo (Brazil); Kraig Adler; Susan E. Olmsted, Librarian at NSF; Santiago Castroviejo-Fisher; Félix Muñoz Garmendia, from the Digital Library of the Real Jardín Botánico, Madrid; Francisco Pando, GBIF Spain, Real Jardín Botánico, Madrid. Sonia Kretschmar helped with the translation of text in gothic German. JMP by the EU Marie Curie Mobility and Training Programme (FP7, proposal 220714).

We are thankful to James Dobreff, Center for Language and Literature, Lund University, leading translator of Rolander's diary, and David Morgan, from the Department of Modern Languages and Literatures at Furman University (Greenville, SC) for helping in elucidating parts of the notes from Rolander's diary. Hans Mejlon (UUZM), Sven Kullander, Bodil Kajrup, and Erik Åhlander (NHRM) provided access and assistance with Linnaean collections.

## References

- Abrunhosa, P.A., Wogel, H. & Pombal Jr., J.P. (2006) Anuran temporal occupancy in a temporary pond from the Atlantic Rain Forest, South-Eastern Brazil. *The Herpetological Journal*, 16 (2), 115–122.
- Acosta-Galvis, A.R., Huertas-Salgado, C. & Rada, M. (2006) Aproximación al conocimiento de los anfibios en una localidad del Magdalena medio (Departamento de Caldas, Colombia). *Revista de la Academia Colombiana de Ciencias*, 30 (115), 291–303.
- Ahl, E. (1933) Über einige neu Frösche aus Brasilien. *Zoologischer Anzeiger*, 104, 25–30.
- Alvarez, B.B., García, J.A.R., Céspedes, J.A., Hernando, A.B., Zaracho, V.H., Calamante, C.C. & Aguirre, R.H. (2009) Herpetofauna, provinces of Chaco and Formosa, Chaco Oriental region, north-eastern Argentina. *Check List*, 5 (1), 74–82.
- Andersson, L.G. (1900) Catalogue of Linnæan type-specimens of Linnaeus's reptilia in The Royal Museum in Stockholm. *Bihang till Kongl. Svenska Vetenskaps-Akademiens Handlingar*, 26 (4) (1), 1–29.
- Andersson, L.G. (1906) On batrachians from Bolivia, Argentina, and Peru collected by Erland Nordenskiöld 1901-1902 and Nils Holmgren 1904–1905. *Arkiv för Zoologi*, 3, 1–24.
- Andersson, L.G. (1945) Batrachians from East Ecuador, collected 1937, 1938 by Wm. Clarke-Macintyre and Rolf Blomberg. *Arkiv för Zoologi*, 37, 1–88.
- Barrio-Amoros, C.L. & Brewer-Carias, C. (2008) Herpetological results of the 2002 expedition to Sarisariñama, a tepui in Venezuelan Guayana, with the description of five new species. *Zootaxa*, 1942, 1–68.
- Bastazini, C.V., Munduruca, J.F.V., Rocha, P.L.B. & Napoli, M.F. (2007) Which environmental variables better explain changes in anuran community composition? A case study in the restinga of Mata de São João, Bahia, Brazil. *Herpetologica*, 63 (4), 459–471.
- Bernarde, P.S. (2007) Ambientes e temporada de vocalização da anurofauna no Município de Espigão do Oeste, Rondônia, Sudoeste da Amazônia - Brasil (Amphibia: Anura). *Biota Neotropica*, 7 (2), 1–6.
- Boie, F. (1827) Naturhistorische Beiträge vermischten Inhaltes. *Isis von Oken* 20, VIII–IX: columns 726–730.
- Borges, F.J.A. & de Freitas, J.R. (2007) Distribuição espacial e temporal de uma comunidade de anuros do município de Morrinhos, Goiás, Brasil (Amphibia: Anura). *Neotropical Biology and Conservation*, 2 (1), 21–27.
- Borges-Martins, M., Colombo, P., Zank, C., Becker, F.G. & Melo, M.T.Q. (2007) Anfíbios. In: Becker FG, Ramos RA, Moura LA (Org.). *Biodiversidade. Regiões da Lagoa do Casamento e dos Butiazais de Tapes, Planície Costeira do Rio Grande do Sul*. Brasília, Ministério do Meio Ambiente, 276–291.

- Boulenger, G.A. (1882) *Catalogue of the Batrachia Salientia s. Ecaudata in the Collection of the British Museum*. Second Edition. London: Taylor and Francis/British Museum of Natural History.
- Brandão, R.A. & Araújo, A.F.B. (2008) Changes in anuran species richness and abundance resulting from hydroelectric dam flooding in Central Brazil. *Biotropica*, 40 (2), 263–266.
- Brocchi, P. (1879) Sur divers batraciens anoures de l'Amérique Centrale. *Bulletin de la Société Philomathique de Paris Ser.* 7 (3), 19–24.
- Cabagna Zenklusen, M.C., Lajmanovich, R.C., Peltzer, P.M., Attademo, A.M., Fiorenza Biancucci, G.S. & Basso, A. (2009) Primeros registros de endoparásitos en cinco especies de anfibios anuros del litoral Argentino. *Cuadernos de Herpetología*, 23 (1), 33–40.
- Cintra, C.E.D., Silva, H.L.R., da Silva, N.J., Garcia, P.C.A. & Zaher, H. (2009) A new species of *Trachycephalus* (Amphibia, Anura, Hylidae) from the State of Goiás, Brazil. *Zootaxa*, 1975, 58–68.
- Cisneros-Heredia, D.F. (2006) Amphibians, Machalilla National Park, province of Manabi, western Ecuador. *Check List*, 2 (1), 45–54.
- Cisneros-Heredia, D.F. (2007) Notes on the natural history of the casque-headed treefrog *Trachycephalus jordani* (Stejneger and Test, 1891). *Herpetozoa*, 20 (1–2), 92–94.
- Clark, A.M. (1953) The ecology, evolution, and distribution of the Vertebrates. *Annual Report of the Board of Regents of the Smithsonian Institution. Showing the operations, expenditures, and condition of the Institution for the year ended June 30 1952*. Publication 4111, 283–303.
- Cope, E.D. (1862) Catalogues of the reptiles obtained during the explorations of the Parana, Paraguay, Vermejo and Uruguay Rivers, by Capt. Thos. J. Page, U.S.N.; and of those procured by Lieut. N. Michler, U.S. Top. Eng., Commander of the expedition conducting the survey of the Atrato River. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 14, 346–359.
- Cope, E.D. (1866) On the structure and distribution of the genera of the arciferous Anura. *Journal of the Academy of Natural Sciences of Philadelphia*, Ser. 2 (6), 67–112.
- Cope, E.D. (1877) Tenth contribution to the herpetology of tropical America. *Proceedings of the American Philosophical Society*, 17, 85–98.
- Daudin, F.M. (1800) *Histoire Naturelle des Quadrupèdes Ovipaires*. Livraison 2. Paris: Marchant et Cie.
- Daudin, F.M. An. XI [1801] *Histoire Naturelle des Rainettes, des Grenouilles et des Crapauds*. Paris: Levrault.
- De Freitas, M.A. & de Oliveira Lima, T. (2009) *Trachycephalus nigromaculatus* (black-spotted casque-headed treefrog). *Herpetological Review*, 40 (3), 363.
- Dobref, J. (2009a) *Daniel Rolander. The invisible naturalist*. In A. Polaszeck (ed.). *Systema Naturae 250*. The Linnean Ark., Boca Raton, London, New York: CRC Press, 11–28.
- Dobref, J. (2009b) *Appendix 1: Concordance of Linnaeus's names for Rolander's insects*. In A. Polaszeck (ed.). *Systema Naturae 250*. The Linnean Ark., Boca Raton, London, New York: CRC Press, 253–264.
- Dubois, A. (2000) Synonymies and related lists in zoology: general proposals, with examples in herpetology. *Dumerilia*, 4, 33–98.
- Dubois, A. & Ohler, A. 1996 [1997] Early scientific names of Amphibia Anura II. An exemplary case: *Rana arborea* Linnaeus, 1758. *Bulletin du Muséum national d'Histoire naturelle*, Paris, 4e série, 18, Section A, n° 3–4, 321–340.
- Duellman, W.E. (1956) The frogs of the hylid genus *Phrynohyas* Fitzinger, 1843. *Miscellaneous Publications. Museum of Zoology, University of Michigan*, 96, 1–47.
- Duellman, W.E. (2001) *Hylid frogs of Middle America, expanded edition*. Ithaca: Society for the Study of Amphibians and Reptiles.
- Duellman, W.E. & Schulte, R. (1992) Description of a new species of *Bufo* from Northern Peru with comments on phenetic groups of South American toads (Anura: Bufonidae). *Copeia*, 1992 (1), 162–172.
- Duméril, A.M.C. & Bibron, G. (1841) *Erpétologie Générale ou Histoire naturelle complète des Reptiles. Tome Huitième, comprenant l'histoire générale des batraciens, et la description des cinquante-deux genres et des cent soixante-trois espèces des deux premiers sous-ordres: Les péromèles qui n'ont pas de membres, et les anoures qui sont privés de la queue*. Paris: Roret.
- Eterovick, P.C. & Sazima, I. (2004) *Anfibios da Serra do Cipó/Amphibians from the Serra do Cipó Minas Gerais – Brasil*. Belo Horizonte: PUC Minas.
- Faivovich, J., Haddad, C.F.B., Garcia, P.C.d.A., Frost, D.R., Campbell, J.A. & Wheeler, W.C. (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: a phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History*, 294, 1–240.
- Fermin, P. (1765) *Histoire naturelle de la Hollande Equinoxiale: Ou description des animaux, plantes, fruits, et autres curiosités naturelles, qui se trouvent dans la Colonie de Surinam; avec leurs noms différents, tant François, que Latins, Hollandais, Indiens Nègre-Anglois*. Amsterdam: M. Magerus.
- Fitzinger, L.J.F.J. (1843) *Systema Reptilium. Fasciculus Primus*. Wien: Braumüller et Seidel.
- Frost, D.R. (2010) *Amphibian Species of the World: an Online Reference*. Version 5.4 (June, 2010). Electronic Database accessible at <http://research.amnh.org/vz/herpetology/amphibia/>. American Museum of Natural History, New York, USA.

- Frost, D.R., Grant, T., Faivovich, J., Bain, R.H., Haas, A., Haddad, C.F.B., de Sá, R.O., Channing, A., Wilkinson, M., Donnellan, S.C., Raxworthy, C.J., Campbell, J.A., Blotto, B.L., Moler, P.E., Drewes, R.C., Nussbaum, R.A., Lynch, J.D., Green, D.M. & Wheeler, W.C. (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History*, 297, 1–370.
- Günther, A.C.L.G. (1858) Neue Batrachier in der Sammlung des britischen Museums. *Archiv für Naturgeschichte*, 24, 319–328.
- Hansen, L. (Editor-in-chief) (2008) *Daniel Rolander's Journal*. Translated by J. Dobreff, C. Dahlman, D. Morgan, and J. Tipton. In *The Linnaeus Apostles - Global science and adventure. Europe, North- South America*. London, Whitby: IK Foundation and Company. Vol. 3 Book 3, 1215–1576.
- Hartwig, G. (1863) *The tropical world: a popular scientific account of the natural history of the animal and vegetable kingdoms in the equatorial regions*. London: Longman, Green, Longman, Roberts, and Green.
- Hartwig, G. (1871) *The polar and tropical worlds: A description of man and nature in the polar and equatorial regions of the globe by G. Hartwig; edited, with additional chapters, by A. H. Guernsey*. San Francisco: A.L. Bancroft.
- Henle, K. (1981) *Argentohyla altamazonica*, ein neuer Hylid mit paarigen lateralen Schallblasen aus Peru (Amphibia: Salientia: Hylidae). *Amphibia-Reptilia*, 2, 133–137.
- Holder, Ch. & Holder, F.J.B. (1885) *Appletons' Science Text-books. Elements of Zoölogy*. New York, Cincinnati, Chicago: American Book Company.
- Hoogmoed, M.S. (1989) South American bufonids (Amphibia: Anura: Bufonidae), an enigma for taxonomists. *Treballs d'Ictiologica i Herpetologia*, 2, 167–180.
- International Commission on Zoological Nomenclature. (1963) Opinion 660. Suppression under the plenary powers of seven specific names of turtles (Reptilia, Testudines). *Bulletin of Zoological Nomenclature*, 20, 187–190.
- International Commission on Zoological Nomenclature. (1999) *International Code of Zoological Nomenclature*. International Trust for Zoological Nomenclature, London. <http://www.iczn.org/iczn/index.jsp>
- Jungfer, K.-H. (2010) The taxonomic status of some spiny-backed treefrogs, genus *Osteocephalus* (Amphibia: Anura: Hylidae). *Zootaxa*, 2407, 28–50.
- Kappler, A. (1887) *Surinam: sein Land, seine Natur, Bevölkerung und seine Kulturverhältnisse mit Bezug auf Kolonisation*. Stuttgart: J.G. Cotta.
- Kitchell, K. & Dundee, H.A. (1994) A trilogy on the Herpetology of Linnaeus's Systema Naturae X. *Smithsonian Herpetological Information Services*, 100, 1–61.
- Kok, P.J.R. & Kalamandeen, M. (2008) Introduction to the taxonomy of the amphibians of Kaieteur National Park, Guyana. *Abc Taxa*, 5, i–ix, 1–278.
- Korn, N. & Smith, H.R. (1959) *Human evolution: readings in physical anthropology*. New York: Holt.
- Kwet, A. & Sole, M. (2008) A new species of *Trachycephalus* (Anura: Hylidae) from the Atlantic Rain Forest in southern Brazil. *Zootaxa*, 1947, 53–67.
- Lacépède, B.G.É. (1788) *Histoire Naturelle des Quadrupèdes Ovipares et des Serpens, des Poisson et des Cetaces*. 16mo version. Volume 2. Paris: Hôtel de Thou.
- Laurenti, J.N. (1768) *Specimen Medicum, Exhibens Synopsin Reptilium Emendatum cum Experimentis Circa Venena et Antidota Reptilium Austriacorum*. Wien, Joan. Thom. nob. de Trattnern.
- Lavilla, E.O., Langone, J.A., Caramaschi, U., Heyer, W.R. & de Sá, R.O. (2010) The identification of *Rana ocellata* Linnaeus, 1758. Nomenclatural impact on the species currently known as *Leptodactylus ocellatus* (Leptodactylidae) and *Osteopilus brunneus* (Gosse, 1851) (Hylidae). *Zootaxa*, 2346, 1–16.
- Linnaeus, C. (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata*. Holmiæ: Salvius.
- Lönnberg, E. (1896) Linnean type-specimens of birds, reptiles, batrachians and fishes in the Zoological Museum of the R. University in Upsala. *Bihang till Kongliga Svenska vetenskaps-akademiens Handlingar*, 22 Afd. 4 (1), 1–45.
- Lutz, B. (1968) New Brazilian forms of *Hyla*. *Pearce-Sellards Series. Texas Memorial Museum*, 10, 3–18.
- McDiarmid, R.W. (1968) Populational variation in the frog genus *Phrynohyas* Fitzinger in Middle America. *Contributions in Science. Los Angeles County Museum*, 134, 1–25.
- Merrem, B. (1820) *Versuch eines Systems der Amphibien*. Marburg: Johann Christian Krieger.
- Mertens, R. (1952) Die Amphibien und Reptilien von El Salvador. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft, Frankfurt am Main*, 487, 1–120.
- Oken, L. (1836) *Allgemeine Naturgeschichte für alle Stände von Professor Oken*. Sechster Band, oder Thierreich, Dritter Band. Stuttgart: Hoffmann'sche Verlags.
- Pimenta, B.V.S. & Canedo, C. (2007) Description of the tadpole of *Itapotihyla langsdorffii* (Anura: Hylidae). *Zootaxa*, 1387, 39–46.
- Pulteney, R. (1805) *A general view of the writings of Linnaeus. The second edition; with corrections, considerable additions, and memoirs of the author; by William George Maton, M. D. F. R.S. F. S.A. to which is annexed the diary of Linnaeus, written by himself, and now translated into English, from the Swedish' manuscript in the possession of the editor*. London: J. Nawman and R. Taylor and Co.
- Ramirez Valverde, T., Sarmiento Rojas, A., Meza Parral, Y. & Martinez Campos, A. (2009) *Trachycephalus venulosus*

- (veined treefrog). *Herpetological Review*, 40 (4), 447–448.
- Rigolo, J.R., Almeida, J.A. & Ananias, F. (2008) Histochemistry of skin glands of *Trachycephalus* aff. *venulosus* Laurenti, 1768 (Anura, Hylidae). *Micron*, 39 (1), 56–60.
- Rodrigues, D.J., Uetanabaro, M. & Lopes, F.S. (2005) Reproductive patterns of *Trachycephalus venulosus* (Laurenti, 1768) and *Scinax fuscovarius* (Lutz, 1925) from the Cerrado, Central Brazil. *Journal of Natural History*, 39 (35), 3217–3226.
- Rodrigues, M.G. (2008) *Chironius exoletus* (common whipsnake): prey and possible diet convergence. *Herpetological Bulletin*, 105, 41–42.
- Savage, J.M. & Heyer, W.R. (1997) Digital webbing formulae for anurans: a refinement. *Herpetological Review*, 28 (3), 131.
- Schneider, J.G. (1799) *Historiae amphibiorum naturalis et literariae. Fasciculus primus continens ranas, calamitas, bufones, salamandras et hydros in genera et species descriptos notisque suis distinctos*. Ienae: Frommann.
- Schoedler, F. (1853) *The book of nature: an elementary introduction to the sciences of physics, astronomy, chemistry, mineralogy, geology, botany, zoology, and physiology. First American edition, with a glossary and other additions and improvements from the second English edition, translated from the sixth German edition by Henry Medlock*. Philadelphia: Blanchard and Lea.
- Schumacher, G.H. (1844) Würdigung Treviranus's als Biologie. In: *Biographische Skizzen verstorbener Bremischer Aerzte und Naturforscher: eine Festgabe für die zwei und zwanzigste Versammlung Deutscher Naturforscher und Aerzte zu Bremen*. Bremen: J.G. Heyse, 483–556.
- Seba, A. (1734) *Locupletissimi rerum naturalium thesauri accurata descriptio, et iconibus artificiosissimis expressio, per universam physices historiam. Opus, cui, in hoc rerum genere, nullum par exstitit. Ex toto terrarum orbe collegit, digessit, et depingendum curavit*. Tomus I. Amstelaedami: Wetstenium, Smith Janssonio–Waesbergios.
- Shannon, F.A. & Humphrey, F.L. (1957) A new species of *Phrynohyas* from Nayarit. *Herpetologica*, 13, 15–18.
- Shaw, G. (1802) *General Zoology, or systematic natural history*. Vol III. Part I: Amphibia. London: G. Kearsley.
- Sherborn, C.D. (1932) *Index Animalium sive index nominum quae ab A.D. MDCCLVIII generibus e specibus animalia imposita sunt. Sectio secunda | A kalendis Ianuariis MDCCCI | usque ad finem Decembris, MDCCCL. Part XXXI. Index to trivialia under genera, pp. 417–654 (Diabasis (Col.)-Lyurus). 1801–1850*. London: British Museum.
- Shufeldt, R.W. (1896) The frogs and their uses. *Appleton's Popular Science Monthly*. June, 1896, 179–185.
- Sonnini de Manoncourt, C.S., Latreille, P.A. & An, X. *Histoire Naturelle des Reptiles, avec Figures dissinées d'après Nature*. Volume 2. Paris: Deterville.
- Speake, J. (ed.) (2003) *Literature of travel and exploration: an encyclopedia*. Volumen 2, G to P. New York and London: Taylor & Francis.
- Spix, J.B.v. (1824) *Animalia nova sive Species novae Testudinum et Ranarum quas in itinere per Brasiliam annis MDCCCXVII–MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae Regis*. Monachii: Franc. Seraph. Hubschmanni.
- Steindachner, F. (1862) Über zwei noch unbeschriebene Batrachier aus des Sammlung des K. K. zoologischn Museum zu Wien. *Archivio per la Zoologia l'Anatomia e la Fisiologia, Genova*, 1 (2), 77–82.
- Taylor, E.H. (1944) The hylid genus *Acrodytes*, with comments on Mexican forms. *University of Kansas Science Bulletin*, 30, 63–69.
- Taylor, E.H. & Smith, H.M. (1945) Summary of the collections of amphibians made in México under the Walter Rathbone Bacon traveling scholarship. *Proceedings of the United States National Museum*, 95, 521–613.
- Thunberg, C.P. (1787) D. D. *Museum naturalium Academiae Upsaliensis cujus partem secundam, Consensu Exp. Fac. Med. Upsal. praeside Carol. Pet. Thunberg (...) publico examini proponit, Laur. Magn. Holmer, stipendiarius regius, Dalia-Wermelandus. In audit. Gust. Maj. D. XIV. April. anni MDCCCLXXVII. H. A. M. S. Upsaliae*: Edman, 17–32.
- Tschudi, J.J.v. (1838) *Classification der Batrachier mit Berücksichtigung der fossilen Thiere dieser Abtheilung der Reptilien*. Neuchatel: Petitpierre.
- Vanzolini, P.E. (1986) *Levantamento herpetológico da área do estado de Rondônia sob a influência da rodovia BR 364. Relatório de pesquisa*. Brasília: CNPq, Assessoria Editorial.
- Various, Authors (1968) *Oxford Latin Dictionary*. Oxford: Clarendon Press (Oxford University Press).
- Wagler, J. (1830) *Natürliches System der Amphibien, mit vorangehender Classification der Säugthiere und Vogel. Ein Beitrag zur vergleichenden Zoologie*. München, Stuttgart und Tübingen: J.G. Cotta
- Wagler, J. (1833) Deutung in der Seba's Thesaurus rerum Naturalium T. 1 et 2. enthaltenen Abbildungen von Lurchen, mit kritische Bemerkungen. *Isis von Oken*, 1833, 17(9), columns 886–905.
- Wallin, L. (2001) *Catalogue of type specimens. 4. Linnaean specimens*. Version 6. Uppsala University, Museum of Evolution, Zoology section. [www.evolutionsmuseet.uu.se/samling/UUZM04\\_Linnaeus.pdf](http://www.evolutionsmuseet.uu.se/samling/UUZM04_Linnaeus.pdf)
- Wilms, T., Kuegelgen, B. & Schreckenbach, T. (2005) Zur Haltung, Pflege und Vermehrung des Baumhoehlen - Kroetenlaubfrosches, *Trachycephalus resinifictrix* (Goeldi, 1907) im Reptilium Landau. *Draco*, 6 (3) 23, 66–72.
- Wogel, H., Weber, L.N. & Abrunhosa, P.A. (2006) The tadpole of the casque-headed frog, *Aparasphenodon brunoi* Miranda-Ribeiro (Anura: Hylidae). *South American Journal of Herpetology*, 1 (1), 54–60.
- Ziegler, T. (2008) "In situ" - und "ex situ" - Amphibienprojekte des Koelner Zoos: Forschung und Nachzucht als unser Beitrag zur Arterhaltung. *Zeitschrift des Koelner Zoo*, 51 (2), 67–88.