



Bookshelf

2016

Introduction to Model Spaces and their Operators

William T. Ross

University of Richmond, wross@richmond.edu

Stephan Ramon Garcia

Javad Mashreghi

Follow this and additional works at: <http://scholarship.richmond.edu/bookshelf>



Part of the [Computer Sciences Commons](#), and the [Mathematics Commons](#)

Recommended Citation

Garcia, Stephan Ramon, Javad Mashreghi, and William T. Ross. *Introduction to Model Spaces and their Operators*. Cambridge: Cambridge University Press. 2016.

NOTE: This PDF preview of *Introduction to Model Spaces and their Operators* includes only the preface and/or introduction. To purchase the full text, please click [here](#).

This Book is brought to you for free and open access by UR Scholarship Repository. It has been accepted for inclusion in Bookshelf by an authorized administrator of UR Scholarship Repository. For more information, please contact scholarshiprepository@richmond.edu.

QA
328
.C325
2011

Introduction to Model Spaces and their Operators

STEPHAN RAMON GARCIA

Pomona College, California

JAVAD MASHREGHI

Université Laval, Québec

WILLIAM T. ROSS

University of Richmond, Virginia

LIBRARY
UNIVERSITY OF RICHMOND
VIRGINIA 23173



CAMBRIDGE
UNIVERSITY PRESS

Preliminaries

1.1 Measure and integral

1.1.1 Borel sets and measures

Most of the “measuring” in this book will take place on the unit circle $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$. Since we assume that the reader has a background in graduate analysis, we quickly review the standard definitions without much fanfare.

We let $m := d\theta/2\pi$ denote *Lebesgue measure* on \mathbb{T} , normalized so that $m(\mathbb{T}) = 1$. A subset of \mathbb{T} is called a *Borel set* if it is contained in the *Borel σ -algebra*, the smallest σ -algebra of subsets of \mathbb{T} that contains all of the open arcs of \mathbb{T} . A *Borel measure* on \mathbb{T} is a countably additive function that assigns a complex number to each Borel subset of \mathbb{T} . Unless otherwise stated, our measures will always be finite. A Borel measure is *positive* if it assigns a non-negative number to each Borel set. We let $M(\mathbb{T})$ denote the set of all complex Borel measures on \mathbb{T} and we let $M_+(\mathbb{T})$ denote the set of all positive Borel measures on \mathbb{T} . A function $f : \mathbb{T} \rightarrow \widehat{\mathbb{C}}$ (where $\widehat{\mathbb{C}}$ denotes the Riemann sphere $\mathbb{C} \cup \{\infty\}$) satisfying the condition that $f^{-1}(U)$ is a Borel set for any open set $U \subset \widehat{\mathbb{C}}$ is called a *Borel function*.

We often need to distinguish between the “support” and a “carrier” of a measure. For $\mu \in M_+(\mathbb{T})$, consider the union \mathcal{U} of all the open subsets $U \subset \mathbb{T}$ for which $\mu(U) = 0$. The complement $\mathbb{T} \setminus \mathcal{U}$ is called the *support* of μ . On the other hand, a Borel set $E \subset \mathbb{T}$ for which

$$\mu(E \cap A) = \mu(A) \tag{1.1}$$

for all Borel subsets $A \subset \mathbb{T}$ is called a *carrier* of μ . The support of μ is certainly a carrier, but a carrier need not be the support. Indeed, a carrier of a measure might not even be closed. For example, if $f \geq 0$ is continuous and $d\mu = f dm$, then a carrier of μ is $\mathbb{T} \setminus f^{-1}(\{0\})$ (which is open) while the support of μ is the closure of this set. The support of a measure is unique while a carrier is not.