

3-2020

## On quasilinear parabolic equations and continuous maximal regularity

Jeremy LeCrone

University of Richmond, [jlecrone@richmond.edu](mailto:jlecrone@richmond.edu)

Gieri Simonett

Follow this and additional works at: <https://scholarship.richmond.edu/mathcs-faculty-publications>

 Part of the [Mathematics Commons](#)

This is a pre-publication author manuscript of the final, published article.

---

### Recommended Citation

LeCrone, Jeremy, and Gieri Simonett. "On Quasilinear Parabolic Equations and Continuous Maximal Regularity." *Evolution Equations and Control Theory* 9, no. 1 (March 2020): 61–86. <https://doi.org/10.3934/eect.2020017>.

This Post-print Article is brought to you for free and open access by the Math and Computer Science at UR Scholarship Repository. It has been accepted for inclusion in Math and Computer Science Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact [scholarshiprepository@richmond.edu](mailto:scholarshiprepository@richmond.edu).

# ON QUASILINEAR PARABOLIC EQUATIONS AND CONTINUOUS MAXIMAL REGULARITY

JEREMY LECRONE AND GIERI SIMONETT

ABSTRACT. We consider a class of abstract quasilinear parabolic problems with lower-order terms exhibiting a prescribed singular structure. We prove well-posedness and Lipschitz continuity of associated semiflows. Moreover, we investigate global existence of solutions and we extend the generalized principle of linearized stability to settings with initial values in critical spaces. These general results are applied to the surface diffusion flow in various settings.

## 1. INTRODUCTION

In this paper, we consider abstract quasilinear parabolic evolution equations given by

$$\begin{cases} \dot{u} + A(u)u = F_1(u) + F_2(u), & \text{for } t > 0, \\ u(0) = x, \end{cases} \quad (1.1)$$

for which we extend previous well-posedness and global existence results in the setting of continuous maximal regularity. As a particular feature, we admit nonlinearities  $F_2$  with a prescribed singular structure.

In more detail, we assume that  $(E_1, E_0)$  is a pair of Banach spaces so that  $E_1$  is densely embedded in  $E_0$ , and we seek solutions of (1.1) in time weighted spaces

$$\mathbb{E}_{1,\mu}(J) := BC_{1-\mu}^1(J, E_0) \cap BC_{1-\mu}(J, E_1),$$

where  $\mu \in (0, 1)$  and  $J = [0, T]$ , see Section 2 for a precise definition. Let  $V_\mu \subset E_\mu$  be an open subset of the continuous interpolation space  $E_\mu := (E_0, E_1)_{\mu, \infty}^0$ , and set  $\mathbb{E}_{0,\mu}(J) := BC_{1-\mu}(J, E_0)$ . Then we assume that the functions  $(A, F_1, F_2)$  satisfy the following conditions.

**(H1)** Local Lipschitz continuity of  $(A, F_1)$ :

$$(A, F_1) \in C^{1-}(V_\mu, \mathcal{M}_\mu(E_1, E_0) \times E_0). \quad (1.2)$$

Here  $\mathcal{M}_\mu(E_1, E_0)$  denotes the set of bounded linear operators  $B \in \mathcal{L}(E_1, E_0)$  for which  $(\mathbb{E}_{1,\mu}(J), \mathbb{E}_{0,\mu}(J))$  is a pair of maximal regularity for some  $T > 0$  (and hence all  $T > 0$ ). Therefore,  $B \in \mathcal{M}_\mu(E_1, E_0)$  if and only if for every  $(f, x) \in \mathbb{E}_{0,\mu}(J) \times E_\mu$  there exists a unique function  $u \in \mathbb{E}_{1,\mu}(J)$  such that

$$\dot{u}(t) + Bu(t) = f(t) \quad \text{for } t \in J, \quad \text{and } u(0) = x.$$

---

2010 *Mathematics Subject Classification.* Primary 35K90, 35K59, 35B30, 35B35; Secondary 53C44, 35K93.

*Key words and phrases.* quasilinear parabolic equations, continuous maximal regularity, generalized principle of linearized stability, critical spaces, well-posedness, surface diffusion flow, stability of cylinders, stability of spheres.

This work was supported by a grant from the Simons Foundation (#426729, Gieri Simonett).

**(H2)** Structural regularity of  $F_2$ :

There exists a number  $\beta \in (\mu, 1)$  such that  $F_2 : V_\mu \cap E_\beta \rightarrow E_0$ . Moreover, there are numbers  $\beta_j \in [\mu, \beta]$ ,  $\rho_j \geq 0$  and  $m \in \mathbb{N}$  with

$$\frac{\rho_j(\beta - \mu) + (\beta_j - \mu)}{1 - \mu} \leq 1, \quad \text{for all } j = 1, \dots, m, \quad (1.3)$$

so that for each  $x_0 \in V_\mu$  and  $R > 0$  there is a constant  $C_R = C_R(x_0) > 0$  for which the estimate

$$|F_2(x_1) - F_2(x_2)|_{E_0} \leq C_R \sum_{j=1}^m (1 + |x_1|_{E_\beta}^{\rho_j} + |x_2|_{E_\beta}^{\rho_j}) |x_1 - x_2|_{E_{\beta_j}} \quad (1.4)$$

holds for all  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, R) \cap (V_\mu \cap E_\beta)$ .

Following the convention introduced in Prüss, Wilke [20], we call the index  $j$  *subcritical* if (1.3) is a strict inequality, and *critical* in case equality holds in (1.3). As  $\beta_j \leq \beta < 1$ , any  $j$  with  $\rho_j = 0$  is subcritical. Furthermore, (1.3) is equivalent to  $\rho_j \beta + \beta_j - 1 \leq \rho_j \mu$ . Hence, the minimum value of  $\mu$  is given by

$$\mu_{\text{crit}} := \beta - \min_{\rho_j \neq 0} \left( \frac{1 - \beta_j}{\rho_j} \right).$$

The number  $\mu_{\text{crit}}$  is called the *critical weight* and  $E_{\mu_{\text{crit}}}$  a *critical space*.

In the last decades, there has been an increasing interest in finding critical spaces for nonlinear parabolic partial differential equations. As a matter of fact, there is no generally accepted definition in the mathematical literature concerning the notion of critical spaces. One possible definition may be based on the idea of a ‘largest space of initial data such that the given PDE is well-posed.’ Critical spaces are often introduced as ‘scaling invariant spaces,’ provided the underlying PDE enjoys a scaling invariance. It has been shown in [17] that the concept of *critical weight* and *critical space* introduced there (and also used in this paper) captures and unifies the idea of largest space and scaling invariant space. In more detail, it has been shown in [17] that  $E_{\mu_{\text{crit}}}$  is, in a generic sense, the largest space of initial data for which the given equation is well-posed, and that  $E_{\mu_{\text{crit}}}$  is scaling invariant, provided the given equation admits a scaling.

Our approach for establishing well-posedness of (1.1) relies on the concept of *continuous maximal regularity* in time-weighted spaces and extends previous results by Angenent [2], Clement and Simonett [5], Lunardi [13], and Asai [3]. The results parallel those in [9, 17, 20], where well-posedness of (1.1) is studied by means of maximal  $L_p$ -regularity in time-weighted function spaces.

Leveraging the singular structure of  $F_2$ , along with inequalities from interpolation theory and continuous maximal regularity, we prove local well-posedness of (1.1) via a fixed point argument. Allowing for rough initial values in  $E_\mu$  with  $\mu \geq \mu_{\text{crit}}$ , we prove Lipschitz continuity of the associated semiflow on  $V_\mu$  and derive conditions for global well-posedness and asymptotic behavior of solutions near normally stable equilibria. A key feature of our results is that dynamic properties of solutions are controlled in the topology of  $E_\mu$ , rather than requiring further control in stronger topologies of  $E_\beta$  or  $E_1$  as solutions regularize.

In particular, we prove that a priori bounds in the topology of  $E_\mu$  yield global existence. Moreover, we extend the generalized principle of linearized stability (c.f. [18, 19]), proving that solutions with initial data that is  $E_\mu$ -close to a normally stable equilibrium will converge exponentially fast to a nearby equilibrium.

As a particular application of our abstract results, we consider the surface diffusion flow, a geometric evolution equation acting on orientable hypersurfaces. Given a fixed reference manifold  $\Sigma \subset \mathbb{R}^n$ , we consider the evolution of surfaces  $\Gamma(t)$  defined in normal direction over  $\Sigma$  via time-dependent height functions  $h : \Sigma \times \mathbb{R}^+ \rightarrow \mathbb{R}$ . The governing equation for surface diffusion is then expressed as a fourth-order, parabolic evolution law acting on  $h = h(t)$  and we look for solutions in the setting of little-Hölder continuous functions; i.e.

$$E_0 := bc^\alpha(\Sigma) \quad \text{and} \quad E_1 := bc^{4+\alpha}(\Sigma), \quad \text{for some } \alpha \in (0, 1).$$

Considering the setting of  $\Sigma = \mathcal{C}_r \subset \mathbb{R}^3$ , an infinite cylinder of radius  $r > 0$ , we work with height functions  $h(t) : \mathcal{C}_r \rightarrow (-r, \infty)$  which produce so-called *axially-definable* surfaces  $\Gamma(h)$ , as in [12]. We show that the resulting surface diffusion flow can be cast as a quasilinear parabolic evolution equation in the form (1.1) with

$$E_\mu := bc^{1+\alpha}(\Sigma) \quad \text{and} \quad E_\beta := bc^{3+\alpha}(\Sigma),$$

from whence we have  $\mu = 1/4$  and  $\beta = 3/4$  in this setting. Explicitly expressing the singular nonlinearity  $F_2$ , we employ interpolation theory estimates to confirm the necessary singular structure **(H2)** is satisfied on  $V_\mu \cap E_\beta$ , where  $V_\mu$  is an appropriately chosen open subset of  $E_\mu$ . The appearance of several critical indices supports the idea that  $bc^{1+\alpha}(\Sigma)$  is in fact a critical space for surface diffusion flow.

Applying our general results to initial data  $h_0 \in bc^{1+\alpha}(\mathcal{C}_r)$  we extend well-posedness from [12, Proposition 3.2] to surfaces with only one Hölder continuous derivative. Further, we extend [11, Proposition 2.2, 2.3] by restricting to functions  $h_0 \in bc_{symm}^{1+\alpha}(\mathcal{C}_r)$  exhibiting azimuthal symmetry around the cylinder  $\mathcal{C}_r$ . Further, enforcing periodicity of  $h_0$  along the central axis of  $\mathcal{C}_r$ , we show stability and instability of cylinders under periodic perturbations with Hölder control on only first order derivatives. In particular, when  $r > 1$ , we show that  $2\pi$ -periodic  $\Gamma(h_0)$  surfaces that are  $bc^{1+\alpha}$ -close to  $\mathcal{C}_r$  give rise to global solutions to surface diffusion flow converging to a nearby cylinder exponentially fast. On the other hand, when  $r < 1$ , we show that there exist  $2\pi$ -periodic perturbations which are arbitrarily close to  $\mathcal{C}_r$  in  $bc^{1+\alpha}$  for which solutions escape a neighborhood of the cylinder. We also direct the reader to [4], for additional information concerning the surface diffusion flow for axisymmetric surfaces.

Taking  $\Sigma$  to be an arbitrary compact, connected, immersed manifold, we demonstrate well-posedness of surface diffusion for initial data in  $bc^{1+\alpha}(\Sigma)$  that are sufficiently close to the manifold  $\Sigma$ , an extension of [6, Theorem 1.1]. Further, in case  $\Sigma \subset \mathbb{R}^n$  is a Euclidean sphere, we apply our generalized stability result to yield stability of the family of spheres under perturbations which require control on only first-order derivatives. This result extends [6, Theorem 1.2], where initial values in  $bc^{2+\alpha}(\Sigma)$  are considered.

Working also in the setting of surfaces parameterized over a sphere, Escher and Mucha [7] show that small perturbations in the topology of Besov spaces  $B_{p,2}^{5/2-4/p}(\Sigma)$  exist globally and converge exponentially fast to a sphere. Although the topologies of these Besov spaces and our little-Hölder spaces are not easily comparable, we note that our stability results hold for any spacial dimension  $n$ , while the regularity of perturbations in [7] changes with  $n$ . In particular, Escher and Mucha enforce the bound  $p > \frac{2n+6}{3}$ , which they note only guarantees existence of lower regularity perturbations in  $C^{1+\alpha}(\Sigma) \setminus C^2(\Sigma)$  when  $n < 9$ . (Notice that the authors in [7] consider surfaces in  $\mathbb{R}^{n+1}$ .)

Regarding a different approach to stability of spheres, we refer to [14, 23, 24] where the lifespan of solutions, and convergence to equilibria, is controlled via  $L_2$ -estimates of the second fundamental form. We observe that our assumptions on initial data allow for initial surfaces on which the second fundamental form may not be defined, so that our results are not contained in [14, 23, 24].

As a final remark on surface diffusion flow, we mention that several authors have considered the flow of surfaces with rough initial data when  $\Gamma \subset \mathbb{R}^{n+1}$  is given as the graph of a function over a domain  $\Omega \subseteq \mathbb{R}^n$ , see [3, 8, 9] for instance. In [8], Koch and Lamm prove global existence of solutions to surface diffusion flow with initial surfaces that are merely Lipschitz continuous, and they prove analytic dependence on initial data. However, Koch and Lamm work in the setting of entire graphs (i.e.  $\Omega = \mathbb{R}^n$ ) and require a smallness condition on Lipschitz norm which seems to make it difficult to translate their result to more general settings. Working also in the setting of entire graphs, the conclusions of Asai in [3] are closest to our current results, as the author works in spaces of little-Hölder continuous functions. We refer to Remark 2.4 for a detailed account of the results in [3]. In [9], the authors approach surface diffusion flow from the setting of  $L_p$  maximal regularity on a bounded domain  $\Omega$ , producing well-posedness for initial data in Besov spaces  $B_{qp}^{4\mu-4/p}(\Omega)$ , for an appropriate choice of  $\mu$ ,  $p$ , and  $q$ .

We briefly outline the current paper. In Section 2, we state and prove our main result, Theorem 2.2. We conclude Section 2 with an extension of well-posedness, giving continuous dependence on initial data in stronger topologies  $E_{\bar{\mu}}$ .

In Section 3, we prove equivalence between measuring stability of equilibria in the space  $E_{\mu}$  and measuring stability in a smaller space  $E_{\bar{\mu}}$ ,  $\bar{\mu} \in [\beta, 1)$ . We then prove the generalized principle of linearized stability for perturbations in  $E_{\mu}$ .

In Section 4 we apply our results to various settings for surface diffusion flow. Beginning with *axially-definable* surfaces parameterized over an infinite cylinder, we conclude well-posedness of surface diffusion flow for general perturbations in  $bc^{1+\alpha}$ . Then we enforce periodicity in the general setting to establish stability / instability of cylinders under perturbations in  $bc_{per}^{1+\alpha}$  (with radius  $r$  above / below the threshold  $r = 1$ ), before producing similar results in the setting of axisymmetric surfaces. We end Section 4 with the setting of surfaces defined over an arbitrary compact reference manifold  $\Sigma$ , and establish well-posedness and stability of spheres with initial data in  $bc^{1+\alpha}$ .

## 2. WELL-POSEDNESS OF (1.1)

In this section, we formulate and prove our main result concerning solvability of (1.1). Moreover, we formulate and prove conditions for global existence of solutions. We start with the definition and elementary properties of *time-weighted continuous spaces* (see [5, Section 2] for more details).

Let  $E$  be an arbitrary Banach space and define the spaces of *time-weighted* continuous functions

$$BC_{1-\mu}(J, E) := \left\{ v \in C(\dot{J}, E) : [t \mapsto t^{1-\mu}v(t)] \in BC(\dot{J}, E), \lim_{t \rightarrow 0^+} |t^{1-\mu}v(t)|_E = 0 \right\}$$

and

$$BC_{1-\mu}^1(J, E) := \{v \in C^1(\dot{J}, E) : v, \dot{v} \in BC_{1-\mu}(J, E)\},$$

where  $J := [0, T]$ ,  $\dot{J} := (0, T]$ , and  $\mu \in (0, 1)$ . We also set

$$BC_0(J, E) := C(J, E), \quad \text{and} \quad BC_0^1(J, E) := C^1(J, E).$$

Given Banach spaces  $E_0$  and  $E_1$  so that  $E_1$  is densely embedded in  $E_0$ , we define

$$\begin{aligned} \mathbb{E}_{1,\mu}(J) &:= BC_{1-\mu}^1(J, E_0) \cap BC_{1-\mu}(J, E_1), \quad \text{and} \\ \mathbb{E}_{0,\mu}(J) &:= BC_{1-\mu}(J, E_0), \end{aligned} \tag{2.1}$$

which are themselves Banach spaces when equipped with the norms

$$\begin{aligned} \|v\|_{\mathbb{E}_{1,\mu}(J)} &:= \sup_{t \in J} t^{1-\mu} (|\dot{v}(t)|_{E_0} + |v(t)|_{E_1}), \quad \text{and} \\ \|v\|_{\mathbb{E}_{0,\mu}(J)} &:= \sup_{t \in J} t^{1-\mu} |v(t)|_{E_0} \end{aligned}$$

respectively. Further, we note that the trace operator  $\gamma : \mathbb{E}_{1,\mu}(J) \rightarrow E_0$  is well-defined and, assuming  $\mathcal{M}_\mu(E_1, E_0) \neq \emptyset$  (as we do throughout), the trace space  $\gamma \mathbb{E}_{1,\mu}(J)$  coincides with the continuous interpolation space  $E_\mu := (E_0, E_1)_{\mu, \infty}^0$ .

*Remark 2.1. (a)* The important inequality (1.3) should be viewed in relation to applications of interpolation we will encounter frequently in the article. In particular, if we set

$$\alpha := \frac{\beta - \mu}{1 - \mu} \quad \text{and} \quad \alpha_j := \frac{\beta_j - \mu}{1 - \mu}$$

then  $E_\beta = (E_\mu, E_1)_{\alpha, \infty}^0$  and  $E_{\beta_j} = (E_\mu, E_1)_{\alpha_j, \infty}^0$ , for  $j = 1, \dots, m$ . Then, given  $x, y \in E_1$  and  $t > 0$ , it follows that

$$\begin{aligned} t^{1-\mu} |x|_{E_\beta}^{\rho_j} |y|_{E_{\beta_j}} &\leq c_j t^{1-\mu} (|x|_{E_\mu}^{\rho_j(1-\alpha)} |x|_{E_1}^{\rho_j \alpha}) (|y|_{E_\mu}^{(1-\alpha_j)} |y|_{E_1}^{\alpha_j}) \\ &\leq C_0 t^{(1-\mu)(1-\rho_j \alpha - \alpha_j)} |x|_{E_\mu}^{\rho_j(1-\alpha)} |t^{1-\mu} x|_{E_1}^{\rho_j \alpha} |y|_{E_\mu}^{(1-\alpha_j)} |t^{1-\mu} y|_{E_1}^{\alpha_j}. \end{aligned} \tag{2.2}$$

Here the constant  $c_j = c_j(\alpha, \alpha_j)$  is the product of interpolation constants from  $E_\beta$  and  $E_{\beta_j}$  (c.f. [1, Proposition 2.2.1]), while  $C_0 = C_0(\alpha, \alpha_1, \dots, \alpha_m)$  is an upper bound for the family of all such constants,  $j = 1, \dots, m$ .

*(b)* In the proof of Theorem 2.2 below, we address both subcritical and critical indices  $j$ . The difference in approaches to these two cases can be viewed in context of (2.2). In particular, note that when  $j$  is subcritical, the exponent

$$(1 - \mu)(1 - \rho_j \alpha - \alpha_j)$$

on  $t$  is strictly positive, since  $\rho_j \alpha + \alpha_j$  is exactly the left-hand side of (1.3). Meanwhile, when  $j$  is critical we have a trivial exponent on  $t$ , but it must hold that  $\rho_j > 0$  in this case. Thus, when  $j$  is critical we focus on the term  $|t^{1-\mu} x|_{E_1}^{\rho_j \alpha}$  which has a positive exponent (a property not necessarily holding in the subcritical case).

**Theorem 2.2.** *Suppose  $A, F_1$  and  $F_2$  satisfy conditions (H1)–(H2).*

*(a) (Local Solutions)* *Given any  $x_0 \in V_\mu$ , there exist positive constants  $\tau = \tau(x_0)$ ,  $\varepsilon = \varepsilon(x_0)$ , and  $\sigma = \sigma(x_0)$  such that (1.1) has a unique solution*

$$u(\cdot, x) \in \mathbb{E}_{1,\mu}([0, \tau]) := BC_{1-\mu}^1([0, \tau], E_0) \cap BC_{1-\mu}([0, \tau], E_1)$$

*for all initial values  $x \in \bar{B}_{E_\mu}(x_0, \varepsilon)$ . Moreover,*

$$\|u(\cdot, x_1) - u(\cdot, x_2)\|_{\mathbb{E}_{1,\mu}([0, \tau])} \leq \sigma |x_1 - x_2|_{E_\mu}, \quad x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon). \tag{2.3}$$

- (b) **(Maximal Solutions)** *Each solution with initial value  $x_0 \in V_\mu$  exists on a maximal interval  $J(x_0) := [0, t^+) = [0, t^+(x_0))$  and enjoys the regularity*

$$u(\cdot, x_0) \in C([0, t^+), E_\mu) \cap C((0, t^+), E_1).$$

- (c) **(Global Solutions)** *If the solution  $u(\cdot, x_0)$  satisfies the conditions:*

- (i)  $u(\cdot, x_0) \in UC(J(x_0), E_\mu)$  and
- (ii) *there exists  $\eta > 0$  so that  $\text{dist}_{E_\mu}(u(t, x_0), \partial V_\mu) > \eta$  for all  $t \in J(x_0)$ , then it holds that  $t^+(x_0) = \infty$  and so  $u(\cdot, x_0)$  is a global solution of (1.1). Moreover, if the embedding  $E_1 \hookrightarrow E_0$  is compact, then condition (i) may be replaced by the assumption:*
  - (i.a) *the orbit  $\{u(t, x_0) : t \in [\tau, t^+(x_0))\}$  is bounded in  $E_\delta$  for some  $\delta \in (\mu, 1]$  and some  $\tau \in (0, t^+(x_0))$ .*

Before proceeding to the proof of the theorem we add some remarks. We first note that the embedding  $\mathbb{E}_{1,\mu}([0, \tau]) \hookrightarrow C([0, \tau], E_\mu)$ , see [5, Lemma 2.2(b)], immediately implies the following result.

**Corollary 2.3.** *Under the assumptions of Theorem 2.2, there exists a positive constant  $c = c(x_0)$  so that*

$$\|u(\cdot, x_1) - u(\cdot, x_2)\|_{C([0, \tau], E_\mu)} \leq c|x_1 - x_2|_{E_\mu} \quad \text{for } x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon).$$

*It thus follows that the map  $[(t, x) \mapsto u(t, x)]$  defines a locally Lipschitz continuous semiflow on  $V_\mu$ .*

**Remark 2.4. (a)** We recall briefly that local Lipschitz continuity of a semiflow on  $V_\mu$  means that

$$\mathcal{D} := \bigcup_{x \in V_\mu} [0, t^+(x)) \times \{x\}$$

is an open set in  $\mathbb{R}_+ \times V_\mu$ , the map  $[(t, x) \mapsto u(t, x)]$  is continuous on  $\mathcal{D}$ , and for all  $(t_0, x_0) \in \mathcal{D}$  there exists a product neighborhood  $U \times V \subset \mathcal{D}$  and  $c > 0$  so that

$$|u(t, x) - u(t, y)|_{V_\mu} \leq c|x - y|_{V_\mu} \quad \text{for } (t, x), (t, y) \in U \times V.$$

**(b)** Local well-posedness of (1.1) was also considered by Asai [3] in the presence of a singular right-hand side  $F : V_\mu \cap E_1 \rightarrow E_0$ . In particular, the author assumes that  $F$  satisfies

$$|F(x_1) - F(x_2)|_{E_0} \leq C_R(1 + |x_1|_{E_1}^p + |x_2|_{E_1}^p)|x_1 - x_2|_{E_\theta}$$

for all  $x_1, x_2 \in E_1 \cap B_{E_\mu}(x_0, R)$ . Here the author has  $p$  and  $E_\theta := (E_0, E_1)_{\theta, \infty}^0$  appropriately chosen, with  $\theta \in [\mu, 1)$ , so that  $p + (\theta - \mu)/(1 - \mu) < 1$ . This setting is similar to our condition **(H2)** if one allows  $j = 1$ ,  $\beta = 1$ ,  $\rho_j = p$ , and  $\beta_j = \theta$ , whereby it follows that Asai only considers subcritical weights. Further, we note that in [3, Theorem 1.1] the author proves Hölder continuous dependence on initial data in  $V_\mu$ , whereas we obtain Lipschitz continuity. No additional geometric properties for solutions are established in [3].

*Proof of Theorem 2.2.* (a) We follow the structure of related proofs in [9] and [20], where  $L_p$ -maximal regularity is assumed. We note that sub-critical and critical indices required distinct proofs in [9] and [20], respectively, whereas both cases can be handled in the same setting here.

Choose  $x_0 \in V_\mu$  and fix  $\varepsilon_0 > 0$  so that

$$\bar{B}_{E_\mu}(x_0, \varepsilon_0) \subset V_\mu.$$

Applying **(H1)** and **(H2)**, we obtain constants  $L = L(\varepsilon_0) > 0$  and  $C_{\varepsilon_0} > 0$  so that

$$\|(A, F_1)(x_1) - (A, F_1)(x_2)\|_{\mathcal{L}(E_1, E_0) \times E_0} \leq L|x_1 - x_2|_{E_\mu}, \quad (2.4)$$

for  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon_0)$ , and

$$|F_2(x_1) - F_2(x_2)|_{E_0} \leq C_{\varepsilon_0} \sum_{j=1}^m (1 + |x_1|_{E_\beta}^{\rho_j} + |x_2|_{E_\beta}^{\rho_j}) |x_1 - x_2|_{E_{\beta_j}} \quad (2.5)$$

for  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon_0) \cap E_\beta$ .

It follows from **(H1)** and [10, Corollary 1] that  $-A(x_0)$  generates a strongly continuous analytic semigroup on  $E_0$  with domain  $E_1$ . For each element  $x \in V_\mu$ , we define

$$u_x^*(t) := e^{-tA(x_0)}x$$

which is in  $\mathbb{E}_{1,\mu}(J_T)$ , for any  $T > 0$  and solves  $\dot{u} + A(x_0)u = 0$ ,  $u(0) = x$ .

Furthermore, we fix positive constants  $T_1 > 0$ ,  $M_1 \geq 1$ , and  $C_1 > 0$  so that, for all  $J_T := [0, T] \subset J_{T_1}$  and  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon_0)$ , we have

$$\|u_{x_0}^* - x_0\|_{C(J_T, E_\mu)} < \frac{\varepsilon_0}{3}, \quad (2.6)$$

$$\|u\|_{C(J_T, E_\mu)} \leq M_1 \|u\|_{\mathbb{E}_{1,\mu}(J_T)}, \quad \text{for } u \in \mathbb{E}_{1,\mu}(J_T) \text{ with } u(0) = 0, \quad (2.7)$$

and

$$M_1 \|u_{x_1}^* - u_{x_2}^*\|_{\mathbb{E}_{1,\mu}(J_T)} + \|u_{x_1}^* - u_{x_2}^*\|_{C(J_T, E_\mu)} \leq C_1 |x_1 - x_2|_{E_\mu}. \quad (2.8)$$

The previous inequalities are justified by strong continuity of the semigroup  $e^{-tA(x_0)}$  in  $E_\mu$ , [5, Lemma 2.2(c)], and [5, Equation (3.7)], respectively.

We will construct a contraction mapping on a closed subset of  $\mathbb{E}_{1,\mu}(J_T)$  given by

$$W_x(J_T, r) := \{v \in \mathbb{E}_{1,\mu}(J_T) : v(0) = x, \text{ and } \|v - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \leq r\}, \quad (2.9)$$

where  $x \in E_\mu$ ,  $J_T \subset J_{T_1}$ , and  $r > 0$ . The mapping we consider will be  $\mathcal{T}_x$  which takes  $v \in W_x(J_T, r)$  to the solution  $w = \mathcal{T}_x(v)$  of the linear initial value problem

$$\begin{cases} \dot{w} + A(x_0)w = (A(x_0) - A(v))v + F_1(v) + F_2(v), & t \in J_T, \\ w(0) = x. \end{cases} \quad (2.10)$$

Note that fixed points  $v = \mathcal{T}_x(v)$  are solutions to the original problem (1.1) on  $J_T$ . We proceed by first proving that  $\mathcal{T}_x$  is well-defined (see Claims 1 and 2 below), then we show that  $\mathcal{T}_x$  is in fact a contraction mapping on  $W_x(J_T, r)$  for  $r, T$  and  $x$  appropriately chosen (see Claims 3 and 4 below).

**Claim 1:** For  $r, T, \varepsilon$  chosen sufficiently small and positive, if  $x \in \bar{\mathbb{B}}_{E_\mu}(x_0, \varepsilon)$  then  $\|v - x_0\|_{C(J_T, E_\mu)} \leq \varepsilon_0$  for all  $v \in W_x(J_T, r)$ .



*Proof of Claim 1.* For any  $v \in W_x(J_T, r)$  note that  $v(0) - u_x^*(0) = 0$ , thus (2.7) and the triangle inequality imply

$$\|v - u_x^*\|_{C(J_T, E_\mu)} \leq M_1 \|v - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} + M_1 \|u_{x_0}^* - u_x^*\|_{\mathbb{E}_{1,\mu}(J_T)}.$$

Applying (2.6), (2.8), and (2.9), we compute

$$\begin{aligned} \|v - x_0\|_{C(J_T, E_\mu)} &\leq \|v - u_x^*\|_{C(J_T, E_\mu)} + \|u_x^* - u_{x_0}^*\|_{C(J_T, E_\mu)} + \|u_{x_0}^* - x_0\|_{C(J_T, E_\mu)} \\ &\leq M_1 r + C_1 |x - x_0|_{E_\mu} + \|u_{x_0}^* - x_0\|_{C(J_T, E_\mu)} \\ &\leq M_1 r + C_1 \varepsilon + \varepsilon_0/3. \end{aligned} \quad (2.11)$$

Claim 1 thus follows by restricting  $r$  and  $\varepsilon$  appropriately so that the last line in (2.11) is bounded by  $\varepsilon_0$ .  $\blacksquare$

Henceforth, we assume  $x$  is sufficiently close to  $x_0$  (in  $E_\mu$ ) and  $r, T, \varepsilon$  are given appropriately small so that Claim 1 holds. It follows that given any  $v_1, v_2 \in W_x(J_T, r)$ , the structural conditions (2.4)–(2.5) hold for  $v_1(s), v_2(t)$ , with  $s, t \in \dot{J}_T$ .

**Claim 2:**  $(A(x_0) - A(v))v, F_1(v), F_2(v) \in \mathbb{E}_{0,\mu}(J_T)$  for each  $v \in W_x(J_T, r)$ .

*Proof of Claim 2.* Regarding regularity of  $(A(x_0) - A(v))v$  and  $F_1(v)$ , for  $v \in W_x(J_T, r)$ , note that continuity of each function into  $E_0$  follows from (2.4) and the fact that  $v \in \mathbb{E}_{1,\mu}(J_T) \hookrightarrow C(J_T, E_\mu)$ . Employing **(H1)** and the bounds (2.4) and (2.11), we compute

$$\begin{aligned} t^{1-\mu} |(A(x_0) - A(v(t)))v(t)|_{E_0} &\leq L |x_0 - v(t)|_{E_\mu} t^{1-\mu} |v(t)|_{E_1} \\ &\leq L \|v - x_0\|_{C(J_T, E_\mu)} \|v\|_{\mathbb{E}_{1,\mu}(J_T)} \\ &\leq L (M_1 r + C_1 \varepsilon + \|u_{x_0}^* - x_0\|_{C(J_T, E_\mu)}) (r + \|u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}) \end{aligned} \quad (2.12)$$

and

$$\begin{aligned} t^{1-\mu} |F_1(v(t))|_{E_0} &\leq t^{1-\mu} |F_1(v(t)) - F_1(x_0)|_{E_0} + t^{1-\mu} |F_1(x_0)|_{E_0} \\ &\leq t^{1-\mu} L |v(t) - x_0|_{E_\mu} + t^{1-\mu} |F_1(x_0)|_{E_0} \\ &\leq T^{1-\mu} (L \varepsilon_0 + |F_1(x_0)|_{E_0}). \end{aligned} \quad (2.13)$$

From (2.12) and (2.13), we draw the following conclusions. For each  $v \in W_x(J_T, r)$ , we see that  $t^{1-\mu} |F_1(v(t))|_{E_0}$  and  $t^{1-\mu} |(A(x_0) - A(v(t)))v(t)|_{E_0}$  are bounded on  $\dot{J}_T$ . Further, as  $T \rightarrow 0^+$  notice that  $\|v\|_{\mathbb{E}_{1,\mu}(J_T)} \rightarrow 0$  and  $T^{1-\mu} \rightarrow 0$ , from which we conclude

$$(A(x_0) - A(v))v, F_1(v) \in \mathbb{E}_{0,\mu}(J_T), \quad \text{for all } v \in W_x(J_T, r). \quad (2.14)$$

As an additional observation, note (2.12) and (2.13) imply  $\|(A(x_0) - A(v))v\|_{\mathbb{E}_{0,\mu}(J_T)}$  and  $\|F_1(v)\|_{\mathbb{E}_{0,\mu}(J_T)}$  are uniformly bounded on  $W_x(J_T, r)$ .

Lastly, we consider the term  $F_2(v)$ , and we first observe that

$$F_2(v) \in C(\dot{J}_T, E_0) \quad \text{for each } v \in W_x(J_T, r). \quad (2.15)$$

Indeed, let  $v \in W_x(J_T, r)$  be given. Then  $v \in C(\dot{J}_T, E_\beta)$ , and by Claim 1, we also know that  $v(t) \in \bar{B}_{E_\mu}(x_0, \varepsilon_0)$  for all  $t \in J_T$ . Hence we have for each  $s, t \in \dot{J}_T$

$$|F_2(v(t)) - F_2(v(s))|_{E_0} \leq C_{\varepsilon_0} \sum_{j=1}^m (1 + |v(t)|_{E_\beta}^{\rho_j} + |v(s)|_{E_\beta}^{\rho_j}) |v(t) - v(s)|_{E_{\beta_j}}.$$

The assertion in (2.15) now follows from the embedding  $E_\beta \hookrightarrow E_{\beta_j}$  and the observation that  $|v(t)|_{E_\beta}$  and  $|v(s)|_{E_\beta}$  are bounded for values  $s, t$  that are bounded away from 0. The latter statement means that for each  $\eta \in \dot{J}_T$  there is a constant  $C_\eta > 0$  such that  $|v(t)|_{E_\beta}, |v(s)|_{E_\beta} \leq C_\eta$  for all  $s, t \in [\eta, T]$ .

In order to show boundedness of  $t^{1-\mu}|F_2(v(t))|_{E_0}$  we choose  $y \in E_1 \cap B_{E_\mu}(x_0, \varepsilon_0)$ , which is feasible by the density of the embedding  $E_1 \hookrightarrow E_\mu$ , and write

$$F_2(v(t)) = (F_2(v(t)) - F_2(y)) + F_2(y), \quad t \in \dot{J}_T.$$

Clearly,  $F_2(y) \in \mathbb{E}_{0,\mu}(J_T)$ . To treat the term  $F_2(v(t)) - F_2(y)$ , we first observe that

$$\begin{aligned} |v(t) - y|_{E_{\beta_j}} &\leq c_j t^{-(1-\mu)\alpha_j} |v(t) - y|_{E_\mu}^{1-\alpha_j} (t^{1-\mu} |v(t) - y|_{E_1})^{\alpha_j} \\ &\leq c_j t^{-(1-\mu)\alpha_j} |v(t) - y|_{E_\mu}^{1-\alpha_j} \|v - y\|_{\mathbb{E}_{1,\mu}(J_T)}^{\alpha_j} \leq \tilde{c} t^{-(1-\mu)\alpha_j} \end{aligned} \quad (2.16)$$

for each  $t \in \dot{J}_T$ . Next, employing (2.2) and (2.16), we have

$$\begin{aligned} t^{1-\mu} |F_2(v(t)) - F_2(y)|_{E_0} &\leq C_{\varepsilon_0} t^{1-\mu} \sum_{j=1}^m (1 + |y|_{E_\beta}^{\rho_j} + |v(t)|_{E_\beta}^{\rho_j}) |v(t) - y|_{E_{\beta_j}} \\ &\leq \tilde{C} \left[ \sum_{j=1}^m t^{(1-\mu)(1-\alpha_j)} (1 + |y|_{E_\beta}^{\rho_j}) + \sum_{j \text{ critical}} \|v\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right. \\ &\quad \left. + \sum_{j \text{ subcritical}} t^{(1-\mu)(1-\rho_j\alpha-\alpha_j)} \|v\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right]. \end{aligned}$$

Since  $\|v\|_{\mathbb{E}_{1,\mu}(J_T)} \rightarrow 0$  as  $T \rightarrow 0^+$ , we see that  $t^{1-\mu}F_2(v(t))$  is bounded in  $E_0$  and  $\|F_2(v)\|_{\mathbb{E}_{0,\mu}(J_T)}$  converges to zero as  $T \rightarrow 0^+$ . Claim 2 is thus proved.  $\blacksquare$

With  $T, r$  and  $x$  chosen as above, we have now shown that the right hand side of equation (2.10) is in  $\mathbb{E}_{0,\mu}(J_T)$  for all  $v \in W_x(J_T, r)$ . Thus, since  $A(x_0) \in \mathcal{M}_\mu(E_1, E_0)$ , we conclude that  $\mathcal{T}_x : W_x(J_T, r) \rightarrow \mathbb{E}_{1,\mu}(J_T)$  is well-defined. Now, to conclude the proof of the theorem, we must show that  $\mathcal{T}_x$  is a contraction mapping on  $W_x(J_T, r)$  for appropriately chosen  $r, T$ , and  $x$ .

**Claim 3:** For  $r, T$ , and  $\varepsilon$  chosen sufficiently small and positive, if  $x \in \bar{B}_{E_\mu}(x_0, \varepsilon)$ , then  $\mathcal{T}_x$  maps  $W_x(J_T, r)$  into itself.

*Proof of Claim 3.* For  $v \in W_x(J_T, r)$ , notice that  $\mathcal{T}_x(v) \in \mathbb{E}_{1,\mu}(J_T)$  and  $\mathcal{T}_x(v)(0) = x$ , by the property of maximal regularity and the definition of the mapping  $\mathcal{T}_x$ . Thus, it suffices to show that  $\|\mathcal{T}_x(v) - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \leq r$  holds for all  $v \in W_x(J_T, r)$ , provided  $r, T$ , and  $\varepsilon$  are chosen sufficiently small.

We begin with the observation

$$\|\mathcal{T}_x(v) - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \leq \|\mathcal{T}_x(v) - u_x^*\|_{\mathbb{E}_{1,\mu}(J_T)} + \|u_x^* - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}. \quad (2.17)$$

Applying (2.8) we get

$$\|u_x^* - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \leq C_1 |x - x_0|_{E_\mu},$$

using the fact that  $M_1 \geq 1$ . Note that choosing  $\varepsilon$  sufficiently small, this term can be bounded by  $r/4$  for all  $x \in \bar{B}_{E_\mu}(x_0, \varepsilon)$ .

Since  $(\mathcal{T}_x(v) - u_x^*)|_{t=0} = 0$ , we apply maximal regularity of  $A(x_0)$  to bound the first term of (2.17)

$$\begin{aligned} & \|\mathcal{T}_x(v) - u_x^*\|_{\mathbb{E}_{1,\mu}(J_T)} \\ & \leq C_{T_1} (\|(A(x_0) - A(v))v\|_{\mathbb{E}_{0,\mu}(J_T)} + \|F_1(v)\|_{\mathbb{E}_{0,\mu}(J_T)} + \|F_2(v)\|_{\mathbb{E}_{0,\mu}(J_T)}), \end{aligned} \quad (2.18)$$

where  $C_{T_1} > 0$  is the constant of maximal regularity for the interval  $[0, T_1]$ ; recalling that  $T_1 > 0$  was introduced before bounds (2.6)–(2.8). The first two terms of (2.18) are bounded as in (2.12) and (2.13), respectively — which are both bounded by  $r/4C_{T_1}$  for  $r, T$ , and  $\varepsilon$  sufficiently small.

Addressing the last term in (2.18), we first split

$$\|F_2(v)\|_{\mathbb{E}_{0,\mu}(J_T)} \leq \|F_2(v) - F_2(u_{x_0}^*)\|_{\mathbb{E}_{0,\mu}(J_T)} + \|F_2(u_{x_0}^*)\|_{\mathbb{E}_{0,\mu}(J_T)},$$

then note that  $\|F_2(u_{x_0}^*)\|_{\mathbb{E}_{0,\mu}(J_T)}$  can be made arbitrarily small by taking  $T$  sufficiently small. Meanwhile, we apply (2.5) and Remark 2.1 to bound

$$\begin{aligned} & \|F_2(v) - F_2(u_{x_0}^*)\|_{\mathbb{E}_{0,\mu}(J_T)} \\ & \leq M_0 \sum_{j=1}^m \left[ T^{(1-\mu)(1-\alpha_j)} + T^{(1-\mu)(1-\rho_j\alpha-\alpha_j)} \left( \|v\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right. \right. \\ & \quad \left. \left. + \|u_{x_0}^*\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right) \right] \|v - u_{x_0}^*\|_{C(J_T, E_\mu)}^{1-\alpha_j} \|v - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}^{\alpha_j} \\ & \leq M_0 \sum_{j=1}^m \left[ T^{(1-\mu)(1-\alpha_j)} + T^{(1-\mu)(1-\rho_j\alpha-\alpha_j)} (2M_3 r^{\rho_j\alpha}) \right] M_2 r, \end{aligned} \quad (2.19)$$

where  $M_0 := C_{\varepsilon_0} C_0$  and  $M_2, M_3$  are constants chosen as follows. Applying part of (2.11) and Young's inequality, we select  $M_2 > 0$  so that

$$\begin{aligned} & \|v - u_{x_0}^*\|_{C(J_T, E_\mu)}^{1-\alpha_j} \|v - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}^{\alpha_j} \\ & \leq (1 - \alpha_j) \|v - u_{x_0}^*\|_{C(J_T, E_\mu)} + \alpha_j \|v - u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \\ & \leq (1 - \alpha_j)(M_1 r + C_1 \varepsilon) + \alpha_j r \\ & \leq M_2 r, \end{aligned} \quad (2.20)$$

for  $\varepsilon \leq r$ . Likewise, applying Claim 1, we select  $M_3 > 0$  so that

$$\begin{aligned} & \|v\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \\ & \leq (\varepsilon_0 + |x_0|_{E_\mu})^{\rho_j(1-\alpha)} (r + \|u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)})^{\rho_j\alpha} \\ & \leq M_3 r^{\rho_j\alpha}, \end{aligned} \quad (2.21)$$

for all  $T > 0$  sufficiently small so that  $\|u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \leq r$ .

Finally, note that all terms in (2.19) have a linear factor of  $r$  and an additional factor that can be made arbitrarily small by restricting the sizes of  $r, T$ , and  $\varepsilon$ . In context of Remark 2.1(b), we note that terms involving subcritical index  $j$  are made small with  $T$  alone, while critical indices  $j$  require restriction on the size of  $r$ . We

conclude that the last term in (2.18) can be bounded by  $r/4$  and (2.17) can thus be bounded by  $r$  for all  $r, T$ , and  $\varepsilon$  chosen sufficiently small. This proves Claim 3.  $\blacksquare$

**Claim 4:** There exist constants  $\kappa = \kappa(r, T, \varepsilon) > 0$  and  $\sigma = \sigma(r, T, \varepsilon) > 0$  so that for all  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon)$  and  $v_i \in W_{x_i}(J_T, r)$ ,  $i = 1, 2$ , it holds that

$$\|\mathcal{T}_{x_1}(v_1) - \mathcal{T}_{x_2}(v_2)\|_{\mathbb{E}_{1,\mu}(J_T)} \leq \kappa \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + \sigma |x_1 - x_2|_{E_\mu}.$$

Further,  $\kappa$  is made arbitrarily small by choosing  $r, T$ , and  $\varepsilon$  sufficiently small.

*Proof of Claim 4.* Let  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon)$  be given and pick  $v_1 \in W_{x_1}(J_T, r)$ ,  $v_2 \in W_{x_2}(J_T, r)$ . By (2.7) and (2.8) we have

$$\begin{aligned} \|v_1 - v_2\|_{C(J_T, E_\mu)} &\leq \|(v_1 - v_2) - (u_{x_1}^* - u_{x_2}^*)\|_{C(J_T, E_\mu)} + \|u_{x_1}^* - u_{x_2}^*\|_{C(J_T, E_\mu)} \\ &\leq M_1 \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + C_1 |x_1 - x_2|_{E_\mu}. \end{aligned} \quad (2.22)$$

Proceeding, we first note that  $((\mathcal{T}_{x_1}(v_1) - \mathcal{T}_{x_2}(v_2)) - (u_{x_1}^* - u_{x_2}^*))|_{t=0} = 0$ , so we compute, applying maximal regularity of  $A(x_0)$  and (2.8),

$$\begin{aligned} &\|\mathcal{T}_{x_1}(v_1) - \mathcal{T}_{x_2}(v_2)\|_{\mathbb{E}_{1,\mu}(J_T)} \\ &\leq \|(\mathcal{T}_{x_1}(v_1) - \mathcal{T}_{x_2}(v_2)) - (u_{x_1}^* - u_{x_2}^*)\|_{\mathbb{E}_{1,\mu}(J_T)} + \|u_{x_1}^* - u_{x_2}^*\|_{\mathbb{E}_{1,\mu}(J_T)} \\ &\leq C_{T_1} \left[ \|(A(v_2) - A(v_1))v_2\|_{\mathbb{E}_{0,\mu}(J_T)} + \|(A(x_0) - A(v_1))(v_1 - v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} \right. \\ &\quad \left. + \|F_1(v_1) - F_1(v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} + \|F_2(v_1) - F_2(v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} \right] \\ &\quad + C_1 |x_1 - x_2|_{E_\mu}. \end{aligned} \quad (2.23)$$

Continuing with individual terms in (2.23), we apply (2.4) and (2.22) to get

$$\begin{aligned} &\|(A(v_2) - A(v_1))v_2\|_{\mathbb{E}_{0,\mu}(J_T)} \\ &\leq L \|v_1 - v_2\|_{C(J_T, E_\mu)} \|v_2\|_{\mathbb{E}_{1,\mu}(J_T)} \\ &\leq L \left( M_1 \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + C_1 |x_1 - x_2|_{E_\mu} \right) (r + \|u_{x_0}^*\|_{\mathbb{E}_{1,\mu}(J_T)}), \\ &\|F_1(v_1) - F_1(v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} \\ &\leq T^{1-\mu} L \left( M_1 \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + C_1 |x_1 - x_2|_{E_\mu} \right), \end{aligned}$$

and, also applying (2.11), we have

$$\begin{aligned} &\|(A(x_0) - A(v_1))(v_1 - v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} \\ &\leq L \|v_1 - x_0\|_{C(J_T, E_\mu)} \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} \\ &\leq L \left( M_1 r + C_1 \varepsilon + \|u_{x_0}^* - x_0\|_{C(J_T, E_\mu)} \right) \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)}. \end{aligned}$$

Meanwhile, by Young's inequality and (2.22), recalling that  $M_1 \geq 1$ , we have

$$\begin{aligned} &\|v_1 - v_2\|_{C(J_T, E_\mu)}^{1-\alpha_j} \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)}^{\alpha_j} \\ &\leq M_1 \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + C_1 |x_1 - x_2|_{E_\mu}, \end{aligned}$$

which we apply in combination with (2.5), Remark 2.1, and (2.21) to bound

$$\begin{aligned}
& \|F_2(v_1) - F_2(v_2)\|_{\mathbb{E}_{0,\mu}(J_T)} \\
& \leq M_0 \sum_{j=1}^m \left[ T^{(1-\mu)(1-\alpha_j)} + T^{(1-\mu)(1-\rho_j\alpha-\alpha_j)} \left( \|v_1\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v_1\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right. \right. \\
& \quad \left. \left. + \|v_2\|_{C(J_T, E_\mu)}^{\rho_j(1-\alpha)} \|v_2\|_{\mathbb{E}_{1,\mu}(J_T)}^{\rho_j\alpha} \right) \right] \|v_1 - v_2\|_{C(J_T, E_\mu)}^{1-\alpha_j} \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)}^{\alpha_j} \\
& \leq M_0 \sum_{j=1}^m \left[ T^{(1-\mu)(1-\alpha_j)} + T^{(1-\mu)(1-\rho_j\alpha-\alpha_j)} (2M_3 r^{\rho_j\alpha}) \right] \\
& \quad \cdot (M_1 \|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)} + C_1 |x_1 - x_2|_{E_\mu}),
\end{aligned}$$

with  $M_0 := C_{\varepsilon_0} C_0$ . Combining all terms involving  $\|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)}$  and likewise terms involving  $|x_1 - x_2|_{E_\mu}$ , note that (2.23) takes on the desired structure for the claim. Moreover, every factor multiplying the terms  $\|v_1 - v_2\|_{\mathbb{E}_{1,\mu}(J_T)}$  can be made arbitrarily small by taking either  $r, T$ , or  $\varepsilon$  sufficiently small. Note that the same cannot be said for every factor of  $|x_1 - x_2|_{E_\mu}$ , as seen in the last term of (2.23). Regardless, we have thus proved Claim 4.  $\blacksquare$

Finally, fix  $r, T$ , and  $\varepsilon$  small enough so that  $\kappa \leq \frac{1}{2}$ . We thus have the estimate

$$\|\mathcal{T}_{x_1}(v_1) - \mathcal{T}_{x_2}(v_2)\|_{\mathbb{E}_{1,\mu}(J_T)} \leq \frac{1}{2} \|v_1 - v_2\|_{\mathbb{E}_{0,\mu}(J_T)} + \sigma |x_1 - x_2|_{E_\mu}, \quad (2.24)$$

for every  $v_i \in W_{x_i}(J_T, r)$  and  $x_i \in \bar{B}_{E_\mu}(x_0, \varepsilon)$ . Let  $x_1 = x_2 = x \in B_{E_\mu}(x_0, \varepsilon)$  be given. Then

$$\|\mathcal{T}_x(v_1) - \mathcal{T}_x(v_2)\|_{\mathbb{E}_{1,\mu}(J_T)} \leq \frac{1}{2} \|v_1 - v_2\|_{\mathbb{E}_{0,\mu}(J_T)} \quad \text{for } v_1, v_2 \in W_x(J_T, r),$$

and so  $\mathcal{T}_x$  is a strict contraction on  $W_x(J_T, r)$ . Applying Banach's fixed point theorem, we obtain a unique fixed point

$$u(\cdot, x) \in W_x(J_T, r) \subset \mathbb{E}_{1,\mu}(J_T), \quad \text{for every } x \in \bar{B}_{E_\mu}(x_0, \varepsilon),$$

which solves (1.1) by construction of the mapping  $\mathcal{T}_x$ . Furthermore, for  $x_1, x_2 \in \bar{B}_{E_\mu}(x_0, \varepsilon)$ , (2.24) implies

$$\|u(\cdot, x_1) - u(\cdot, x_2)\|_{\mathbb{E}_{1,\mu}(J_T)} \leq 2\sigma |x_1 - x_2|_{E_\mu},$$

which completes the proof of the first statement of the theorem.

(b) By a standard argument, we can extend the local solution obtained in part (a) to a maximal solution on some right-open interval  $[0, t^+(x_0))$ . To confirm this maximal solution satisfies the stated regularity, we consider a portion of this extension argument. In particular, with  $x_0 \in V_\mu$  given, we apply part (a) to produce the solution  $u_1(\cdot, x_0) \in \mathbb{E}_{1,\mu}([0, \tau_1])$  on some interval  $[0, \tau_1]$ . Then, we note that  $x_1 := u_1(\tau_1, x_0) \in E_1 \cap V_\mu$ , and so we may apply part (a) again to produce the solution  $u_2(\cdot, x_1) \in \mathbb{E}_{1,\mu}([0, \tau_2])$  on a second interval  $[0, \tau_2]$ . It follows that

$$u(t) := \begin{cases} u_1(t), & \text{for } t \in [0, \tau_1] \\ u_2(t - \tau_1), & \text{for } t \in [\tau_1, \tau_1 + \tau_2] \end{cases}$$

satisfies (1.1) with  $u(0) = x_0$  and regularity  $u \in \mathbb{E}_{1,\mu}([0, \tau_1 + \tau_2])$ . To prove this last claim, it suffices to show that  $u_2 \in C([0, \tau_2], E_1)$ , in particular  $\lim_{t \rightarrow 0^+} u_2(t) = x_1$ . For that purpose, we fix  $\varepsilon > 0$  so that the result of part (a) holds for  $x \in \bar{B}_{E_\mu}(x_1, \varepsilon)$  and choose  $\delta \in (0, \tau_2)$  sufficiently small that  $u_1(\tau_1 - \delta) \in \bar{B}_{E_\mu}(x_1, \varepsilon)$ . Now let  $v \in \mathbb{E}_{1,\mu}([0, \tau_2])$  denote the solution to (1.1) with initial value  $u_1(\tau_1 - \delta)$ . By uniqueness of solutions, it follows that

$$v|_{[0, \delta]} = u_1|_{[\tau_1 - \delta, \tau_1]} \quad \text{and} \quad v|_{[\delta, \tau_2]} = u_2|_{[0, \tau_2 - \delta]},$$

and the desired regularity of  $u_2$  now follows by the regularity of  $v \in \mathbb{E}_{1,\mu}([0, \tau_2])$  away from  $t = 0$ .

(c) The proof of global existence follows exactly as in [5, Theorem 4.1 (c)–(d)], with the regularity of maximal solutions confirmed in (b) above.  $\square$

We conclude this section on well-posedness with the following extension of (2.3), accounting for the dependence of solutions on initial data residing in smaller spaces  $E_{\bar{\mu}} \subset E_\mu$ . This result will be useful in the following section as we consider long-term dynamics of solutions that start in  $E_\mu$  and instantaneously regularize to spaces  $E_{\bar{\mu}}$ .

**Proposition 2.5.** *Suppose the assumptions of Theorem 2.2 hold,  $\bar{\mu} \in [\beta, 1)$ , and  $x_0 \in V_\mu \cap E_{\bar{\mu}}$ . Then Theorem 2.2(a) holds true with  $\mu$  replaced by  $\bar{\mu}$ . In particular,*

$$\|u(\cdot, x_1) - u(\cdot, x_2)\|_{\mathbb{E}_{1,\bar{\mu}}([0, \tau])} \leq \sigma |x_1 - x_2|_{E_{\bar{\mu}}}, \quad x_1, x_2 \in \bar{B}_{E_{\bar{\mu}}}(x_0, \varepsilon). \quad (2.25)$$

*Proof.* We recall that  $\mathcal{M}_\mu(E_1, E_0) \subset \mathcal{M}_{\bar{\mu}}(E_1, E_0)$ , see [5, Lemma 2.6]. Hence it follows from **(H1)**–**(H2)** that

$$(A, F) \in C^{1-}(V_\mu \cap E_{\bar{\mu}}, \mathcal{M}_{\bar{\mu}}(E_1, E_0) \times E_0), \quad \text{where } F := F_1 + F_2. \quad (2.26)$$

Existence of a unique solution  $u_{\bar{\mu}} = u_{\bar{\mu}}(\cdot, x) \in \mathbb{E}_{1,\bar{\mu}}([0, \tau])$  with property (2.25) follows as in the proof of Theorem 2.2(a), with  $F_1 = F$  and  $F_2 = 0$ . In both cases, the solution is obtained as a fixed point of a strict contraction  $\mathcal{T}_x : \mathbf{M}_\nu \rightarrow \mathbf{M}_\nu$ , where  $\mathbf{M}_\nu$  is a closed subset of  $\mathbb{E}_{1,\nu}([0, \tau])$ , respectively, with  $\nu \in \{\mu, \bar{\mu}\}$ . But

$$\mathcal{T}_x : \mathbf{M}_\mu \cap \mathbf{M}_{\bar{\mu}} \rightarrow \mathbf{M}_\mu \cap \mathbf{M}_{\bar{\mu}}$$

is a strict contraction as well, and thus has a unique fixed point  $u_\star \in \mathbf{M}_\mu \cap \mathbf{M}_{\bar{\mu}}$ . Therefore,  $u_\mu = u_{\bar{\mu}} = u_\star$  on  $[0, \tau]$ , where  $u_\mu = u_\mu(\cdot, x) \in \mathbb{E}_{1,\mu}([0, \tau])$  is the solution obtained in Theorem 2.2(a). This shows, in particular, that each solution  $u(\cdot, x)$ , with  $x \in B_{E_{\bar{\mu}}}(x_0, \varepsilon)$ , obtained in Theorem 2.2(a), also belongs to  $\mathbb{E}_{1,\bar{\mu}}([0, \tau])$ .  $\square$

### 3. NORMAL STABILITY

With well-posedness of (1.1) established, we investigate the long-term behavior of solutions that start near equilibria. In particular, in this section we demonstrate that the so-called generalized principle of linearized stability (c.f. [18, 19]) continues to hold on  $E_\mu$ , provided the pertinent assumptions are satisfied. As a first step in this direction, we prove that stability of equilibria can be tracked in either the topology of  $E_\mu$  or, equivalently, in the stronger topology of  $E_{\bar{\mu}}$ .

**Proposition 3.1.** *Suppose the assumptions of Theorem 2.2 hold,  $\bar{\mu} \in [\beta, 1)$ , and suppose  $u_\star \in V_\mu \cap E_1$  is an equilibrium for (1.1). Then  $u_\star$  is stable in the topology of  $E_\mu \iff u_\star$  is stable in the topology of  $E_{\bar{\mu}}$ .*

*Proof.* By Theorem 2.2(a) and Proposition 2.5 there are constants  $\tau = \tau(u_*)$ ,  $\eta = \eta(u_*)$  and  $c_1 = c_1(u_*)$ , corresponding to the initial value  $u_*$ , such that

$$\|u(\cdot, y_0) - u_*\|_{\mathbb{E}_{1,\nu}([0,2\tau])} \leq c_1 |y_0 - u_*|_{E_\nu}, \quad \nu \in \{\mu, \bar{\mu}\}, \quad (3.1)$$

for any  $y_0 \in B_{E_\nu}(u_*, \eta)$ . Moreover, one readily verifies that there is a constant  $c_2 = c_2(\tau, \mu, \bar{\mu})$  such that

$$\|v - u_*\|_{BC([\tau, 2\tau], E_{\bar{\mu}})} \leq c_2 \|v - u_*\|_{\mathbb{E}_{1,\mu}([0, 2\tau])} \quad (3.2)$$

for any function  $v \in \mathbb{E}_{1,\mu}([0, 2\tau])$ . In the sequel, we denote the embedding constant of  $E_{\bar{\mu}} \hookrightarrow E_\mu$  by  $c_\mu$ . Consequently,

$$B_{E_{\bar{\mu}}}(u_*, \alpha) \subset B_{E_\mu}(u_*, c_\mu \alpha). \quad (3.3)$$

Suppose  $u_*$  is stable in  $E_\mu$ . Let  $\varepsilon > 0$  be given and set  $\varepsilon_\mu := \min\{\varepsilon/(c_1 c_2), \eta\}$ . By assumption, there is a number  $\delta_\mu$  such that every solution of (1.1) with initial value  $x_0 \in B_{E_\mu}(u_*, \delta_\mu)$  exists globally and satisfies

$$|u(t, x_0) - u_*|_{E_\mu} < \varepsilon_\mu, \quad \text{for all } t \geq 0. \quad (3.4)$$

Next, we choose  $\delta \in (0, \delta_\mu/c_\mu]$  sufficiently small such that

$$|u(t, x_0) - u_*|_{E_{\bar{\mu}}} < \varepsilon, \quad \text{for all } t \in [0, \tau], \quad x_0 \in B_{E_{\bar{\mu}}}(u_*, \delta). \quad (3.5)$$

Here we note that (3.5) follows from continuous dependence on the initial data, see (3.1). As a consequence of (3.3), every solution  $u(\cdot, x_0)$  with  $x_0 \in B_{E_{\bar{\mu}}}(u_*, \delta)$  exists globally and satisfies (3.4) as well as (3.5). Next we will show by induction that  $u(t, x_0) \in B_{E_{\bar{\mu}}}(u_*, \varepsilon)$  for all  $t \geq 0$ . Suppose we have already shown that  $|u(t, x_0) - u_*|_{E_{\bar{\mu}}} < \varepsilon$  for  $t \in [0, (k+1)\tau]$  and  $k \in \mathbb{N}$ . We note that the case  $k = 0$  is exactly (3.5). From the definition of  $\varepsilon_\mu$  and (3.1)-(3.2) as well as (3.4) follows

$$|u(k\tau + s, x_0) - u_*|_{E_{\bar{\mu}}} \leq c_1 c_2 |u(k\tau, x_0) - u_*|_{E_\mu} < \varepsilon, \quad \tau \leq s \leq 2\tau. \quad (3.6)$$

Since this step works for any  $k \in \mathbb{N}$ , we obtain stability of  $u_*$  in  $E_{\bar{\mu}}$ .

Suppose that  $u_*$  is stable in  $E_{\bar{\mu}}$ . Let  $\varepsilon > 0$  be given and set  $\varepsilon_{\bar{\mu}} = \varepsilon/c_\mu$ . By the stability assumption, there exists a number  $\delta_{\bar{\mu}}$  such that every solution of (1.1) with initial value  $x_0 \in B_{E_{\bar{\mu}}}(u_*, \delta_{\bar{\mu}})$  exists globally and satisfies

$$|u(t, x_0) - u_*|_{E_{\bar{\mu}}} < \varepsilon_{\bar{\mu}}, \quad \text{for all } t \geq 0. \quad (3.7)$$

Next, by continuous dependence on initial data, we can choose  $\delta \in (0, \eta)$  sufficiently small such that

$$|u(t, x_0) - u_*|_{E_\mu} < \delta_{\bar{\mu}}/(c_1 c_2), \quad \text{for all } t \in [0, \tau], \quad x_0 \in B_{E_\mu}(u_*, \delta). \quad (3.8)$$

It follows from (3.1)-(3.2) and (3.8) that  $|u(\tau, x_0) - u_*|_{E_{\bar{\mu}}} \leq c_1 c_2 |x_0 - u_*|_{E_\mu} < \delta_{\bar{\mu}}$ , for all  $x_0 \in B_{E_\mu}(u_*, \delta)$ . In particular, after a short time, we are simply tracking the solutions  $u(\cdot, x_0)$  in the stronger topology of  $E_{\bar{\mu}}$ . Hence, by (3.3) and (3.7),  $u(t, x_0) \in B_{E_{\bar{\mu}}}(u_*, \varepsilon)$  for any initial value  $x_0 \in B_{E_\mu}(u_*, \delta)$ . This completes the proof of Proposition 3.1.  $\square$

In addition to **(H1)**-(**H2**) we now assume that

$$(A, F_1, F_2) \in C^1(V_\mu \cap E_{\bar{\mu}}, \mathcal{L}(E_1, E_0) \times E_0 \times E_0), \quad (3.9)$$

where  $\bar{\mu} \in [\beta, 1)$  is a fixed number. Here we note that  $V_\mu \cap E_{\bar{\mu}} \subset E_{\bar{\mu}}$  is open, and that differentiability is understood with respect to the topology of  $E_{\bar{\mu}}$ .

Let  $\mathcal{E} \subset V_\mu \cap E_1$  denote the set of equilibrium solutions of (1.1), which means that

$$u \in \mathcal{E} \text{ if and only if } u \in V_\mu \cap E_1 \text{ and } A(u)u = F_1(u) + F_2(u).$$

Given an element  $u_* \in \mathcal{E}$ , we assume that  $u_*$  is contained in an  $m$ -dimensional manifold of equilibria. This means that there is an open subset  $U \subset \mathbb{R}^m$ ,  $0 \in U$ , and a  $C^1$ -function  $\Psi : U \rightarrow E_1$ , such that

- $\Psi(U) \subset \mathcal{E}$  and  $\Psi(0) = u_*$ ,
  - the rank of  $\Psi'(0)$  equals  $m$ , and
  - $A(\Psi(\zeta))\Psi(\zeta) = F(\Psi(\zeta))$ ,  $\zeta \in U$ .
- (3.10)

We assume furthermore that near  $u_*$  there are no other equilibria than those given by  $\Psi(U)$ , i.e.  $\mathcal{E} \cap B_{E_1}(u_*, r_1) = \Psi(U)$ , for some  $r_1 > 0$ .

For  $u_* \in \mathcal{E}$ , we define

$$A_0 v = A(u_*)v + (A'(u_*)v)u_* - F'_1(u_*)v - F'_2(u_*)v, \quad v \in E_1, \quad (3.11)$$

where  $A'$ ,  $F'_1$  and  $F'_2$  denote the Fréchet derivatives of the respective functions. We denote by  $N(A_0)$  and  $R(A_0)$  the kernel and range, respectively, of the operator  $A_0$ .

After these preparations we can state the following result on convergence of solutions starting near  $u_*$ .

**Theorem 3.2.** *Suppose  $u_* \in V_\mu \cap E_1$  is an equilibrium of (1.1), and suppose that the functions  $(A, F_1, F_2)$  satisfy (H1)-(H2) as well as (3.9). Finally, suppose that  $u_*$  is normally stable, i.e.,*

- (i) *near  $u_*$  the set of equilibria  $\mathcal{E}$  is a  $C^1$ -manifold in  $E_1$  of dimension  $m \in \mathbb{N}$ ,*
- (ii) *the tangent space for  $\mathcal{E}$  at  $u_*$  is given by  $N(A_0)$ ,*
- (iii) *0 is a semi-simple eigenvalue of  $A_0$ , i.e.,  $N(A_0) \oplus R(A_0) = E_0$ ,*
- (iv)  *$\sigma(-A_0) \setminus \{0\} \subset \{z \in \mathbb{C} : \operatorname{Re} z < 0\}$ .*

*Then  $u_*$  is stable in  $E_\mu$ . Moreover, there exists a constant  $\delta = \delta(\bar{\mu}) > 0$  such that each solution  $u(\cdot, x_0)$  of (1.1) with initial value  $x_0 \in B_{E_\mu}(u_*, \delta)$  exists globally and converges to some  $u_\infty \in \mathcal{E}$  in  $E_{\bar{\mu}}$  at an exponential rate as  $t \rightarrow \infty$ .*

*Proof.* Example 2 and Theorem 3.1 in [19] imply stability of  $u_*$  in  $E_{\bar{\mu}}$ . Moreover, the same theorem ensures that there exists  $\delta_1 > 0$  such that each solution  $u(\cdot, y_0)$  of (1.1) with initial value  $y_0 \in B_{E_{\bar{\mu}}}(u_*, \delta_1)$  exists globally and converges to some  $u_\infty \in \mathcal{E}$  in the topology of  $E_{\bar{\mu}}$  at an exponential rate.

By Proposition (3.1),  $u_*$  is stable in  $E_\mu$  as well. Employing (3.1)-(3.2), we deduce that there exists  $\delta = \delta(\delta_1, \bar{\mu}) > 0$  such that

$$|u(\tau, x_0) - u_*|_{E_{\bar{\mu}}} < \delta_1 \text{ for each } x_0 \in B_{E_\mu}(u_*, \delta).$$

As  $u(\tau, x_0) \in E_{\bar{\mu}}$  and  $u(t, u(\tau, x_0)) = u(t + \tau, x_0)$  for  $t \geq \tau$ , the convergence assertion of the Theorem follows from the first part of the proof.  $\square$

*Remark 3.3.* Theorem 3.2 yields convergence of  $u(\cdot, x_0)$  in the stronger norm of  $E_{\bar{\mu}}$  for initial values in  $E_\mu$ . We note that this holds true for any  $\bar{\mu} \in [\beta, 1)$ , with  $\delta$  depending on  $\bar{\mu}$ .



## 4. APPLICATIONS TO SURFACE DIFFUSION FLOW

In this section, we apply the theory from the previous sections to extend results regarding the surface diffusion flow in various settings. First, we extend [12, Proposition 3.2] regarding well-posedness of the surface diffusion flow in the setting of so-called *axially-definable* surfaces. We then prove nonlinear stability of cylinders with radius  $r > 1$  (as equilibria of surface diffusion flow) under a general class of periodic perturbations which only require control of first-order derivatives; this result extends [12, Theorem 4.3] and [11, Theorem 4.9] where control of second-order derivatives was also required. At the conclusion of the section, we establish general well-posedness for surface diffusion flow acting on surfaces parameterized over a compact reference manifold  $\Sigma \subset \mathbb{R}^n$ , and conclude normal stability of Euclidean spheres under  $bc^{1+\alpha}$  perturbations.

**4.1. Axially-Definable Setting: Well-Posedness.** We begin with a brief introduction to the axially-definable setting and formulation of the problem; for a more detailed account we direct the reader to [12, Sections 2 and 3].

First, given  $r > 0$ , let

$$\mathcal{C}_r := \{(x, r \cos(\theta), r \sin(\theta)) : x \in \mathbb{R}, \theta \in \mathbb{T}\}$$

denote the unbounded cylinder in  $\mathbb{R}^3$  of radius  $r$ , where  $\mathbb{T} := [0, 2\pi]$  denotes the one-dimensional torus, with 0 and  $2\pi$  identified. Next, we fix a parameter  $\alpha \in (0, 1)$  and define the Banach spaces

$$E_0 := bc^\alpha(\mathcal{C}_r) \quad \text{and} \quad E_1 := bc^{4+\alpha}(\mathcal{C}_r), \quad (4.1)$$

where  $bc^{k+\alpha}$ ,  $k \in \mathbb{N}$ , denotes the family of  $k$ -times differentiable *little-Hölder* regular functions. In particular, on an open set  $U \subset \mathbb{R}^n$ ,  $bc^\alpha(U)$  is defined as the closure of the bounded smooth functions  $BC^\infty(U)$  in the topology of  $BC^\alpha(U)$ , the Banach space of all bounded Hölder-continuous functions of exponent  $\alpha$ . Then  $bc^{k+\alpha}(U)$  consists of functions having continuous and bounded derivatives of order  $k$ , whose  $k^{\text{th}}$ -order derivatives are in  $bc^\alpha(U)$ . The space  $bc^{k+\alpha}(\mathcal{C}_r)$  is defined via an atlas of local charts.

Taking  $\mu = 1/4$  and  $\beta = 3/4$ , we define the continuous interpolation spaces  $E_\mu := (E_0, E_1)_{\mu, \infty}^0$  and  $E_\beta := (E_0, E_1)_{\beta, \infty}^0$ . It is well-known that the scale of little-Hölder spaces is closed under continuous interpolation (c.f. [13] and [22]) and so these spaces are likewise identified as

$$E_\mu = bc^{1+\alpha}(\mathcal{C}_r) \quad \text{and} \quad E_\beta = bc^{3+\alpha}(\mathcal{C}_r).$$

With the spaces  $E_0, E_\mu, E_\beta, E_1$  thus set, note that condition (1.3) becomes

$$\frac{\rho_j}{2} + \beta_j \leq 1,$$

so that we have a critical index  $j$  exactly when  $\rho_j/2 + \beta_j = 1$ . Further, with  $\varepsilon > 0$  fixed, we define the family of admissible initial values (which coincides with surfaces that remain bounded away from the central axis of rotation)

$$V_\mu := E_\mu \cap \{h : \mathcal{C}_r \rightarrow \mathbb{R} \mid h(p) > \varepsilon - r \text{ for all } p \in \mathcal{C}_r\}.$$

We say that a surface  $\Gamma \subset \mathbb{R}^3$  is *axially-definable* if it can be parameterized as

$$\Gamma = \Gamma(h) = \{p + h(p)\nu(p) : p \in \mathcal{C}_r\}$$

for some *height function*  $h : \mathcal{C}_r \rightarrow \mathbb{R}$  satisfying  $h > -r$  on  $\mathcal{C}_r$ , where  $\nu$  denotes the outer unit normal field over  $\mathcal{C}_r$ . In the setting of axially-definable surfaces, the surface diffusion flow is expressed as the following evolution equation for time-dependent height functions  $h = h(t, p) = h(t, x, \theta)$ :

$$\begin{cases} h_t(t, p) = [G(h(t))](p), & \text{for } t > 0, p \in \mathcal{C}_r \\ h(0) = h_0, & \text{on } \mathcal{C}_r. \end{cases} \quad (4.2)$$

As shown in [12, Section 2.2], the evolution operator  $G$  takes the form

$$\begin{aligned} G(h) := \frac{1}{(r+h)} & \left\{ \partial_x \left[ \frac{(r+h)^2 + h_\theta^2}{\sqrt{\mathcal{G}}} \partial_x \mathcal{H}(h) - \frac{h_x h_\theta}{\sqrt{\mathcal{G}}} \partial_\theta \mathcal{H}(h) \right] \right. \\ & \left. + \partial_\theta \left[ \frac{1 + h_x^2}{\sqrt{\mathcal{G}}} \partial_\theta \mathcal{H}(h) - \frac{\rho_x h_\theta}{\sqrt{\mathcal{G}}} \partial_x \mathcal{H}(h) \right] \right\}, \end{aligned} \quad (4.3)$$

where  $\mathcal{H}(h)$  denotes the mean curvature of the surface  $\Gamma(h)$  and  $\mathcal{G} = \mathcal{G}(h)$  is the determinant of the first fundamental form  $[g_{ij}] = [g_{ij}(h)]$  on  $\Gamma(h)$ .

Using [12, Equations (2.2)–(2.3)], one can expand (4.3) to see that  $G(h)$  is a fourth-order quasilinear operator of the form

$$\begin{aligned} G(h) &= -A(h)h + F_1(h) + F_2(h) \\ &:= - \left( \sum_{|\eta|=4} b_\eta(h, \partial^1 h) \partial^\eta h \right) + F_1(h, \partial^1 h) + F_2(h, \partial^1 h, \partial^2 h, \partial^3 h), \end{aligned}$$

where  $\eta = (\eta_1, \eta_2) \in \mathbb{N}^2$  is a multi-index,  $|\eta| := \eta_1 + \eta_2$  its length,  $\partial^\eta := \partial_x^{\eta_1} \partial_\theta^{\eta_2}$  the mixed partial derivative operator, and  $\partial^k h$  denotes the vector of all derivatives  $\partial^\eta h$  for  $|\eta| = k$ . We note that  $A(h)$  here contains only the highest-order terms of the operator  $\mathcal{A}(h)$  expressed in [12, Section 3.2] — which is essential in the current setting to ensure the coefficients  $b_\eta(h, \partial^1 h)$  are well-defined for  $h \in E_\mu = bc^{1+\alpha}(\mathcal{C}_r)$ . It follows that the principal symbols  $\sigma[A(h)]$  and  $\sigma[\mathcal{A}(h)]$  coincide, so by [12, Equation (3.3)] we have

$$\sigma[A(h)](p, \xi) \geq \frac{1}{\mathcal{G}^2} \left( (r+h)^2 \xi_1^2 + \xi_2^2 \right)^2 \quad \text{for } (p, \xi) \in \mathcal{C}_r \times \mathbb{R}^2. \quad (4.4)$$

This last result implies uniform ellipticity of  $A(h)$  on  $\mathcal{C}_r$  and thus with [12, Proposition 3.1] we have

$$\begin{aligned} (A, F_1) &\in C^\omega(V_\mu, \mathcal{M}_\mu(E_1, E_0) \times E_0) \\ F_2 &\in C^\omega(V_\mu \cap E_\beta, E_0). \end{aligned} \quad (4.5)$$

We have now confirmed that the mappings  $A, F_1$  and  $F_2$  satisfy properties **(H1)** and (3.9). Regarding confirmation of the structural conditions **(H2)**, we expand terms of (4.3) to confirm

$$\begin{aligned} F_2(h) &= \sum_{\substack{|\eta|=3 \\ |\tau| \leq 2}} c_{\eta, \tau}(h) \partial^\tau h \partial^\eta h \\ &\quad + \sum_{\max\{|\eta|, |\tau|, |\sigma|\}=2} d_{\eta, \tau, \sigma}(h) \partial^\eta h \partial^\tau h \partial^\sigma h, \end{aligned} \quad (4.6)$$

where the functions  $c_{\eta, \tau}, d_{\eta, \tau, \sigma}$  depend only upon  $h$  and  $\partial^1 h$ , and are analytic by (4.5). Of particular importance in (4.6), we note that third-order derivatives of  $h$

appear linearly in terms with at most linear factors of  $\partial^2 h$ , while lower-order terms include at most cubic factors of  $\partial^2 h$ .

Letting  $h_0 \in V_\mu$  and  $R > 0$ , we choose  $h_1, h_2 \in \bar{B}_{E_\mu}(h_0, R) \cap (V_\mu \cap E_\beta)$  and use (4.6) to bound  $|F_2(h_1) - F_2(h_2)|_{E_0}$ . Throughout the following computations, we use  $\tilde{C}$  to denote a generic constant that depends only upon  $R$  and  $|h_0|_{E_\mu} = |h_0|_{bc^{1+\alpha}}$ .

Considering the first term in (4.6), when  $|\tau| \leq 1$  we incorporate  $\partial^\tau h_i$  into  $c_{\eta,\tau}(h_i)$  to derive bounds depending only on zeroth and first-order derivatives,  $i = 1, 2$ . Hence, when  $|\tau| \leq 1$  we have

$$|c_{\eta,\tau}(h_1) \partial^\eta h_1 \partial^\tau h_1 - c_{\eta,\tau}(h_2) \partial^\eta h_2 \partial^\tau h_2|_{E_0} \leq \tilde{C} |h_1 - h_2|_{bc^{3+\alpha}},$$

which is a term as in (1.4) corresponding to  $(\rho_j, \beta_j) = (0, 3/4)$  (which is a subcritical index since  $\rho_j/2 + \beta_j < 1$ ). Meanwhile, when  $|\tau| = 2$ , we rewrite

$$\begin{aligned} c_{\eta,\tau}(h_1) \partial^\eta h_1 \partial^\tau h_1 - c_{\eta,\tau}(h_2) \partial^\eta h_2 \partial^\tau h_2 = \\ c_{\eta,\tau}(h_1) \left( \partial^\eta h_1 (\partial^\tau h_1 - \partial^\tau h_2) + \partial^\tau h_2 (\partial^\eta h_1 - \partial^\eta h_2) \right) \\ + \partial^\eta h_2 \partial^\tau h_2 (c_{\eta,\tau}(h_1) - c_{\eta,\tau}(h_2)) \end{aligned} \quad (4.7)$$

and note that  $c_{\eta,\tau}$  is locally Lipschitz continuous in  $E_\mu$  in order to bound the previous expression in  $E_0$  by

$$\tilde{C} \left( |h_1|_{E_\beta} |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{bc^{2+\alpha}} |h_1 - h_2|_{bc^{3+\alpha}} + |h_2|_{bc^{2+\alpha}} |h_2|_{E_\beta} |h_1 - h_2|_{bc^{1+\alpha}} \right).$$

Further, by the reiteration theorem for continuous interpolation (c.f. [1, Section I.2.8]), we have  $bc^{2+\alpha} = (E_\mu, E_\beta)_{1/2, \infty}^0$ . Thus,

$$|h_i|_{bc^{2+\alpha}} \leq \tilde{C} |h_i|_{bc^{3+\alpha}}^{1/2} = \tilde{C} |h_i|_{E_\beta}^{1/2} \quad (4.8)$$

and so, when  $|\tau| = 2$ , we bound

$$\begin{aligned} |c_{\eta,\tau}(h_1) \partial^\eta h_1 \partial^\tau h_1 - c_{\eta,\tau}(h_2) \partial^\eta h_2 \partial^\tau h_2|_{E_0} \leq \tilde{C} \left( |h_1|_{E_\beta} |h_1 - h_2|_{bc^{2+\alpha}} \right. \\ \left. + |h_2|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{3+\alpha}} + |h_2|_{E_\beta}^{3/2} |h_1 - h_2|_{bc^{1+\alpha}} \right), \end{aligned}$$

which are three critical terms with  $(\rho_j, \beta_j) = (1, 1/2)$ ,  $(\rho_j, \beta_j) = (1/2, 3/4)$ , and  $(\rho_j, \beta_j) = (3/2, 1/4)$ , respectively.

Moving on to the second term in (4.6), we first rewrite the difference

$$\begin{aligned} d_{\eta,\tau,\sigma}(h_1) \partial^\eta h_1 \partial^\tau h_1 \partial^\sigma h_1 - d_{\eta,\tau,\sigma}(h_2) \partial^\eta h_2 \partial^\tau h_2 \partial^\sigma h_2 \\ = d_{\eta,\tau,\sigma}(h_1) \left( \partial^\eta h_1 \partial^\tau h_1 (\partial^\sigma h_1 - \partial^\sigma h_2) \right. \\ \left. + \partial^\eta h_1 \partial^\sigma h_2 (\partial^\tau h_1 - \partial^\tau h_2) + \partial^\tau h_2 \partial^\sigma h_2 (\partial^\eta h_1 - \partial^\eta h_2) \right) \\ \left. + \partial^\eta h_2 \partial^\tau h_2 \partial^\sigma h_2 (d_{\eta,\tau,\sigma}(h_1) - d_{\eta,\tau,\sigma}(h_2)) \right) \end{aligned} \quad (4.9)$$

and then we proceed by splitting our analysis into three cases.

First, consider the case when only one of the terms  $|\eta|$ ,  $|\tau|$ , or  $|\sigma|$  is equal to two. In this case, we again employ local Lipschitz continuity of the functions  $d_{\eta,\tau,\sigma}$  and properties of interpolation to bound (4.9) in  $E_0$  by

$$\tilde{C} \left( |h_1|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{1+\alpha}} + |h_2|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{1+\alpha}} + |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{1+\alpha}} \right),$$

which results in two subcritical terms (after combining expressions) with  $(\rho_j, \beta_j) = (1/2, 1/4)$  and  $(\rho_j, \beta_j) = (0, 1/2)$ .

Next, we consider the case when exactly two of the terms  $|\eta|$ ,  $|\tau|$ , and  $|\sigma|$ , equal two. Without loss of generality, we take  $|\sigma| = 1$ , since we can arrive at similar expressions to (4.9) with any combination of multi-indices contained in the cross-term  $\partial^\eta h_1 \partial^\sigma h_2$ . Similar to the previous case, we majorize (4.9) in  $E_0$ , using (4.8),

$$\begin{aligned} & \tilde{C} \left( |h_1|_{bc^{2+\alpha}}^2 |h_1 - h_2|_{bc^{1+\alpha}} + |h_1|_{bc^{2+\alpha}} |h_1 - h_2|_{bc^{2+\alpha}} \right. \\ & \quad \left. + |h_2|_{bc^{2+\alpha}} |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{bc^{2+\alpha}}^2 |h_1 - h_2|_{bc^{1+\alpha}} \right) \\ & \leq \tilde{C} \left( |h_1|_{E_\beta} |h_1 - h_2|_{bc^{1+\alpha}} + |h_1|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{2+\alpha}} \right. \\ & \quad \left. + |h_2|_{E_\beta}^{1/2} |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{E_\beta} |h_1 - h_2|_{bc^{1+\alpha}} \right), \end{aligned}$$

which contributes two additional subcritical terms with  $(\rho_j, \beta_j) = (1, 1/4)$  and  $(\rho_j, \beta_j) = (1/2, 1/2)$ .

Finally, in case  $|\eta| = |\tau| = |\sigma| = 2$ , we bound (4.9) by, again employing (4.8),

$$\begin{aligned} & \tilde{C} \left( (|h_1|_{E_\beta} + |h_1|_{E_\beta}^{1/2} |h_2|_{E_\beta}^{1/2} + |h_2|_{E_\beta}) |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{E_\beta}^{3/2} |h_1 - h_2|_{bc^{1+\alpha}} \right) \\ & \leq \tilde{C} \left( |h_1|_{E_\beta} |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{E_\beta} |h_1 - h_2|_{bc^{2+\alpha}} + |h_2|_{E_\beta}^{3/2} |h_1 - h_2|_{bc^{1+\alpha}} \right), \end{aligned}$$

where we have also applied Young's inequality to get

$$|h_1|_{E_\beta}^{1/2} |h_2|_{E_\beta}^{1/2} \leq \frac{1}{2} |h_1|_{E_\beta} + \frac{1}{2} |h_2|_{E_\beta}.$$

We thus produce two critical terms in this case, with  $(\rho_j, \beta_j) = (1, 1/2)$  and  $(\rho_j, \beta_j) = (3/2, 1/4)$ .

Together with the analysis for the first term of (4.6), we have thus demonstrated that  $F_2$  satisfies the structural condition **(H2)** and we can now apply Theorem 2.2 to produce the following results. Moreover, we note that  $\mu = 1/4$  is the critical weight for  $F_2$ , which indicates that  $bc^{1+\alpha}(\mathcal{C}_r)$  is the critical space for (4.2).

**Theorem 4.1 (Well-Posedness of (4.2)).** *Fix  $\varepsilon > 0$  and  $\alpha \in (0, 1)$ .*

(a) *For each initial value*

$$h_0 \in V_\mu := bc^{1+\alpha}(\mathcal{C}_r) \cap [h > \varepsilon - r],$$

*there exists a unique maximal solution to (4.2) with the addition property*

$$h(\cdot, h_0) \in C([0, t^+(h_0)), bc^{1+\alpha}(\mathcal{C}_r)) \cap C((0, t^+(h_0)), bc^{4+\alpha}(\mathcal{C}_r)).$$

*Further, it follows that the map  $[(t, h_0) \mapsto h(t, h_0)]$  defines a semiflow on  $V_\mu$  which is analytic for  $t > 0$  and Lipschitz continuous for  $t \geq 0$ .*

(b) *Moreover, if the solution  $h(\cdot, h_0)$  satisfies:*

- (i)  $h(\cdot, h_0) \in UC(J(h_0), bc^{1+\alpha}(\mathcal{C}_r))$ , and
  - (ii) *there exists  $M > 0$  so that, for all  $t \in J(h_0) := [0, t^+(h_0))$ ,*
    - (ii.a)  $h(t, h_0)(p) \geq 1/M - r$  for all  $p \in \mathcal{C}_r$ , and
    - (ii.b)  $|h(t, h_0)|_{bc^{1+\alpha}(\mathcal{C}_r)} \leq M$ ,
- then it holds that  $t^+(h_0) = \infty$  and  $h(\cdot, h_0)$  is a global solution of (4.2).*

*Remark 4.2.* (a) Of particular note in Theorem 4.1, control of first-order derivatives of  $h$  is sufficient to determine the lifespan of maximal solutions. This improves previous results in the axially-definable setting (c.f. [12, Proposition 3.2]) which required control of second-order derivatives of solutions as well.

(b) Regarding analyticity of the semiflow  $[(t, h_0) \mapsto h(t, h_0)]$  for  $t > 0$ : for any  $\tau > 0$  we note that

$$h(\tau, h_0) \in V_\beta := bc^{3+\alpha}(\mathcal{C}_r) \cap [h > \varepsilon - r].$$

Thus, analyticity holds for  $t > \tau$  in  $V_\beta$  by [5, Theorem 6.1] and (4.5), and then analyticity also holds in  $V_\mu$  by embedding.

(c) In the setting of surfaces expressed as graphs over  $\mathbb{R}^n$ , existence and uniqueness of solutions with initial data in  $bc^{1+\alpha}(\mathbb{R}^n)$  was established in [3, Theorem 4.2]. However, we note that the author requires initial values to be slightly more regular than those considered here, due to the fact that he tracks regularity of solutions in a different topology than that of the space where he takes initial data.

**4.2. Axially-Definable Setting: Stability of Cylinders.** Considering the stability of cylinders as equilibria for (4.2), we first introduce the  $2\pi$ -periodic little-Hölder spaces  $bc_{per}^{k+\alpha}$ ,  $k \in \mathbb{N}$ , defined as the subspace of functions  $h \in bc^{k+\alpha}(\mathcal{C}_r)$  exhibiting  $2\pi$ -periodicity along the  $x$ -axis; i.e.

$$h(x \pm 2\pi, \theta) = h(x, \theta) \quad \text{for all } (x, \theta) \in \mathcal{C}_r.$$

As shown in [12, Sections 3.4–4.1], working in this setting allows access to Fourier series representations for height functions  $h$  and guarantees the linearized operator  $DG(h)$  has a discrete spectrum.

Regarding well-posedness in the periodic setting, it was shown in [12, Proposition 3.4] that  $G$  preserves periodicity, so Theorem 4.1(a) continues to hold verbatim with  $bc_{per}$  replacing  $bc$  throughout. Meanwhile, we note that global solutions in the periodic setting differ slightly from Theorem 4.1(b) owing to the compactness of the embedding  $bc_{per}^{4+\alpha}(\mathcal{C}_r) \hookrightarrow bc_{per}^\alpha(\mathcal{C}_r)$  (c.f. Theorem 2.2(c)).

Noting that  $h_* \equiv 0$  is always an equilibrium of (4.2) (which coincides with the observation that the cylinder  $\mathcal{C}_r$  is an equilibrium of surface diffusion flow), we consider the stability of  $h_*$  under perturbations in

$$V_{\mu, per} := bc_{per}^{1+\alpha}(\mathcal{C}_r) \cap [h > -r].$$

Further, we denote by  $\mathcal{M}_{cyl}$  the family of height functions  $\bar{h}$  such that  $\Gamma(\bar{h})$  defines a cylinder  $C(\bar{y}, \bar{z}, \bar{r})$  — symmetric about axis  $(\cdot, \bar{y}, \bar{z})$  in  $\mathbb{R}^3$  with radius  $\bar{r} > 0$  — in a neighborhood of  $\mathcal{C}_r$ . With these preparations, we state the following stability result.

**Theorem 4.3** (Global existence, stability / instability of cylinders). *Fix  $\alpha \in (0, 1)$ .*

(a) **(Global Existence)** *Let  $h_0 \in V_{\mu, per}$  and suppose there exists a constant  $M > 0$  so that*

- (i)  $h(t, h_0)(p) \geq 1/M - r$  for all  $t \in J(h_0) = [0, t^+(h_0))$ ,  $p \in \mathcal{C}_r$ , and
- (ii)  $|h(t, h_0)|_{bc^{1+\delta}} \leq M$  for all  $t \in [\tau, t^+(h_0))$ , for some  $\tau \in J(h_0)$  and  $\delta \in (\alpha, 1)$ ,

*then  $t^+(h_0) = \infty$ , so that  $h(\cdot, h_0)$  is a global solution.*

- (b) **(Stability)** Fix  $r > 1$  and  $\bar{\mu} \in (0, 1)$ . There exists a positive constant  $\delta > 0$  such that, given any admissible periodic perturbation

$$h_0 \in V_{\mu, \text{per}} := bc_{\text{per}}^{1+\alpha}(\mathcal{C}_r) \cap [h > -r]$$

with  $|h_0|_{bc^{1+\alpha}(\mathcal{C}_r)} < \delta$ , the solution  $h(\cdot, h_0)$  exists globally in time and converges to some  $\bar{h} \in \mathcal{M}_{\text{cyl}}$  at an exponential rate, in the topology of  $E_{\bar{\mu}}$ .

- (c) **(Instability)** For  $0 < r < 1$  the function  $h_* \equiv 0$  is unstable in the topology of  $bc_{\text{per}}^{1+\alpha}(\mathcal{C}_r)$ .

*Proof.* (a) This result follows from Theorem 2.2(c), noting that  $bc_{\text{per}}^{4+\alpha}(\mathcal{C}_r) \hookrightarrow bc_{\text{per}}^{\alpha}(\mathcal{C}_r)$  is a compact embedding in this periodic setting. Conditions (i)–(ii) guarantee the solution remains bounded away from the boundary  $\partial V_{\mu, \text{per}}$ .

(b) First note that restricting the domains of  $(A, F_1, F_2)$  to periodic little–Hölder spaces will maintain the conditions **(H1)**–**(H2)** and (4.5), all confirmed in Section 4.1. From the proof of [12, Theorem 4.3] we know that  $h_*$  is normally stable when  $r > 1$ . The conclusion thus follows from Theorem 3.2.

(c) It follows from [12, Theorem 4.3(b)] that  $h_*$  is unstable in the topology of  $bc_{\text{per}}^{3+\alpha}(\mathcal{C}_r)$ . Instability in  $bc_{\text{per}}^{1+\alpha}(\mathcal{C}_r)$  then follows from Proposition 3.1.  $\square$

**4.3. Axisymmetric Setting.** We turn now to consider (4.2) acting on the scale of axisymmetric little–Hölder spaces  $bc_{\text{sym}}^{k+\alpha}(\mathcal{C}_r)$ ,  $k \in \mathbb{N}$ , defined as the subspace of functions  $h \in bc^{k+\alpha}(\mathcal{C}_r)$  exhibiting symmetry around the  $x$ -axis; i.e.

$$h(x, \theta_1) = h(x, \theta_2) \quad \text{for all } x \in \mathbb{R} \text{ and } \theta_1, \theta_2 \in \mathbb{T}.$$

These functions naturally coincide with surfaces  $\Gamma(h)$  which are symmetric about the central  $x$ -axis, as considered in [11]; although we relax the setting slightly by not enforcing axial–periodicity for our well–posedness result.

For all such functions with sufficient regularity, it follows that  $\partial_{\theta} h \equiv 0$  and the application of the evolution operator  $G$  to  $h \in bc_{\text{sym}}^{4+\alpha}(\mathcal{C}_r)$  produces the simplified expression

$$G(h) = \frac{1}{(r+h)} \partial_x \left[ \frac{(r+h)}{\sqrt{1+h_x^2}} \partial_x \left( \frac{1}{(r+h)\sqrt{1+h_x^2}} - \frac{h_{xx}}{(1+h_x^2)^{\frac{3}{2}}} \right) \right]. \quad (4.10)$$

In fact, a complete expansion of individual terms for the operator  $G$  is provided in [11, Equations (2.1)–(2.3)] from which we deduce

$$A(h) = \frac{1}{(1+h_x^2)^2} \partial_x^4 \quad F_1(h) = \frac{h_x^2}{(r+h)^3(1+h_x^2)}$$

and

$$\begin{aligned} F_2(h) = & \frac{2h_x(1+h_x^2)}{(r+h)(1+h_x^2)^3} h_{xxx} + \frac{-6(r+h)h_x}{(r+h)(1+h_x^2)^3} h_{xx}h_{xxx} \\ & + \frac{h_x^2-1}{(r+h)^2(1+h_x^2)^2} h_{xx} + \frac{6h_x^2-1}{(r+h)(1+h_x^2)^3} h_{xx}^2 + \frac{3-15h_x^2}{(1+h_x^2)^4} h_{xx}^3, \end{aligned}$$

where we can explicitly observe the structure of (4.6).

To apply the results of Section 4.1 to the axisymmetric setting, it suffices to note that the property of axisymmetry is preserved by (4.2). This claim is clear from a purely geometric perspective, since the evolution equation (4.2) is completely

determined by the geometry of the surfaces  $\Gamma(h(t))$ , and axisymmetry of the surface imparts the same symmetry onto the geometric structure. However, one can also confirm preservation of axisymmetry analytically by confirming that  $G$  commutes with the azimuthal shift operators  $T_\phi$ , for  $\phi \in \mathbb{T}$ ; defined by

$$T_\phi(h(x, \theta)) := h(x, \theta + \phi) \quad \text{for } (x, \theta) \in \mathcal{C}_r.$$

Indeed, we note that axisymmetry of  $h$  can be characterized by the property that  $T_\phi h = h$  on  $\mathcal{C}_r$  for all  $\phi \in \mathbb{T}$ . Then, by direct computation one confirms that  $T_\phi G(h) = G(T_\phi h)$ . From here we apply Theorems 4.1 and 4.3, along with an argument similar to the proof of [12, Proposition 3.6] to produce the following extensions of [11, Propositions 2.2 and 2.3].

**Theorem 4.4** (Well-Posedness). *Fix  $\varepsilon > 0$  and  $\alpha \in (0, 1)$ .*

(a) *For each admissible axisymmetric initial value*

$$h_0 \in V_{\mu, \text{sym}} := bc_{\text{sym}}^{1+\alpha}(\mathcal{C}_r) \cap [h > \varepsilon - r],$$

*there exists a unique maximal solution to (4.2) with the additional property*

$$h(\cdot, h_0) \in C([0, t^+(h_0)), bc_{\text{sym}}^{1+\alpha}(\mathcal{C}_r)) \cap C((0, t^+(h_0)), bc_{\text{sym}}^{4+\alpha}(\mathcal{C}_r)).$$

*Further, the map  $[(t, h_0) \mapsto h(t, h_0)]$  defines a semiflow on  $V_{\mu, \text{sym}}$  which is analytic for  $t > 0$  and Lipschitz continuous for  $t \geq 0$ .*

(b) *Moreover, if the solution  $h(\cdot, h_0)$  satisfies:*

- (i)  $h(\cdot, h_0) \in UC(J(h_0), bc_{\text{sym}}^{1+\alpha}(\mathcal{C}_r))$ , and
- (ii) *there exists  $M > 0$  so that, for all  $t \in J(h_0) := [0, t^+(h_0))$ ,*
  - (ii.a)  $h(t, h_0)(p) \geq 1/M - r$  for all  $p \in \mathcal{C}_r$ , and
  - (ii.b)  $|h(t, h_0)|_{bc^{1+\alpha}(\mathcal{C}_r)} \leq M$ ,

*then it holds that  $t^+(h_0) = \infty$  and  $h(\cdot, h_0)$  is a global solution of (4.2).*

With the additional assumption of periodicity along the  $x$ -axis, we likewise define  $bc_{\text{sym}, \text{per}}^{k+\alpha}(\mathcal{C}_r)$ ,  $k \in \mathbb{N}$ , as the subspace of periodic functions  $h \in bc_{\text{sym}}^{k+\alpha}(\mathcal{C}_r)$ , with  $h(x \pm 2\pi, \theta) = h(x, \theta)$  for all  $(x, \theta) \in \mathcal{C}_r$ .

**Theorem 4.5** (Global existence, stability and instability of cylinders). *Fix  $\alpha \in (0, 1)$ .*

(a) **(Global Existence)** *Let  $h_0 \in V_{\mu, \text{sym}, \text{per}}$  and suppose there exists a constant  $M > 0$  so that, for all  $t \in J(h_0)$ ,*

- (i)  $h(t, h_0)(p) \geq 1/M - r$  for all  $t \in J(h_0) = [0, t^+(h_0))$ ,  $p \in \mathcal{C}_r$ , and
- (ii)  $|h(t, h_0)|_{bc^{1+\delta}} \leq M$  for all  $t \in [\tau, t^+(h_0))$ , for some  $\tau \in J(h_0)$  and  $\delta \in (\alpha, 1)$ ,

*then  $t^+(h_0) = \infty$ , so that  $h(\cdot, h_0)$  is a global solution.*

(b) **(Stability)** *Fix  $r > 1$  and  $\bar{\mu} \in (0, 1)$ . There exists a positive constant  $\delta > 0$  such that, given any admissible periodic axisymmetric perturbation*

$$h_0 \in V_{\mu, \text{sym}, \text{per}} := bc_{\text{sym}, \text{per}}^{1+\alpha}(\mathcal{C}_r) \cap [h > -r]$$

*with  $|h_0|_{bc^{1+\alpha}(\mathcal{C}_r)} < \delta$ , the solution  $h(\cdot, h_0)$  exists globally in time and converges to some  $\bar{h} \in \mathcal{M}_{\text{cyl}}$  at an exponential rate, in the topology of  $E_{\bar{\mu}}$ .*

(c) **(Instability)** *For  $0 < r < 1$  the function  $h_* \equiv 0$  is unstable in the topology of  $bc_{\text{per}}^{1+\alpha}(\mathcal{C}_r)$ .*

**4.4. Surfaces Near Compact Hypersurfaces.** We conclude the paper by looking at the flow of surfaces parameterized over a fixed reference manifold and extend results in [6]. In particular, let  $\Sigma$  denote a smooth, closed, compact, immersed, oriented hypersurface in  $\mathbb{R}^n$ , and let  $\nu_\Sigma$  be a unit normal vector field on  $\Sigma$ , compatible with the chosen orientation. It follows that there exists a constant  $a > 0$  and an open atlas  $\{U_\ell : \ell \in \mathcal{L}\}$  for  $\Sigma$  so that

$$X_\ell : U_\ell \times (-a, a) \rightarrow \mathbb{R}^n, \quad X_\ell(p, r) := p + r\nu_\Sigma(p), \quad (4.11)$$

is a smooth diffeomorphism onto the range  $R_\ell := \text{im}(X_\ell)$ , for  $\ell \in \mathcal{L}$ . We capture the evolution of surfaces that are  $C^1$ -close to  $\Sigma$  via time-dependent height functions  $h : \mathbb{R}^+ \times \Sigma \rightarrow (-a, a)$ . In particular, to  $h(t) := h(t, \cdot)$  we associate the surface

$$\Gamma(h(t)) := \bigcup_{\ell \in \mathcal{L}} \{X_\ell(p, h(t, p)) : p \in U_\ell\},$$

which is parametrized by the mapping

$$\Psi_{h(t)} : \Sigma \rightarrow \mathbb{R}^n, \quad \Psi_{h(t)}(p) := p + h(t, p)\nu_\Sigma(p). \quad (4.12)$$

As in the previous settings, we let  $\alpha \in (0, 1)$  and work in spaces of little-Hölder continuous functions

$$E_0 := bc^\alpha(\Sigma), \quad E_\mu = bc^{1+\alpha}(\Sigma), \quad E_\beta = bc^{3+\alpha}(\Sigma), \quad E_1 := bc^{4+\alpha}(\Sigma),$$

with  $\mu = 1/4$  and  $\beta = 3/4$ . Further, we define

$$V_\mu := bc^{1+\alpha}(\Sigma) \cap [h]_{C(\Sigma)} < a].$$

To express the equations for surface diffusion flow of  $\Gamma(h(t))$  as an evolution equation acting on the height functions  $h$ , we direct the reader to [6, Section 2] and [21, Section 5] where details are given for pulling back the governing equation  $V_\Gamma = \Delta_\Gamma \mathcal{H}_\Gamma$ , defined on  $\Gamma(h(t))$ , to an equivalent equation on the reference manifold  $\Sigma$ . We thus arrive at an expression

$$\begin{cases} h_t(t, p) = [G(h(t))](p) & \text{for } t > 0, p \in \Sigma, \\ h(0) = h_0 & \text{on } \Sigma, \end{cases} \quad (4.13)$$

where the evolution operator  $G$  takes the form (c.f. [21, Section 5])

$$G(h) := -\frac{1}{\beta(h, \partial^1 h)} \Delta_h \mathcal{H}_h. \quad (4.14)$$

Utilizing expressions given in [15, Sections 3.2–3.5] or [16, Section 2.2], we expand (4.14) and confirm properties **(H1)**, **(H2)** and (3.9). For the structure of the Laplace–Beltrami operator in local coordinates, we have (employing the standard summation convention over repeated instances of  $i, j, k$  taking values from 1 to  $(n-1)$  — the dimension of the manifold)

$$\Delta_h \varphi = a^{ij}(h, \partial^1 h) \partial_i \partial_j \varphi_* + b^k(h, \partial^1 h, \partial^2 h) \partial_k \varphi_*, \quad (4.15)$$

where  $\varphi$  is a scalar function on  $\Gamma(h)$  and  $\varphi_* := \Psi_h^* \varphi$  its pull-back to  $\Sigma$  through the parameterization  $\Psi_h$ . The coefficient functions  $a^{ij}$  and  $b^k$  are expressed as

$$\begin{aligned} a^{ij}(h, \partial^1 h) &= \left( P_{\Gamma(h)} M_0(h) \tau_\Sigma^i \middle| P_{\Gamma(h)} M_0(h) \tau_\Sigma^j \right) \quad \text{and} \\ b^k(h, \partial^1 h, \partial^2 h) &= \left( \partial_i (M_0(h) P_{\Gamma(h)}) P_{\Gamma(h)} M_0(h) \tau_\Sigma^i \middle| \tau_\Sigma^k \right), \end{aligned}$$



where  $\tau_i^\Sigma|_p$  denote elements of a basis for the tangent space  $T_p\Sigma$  to a point  $p \in \Sigma$ , while  $\tau_\Sigma^i|_p$  make up a corresponding basis for the dual of  $T_p\Sigma$ , and  $(\cdot|\cdot)$  is the inner product in Euclidean space  $\mathbb{R}^n$ . Further,  $M_0(h) := (I - hL_\Sigma)^{-1}$  depends upon  $h$  and the Weingarten tensor  $L_\Sigma$  on  $\Sigma$  (i.e. no derivatives of  $h$  appear in  $M_0(h)$ ) and

$$P_{\Gamma(h)} := I - \nu_{\Gamma(h)} \otimes \nu_{\Gamma(h)}$$

projects onto the tangent space  $\Gamma(h)$ . Here the normal vector to  $\Gamma(h)$  is

$$\nu_{\Gamma(h)} = \beta(h, \partial^1 h)(\nu_\Sigma - M_0(h)\nabla_\Sigma h),$$

and hence  $P_{\Gamma(h)}$  only depends upon first-order derivatives of  $h$ . Therefore, second-order derivatives of  $h$  only appear in  $b^k$  when the derivative  $\partial_i$  acts on  $P_{\Gamma(h)}$  in the first term of the inner product. With no further factors of  $\partial^2 h$  appearing in the expression, it follows that  $\partial^2 h$  only appears linearly in the functions  $b^k(h, \partial^1 h, \partial^2 h)$ .

By [16, Section 2.2] or [15, Section 3.5], the mean curvature function has the following structure in local coordinates,

$$\mathcal{H}_h = c^{ij}(h, \partial^1 h) \partial_i \partial_j h + d(h, \partial^1 h). \quad (4.16)$$

Thus, applying (4.15) to (4.16), one confirms that  $G$  exhibits the quasilinear structure

$$G(h) = -A(h, \partial^1 h)h + F_1(h, \partial^1 h) + F_2(h, \partial^1 h, \partial^2 h, \partial^3 h).$$

Considering condition **(H1)**, note that, in every local chart  $U_\ell$ , the principal symbol  $\sigma[A(h)]_\ell$  coincides with the expression given for the principal symbol  $\hat{\sigma}[P(h)]_\ell$  in [21, Section 5]. Thus, we have

$$A(h) \in \mathcal{M}_\mu(E_1, E_0) \quad \text{for all } h \in V_\mu,$$

and we likewise conclude

$$\begin{aligned} (A, F_1) &\in C^\omega(V_\mu, \mathcal{M}_\mu(E_1, E_0) \times E_0) \\ F_2 &\in C^\omega(V_\mu \cap E_\beta, E_0). \end{aligned} \quad (4.17)$$

Considering condition **(H2)**, by the argument in Section 4.1, it suffices to show that  $F_2$  exhibits the same structure as (4.6). Thus, applying (4.15) to (4.16) (noting that (4.16) is the pulled back expression of mean curvature), we first consider the four scenarios where third-order derivatives arise in  $G(h)$ , namely:

- $a^{ij}(h, \partial^1 h) \partial_i c^{kl}(h, \partial^1 h) \partial_j \partial_k \partial_l h$ ,
- $a^{ij}(h, \partial^1 h) \partial_i \partial_j c^{kl}(h, \partial^1 h) \partial_k \partial_l h$ ,
- $b^k(h, \partial^1 h, \partial^2 h) c^{ij}(h, \partial^1 h) \partial_k \partial_i \partial_j h$ , and
- $a^{ij}(h, \partial^1 h) \partial_i \partial_j d(h, \partial^1 h)$ .

In all such scenarios, when  $\partial^3 h$  is produced it appears linearly and it multiplies factors of  $\partial^2$  that appear at most linearly. Next, considering all cases within which second-order derivatives arise in  $G(h)$  — without accompanying factors of  $\partial^3 h$  — one likewise confirms that at most cubic factors of  $\partial^2 h$  appear. Therefore, we conclude that conditions **(H1)**, **(H2)** and (3.9) all hold for (4.13), and we thus produce the following extension of [6, Theorem 2.2] by application of Theorem 2.2. Note that the embeddings  $bc^{k+\alpha}(\Sigma) \hookrightarrow bc^\alpha(\Sigma)$  are compact here, since the domain  $\Sigma$  is itself compact.

**Theorem 4.6** (Well-Posedness). *Fix  $\alpha \in (0, 1)$  and  $a > 0$ . Let  $\Sigma$  be a smooth, closed, compact, immersed, oriented hypersurface in  $\mathbb{R}^n$  on which there exists an open atlas  $\{U_\ell : \ell \in \mathcal{L}\}$  where  $X_\ell : [(p, r) \mapsto p + r\nu_\Sigma(p)] : \Sigma \times (-a, a) \rightarrow \mathbb{R}^n$  is a smooth diffeomorphism onto  $R_\ell := \text{im}(X_\ell)$ , for  $\ell \in \mathcal{L}$ .*

(a) *For each admissible initial value*

$$h_0 \in V_\mu := bc^{1+\alpha}(\Sigma) \cap \{h \mid |h|_{C(\Sigma)} < a\},$$

*there exists a unique maximal solution to (4.13) with the additional property*

$$h(\cdot, h_0) \in C([0, t^+(h_0)), bc^{1+\alpha}(\Sigma)) \cap C((0, t^+(h_0)), bc^{4+\alpha}(\Sigma)).$$

*Further, the map  $[(t, h_0) \mapsto h(t, h_0)]$  defines a semiflow on  $V_\mu$  which is analytic for  $t > 0$  and Lipschitz continuous for  $t \geq 0$ .*

(b) *Moreover, if there exists  $M > 0$  and  $\delta \in (\alpha, 1)$  so that*

- (i)  $|h(t, h_0)(p)| \leq a - 1/M$ , for all  $p \in \Sigma$  and  $t \in J(h_0)$ , and
  - (ii)  $|h(t, h_0)|_{bc^{1+\delta}(\Sigma)} \leq M$ , for all  $t \in [\tau, t^+(h_0))$ , and some  $\tau \in (0, t^+(h_0))$ ,
- then it holds that  $t^+(h_0) = \infty$  and  $h(\cdot, h_0)$  is a global solution of (4.13).*

**Remark 4.7.** (a) We note that the global existence result in Theorem 4.6(b) is limited, as it fails to account for the possibility of updating the reference manifold  $\Sigma$  as  $\Gamma(h(t))$  is leaving the tubular neighborhood, but this result is sufficient for considerations of stability/instability when  $\Sigma$  is an equilibrium.

(b) Further regularity of the surfaces  $\Gamma(h(t))$  have been shown in certain settings. In particular, when  $\Sigma$  is additionally assumed to be a smooth embedded surface, it follows from [21, Theorem 5.2], and instantaneous regularization of solutions, that  $\Gamma(h(t))$  is also smooth for all  $t \in J(h_0)$ . Likewise, if  $\Sigma$  is real analytic and embedded, then  $\Gamma(h(t))$  is also real analytic.

**4.5. Stability of Euclidean Spheres.** In the particular case that  $\Sigma$  is a Euclidean sphere, the function  $h_* \equiv 0$  is normally stable in  $E_1$  by [6, Section 3] and we thus conclude the following extension of [6, Theorem 1.2]. Note that our result shows stability of spheres under surface diffusion flow with control on only first derivatives of perturbations.

**Theorem 4.8.** *Fix  $\alpha \in (0, 1)$ ,  $\bar{\mu} \in (0, 1)$ . Let  $\Sigma$  be a Euclidean sphere in  $\mathbb{R}^n$  and choose  $a > 0$  so that the mapping  $[(p, r) \mapsto p + r\nu_\Sigma(p)]$  is a diffeomorphism on  $\Sigma \times (-a, a)$ . There exists a constant  $\delta > 0$  such that, given any admissible perturbation  $\Gamma(h_0)$  for  $h_0 \in V_\mu$  with  $|h_0|_{bc^{1+\alpha}(\Sigma)} < \delta$ , the solution  $h(\cdot, h_0)$  exists globally in time and converges to some  $\bar{h} \in \mathcal{M}_{sph}$  at an exponential rate, in the topology of  $E_{\bar{\mu}}$ . Here  $\mathcal{M}_{sph}$  denotes the family of all spheres which are sufficiently close to  $\Sigma$  in  $\mathbb{R}^n$ .*

An immediate consequence of Theorem 4.8 is a relaxation of convexity constraints for stable perturbations of a sphere. In particular, note that every  $bc^{1+\alpha}$ -neighborhood of a sphere contains non-convex hypersurfaces. This corollary provides a different approach to the same result in [7], where the authors prove the claim by showing that non-convex perturbations of spheres exist in  $B_{p,2}^{5/2-4/p}(\Sigma)$ .

**Corollary 4.9.** *There exist non-convex hypersurfaces  $\Gamma_0$  such that the solution  $h(\cdot, h_0)$  to (4.13), with  $\Gamma(h_0) = \Gamma_0$ , exists globally in time and converges exponentially fast to a sphere.*

## REFERENCES

- [1] Herbert Amann. *Linear and quasilinear parabolic problems. Vol. I*, volume 89 of *Monographs in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1995. Abstract linear theory.
- [2] Sigurd B. Angenent. Nonlinear analytic semiflows. *Proc. Roy. Soc. Edinburgh Sect. A*, 115(1-2):91–107, 1990.
- [3] Tomoro Asai. Quasilinear parabolic equation and its applications to fourth order equations with rough initial data. *J. Math. Sci. Univ. Tokyo*, 19(4):507–532 (2013), 2012.
- [4] Andrew J. Bernoff, Andrea L. Bertozzi, and Thomas P. Witelski. Axisymmetric surface diffusion: dynamics and stability of self-similar pinchoff. *J. Statist. Phys.*, 93(3-4):725–776, 1998.
- [5] Philippe Clément and Gieri Simonett. Maximal regularity in continuous interpolation spaces and quasilinear parabolic equations. *J. Evol. Equ.*, 1(1):39–67, 2001.
- [6] Joachim Escher, Uwe F. Mayer, and Gieri Simonett. The surface diffusion flow for immersed hypersurfaces. *SIAM J. Math. Anal.*, 29(6):1419–1433, 1998.
- [7] Joachim Escher and Piotr B. Mucha. The surface diffusion flow on rough phase spaces. *Discrete Contin. Dyn. Syst.*, 26(2):431–453, 2010.
- [8] Herbert Koch and Tobias Lamm. Geometric flows with rough initial data. *Asian J. Math.*, 16(2):209–235, 2012.
- [9] Jeremy LeCrone, Jan Prüss, and Mathias Wilke. On quasilinear parabolic evolution equations in weighted  $L_p$ -spaces II. *J. Evol. Equ.*, 14(3):509–533, 2014.
- [10] Jeremy LeCrone and Gieri Simonett. Continuous maximal regularity and analytic semigroups. *Discrete Contin. Dyn. Syst.*, (Dynamical systems, differential equations and applications. 8th AIMS Conference. Suppl. Vol. II):963–970, 2011.
- [11] Jeremy LeCrone and Gieri Simonett. On well-posedness, stability, and bifurcation for the axisymmetric surface diffusion flow. *SIAM J. Math. Anal.*, 45(5):2834–2869, 2013.
- [12] Jeremy LeCrone and Gieri Simonett. On the flow of non-axisymmetric perturbations of cylinders via surface diffusion. *J. Differential Equations*, 260(6):5510–5531, 2016.
- [13] Alessandra Lunardi. *Analytic semigroups and optimal regularity in parabolic problems*. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1995. [2013 reprint of the 1995 original] [MR1329547].
- [14] James McCoy, Glen Wheeler, and Graham Williams. Lifespan theorem for constrained surface diffusion flows. *Math. Z.*, 269(1-2):147–178, 2011.
- [15] Jan Prüss and Gieri Simonett. On the manifold of closed hypersurfaces in  $\mathbb{R}^n$ . *Discrete Contin. Dyn. Syst.*, 33(11-12):5407–5428, 2013.
- [16] Jan Prüss and Gieri Simonett. *Moving interfaces and quasilinear parabolic evolution equations*, volume 105 of *Monographs in Mathematics*. Birkhäuser/Springer, [Cham], 2016.
- [17] Jan Prüss, Gieri Simonett, and Mathias Wilke. Critical spaces for quasilinear parabolic evolution equations and applications. *J. Differential Equations*, 264(3):2028–2074, 2018.
- [18] Jan Prüss, Gieri Simonett, and Rico Zacher. On convergence of solutions to equilibria for quasilinear parabolic problems. *J. Differential Equations*, 246(10):3902–3931, 2009.
- [19] Jan Prüss, Gieri Simonett, and Rico Zacher. On normal stability for nonlinear parabolic equations. *Discrete Contin. Dyn. Syst.*, (Dynamical systems, differential equations and applications. 7th AIMS Conference, suppl.):612–621, 2009.
- [20] Jan Prüss and Mathias Wilke. Addendum to the paper “On quasilinear parabolic evolution equations in weighted  $L_p$ -spaces II” [MR3250797]. *J. Evol. Equ.*, 17(4):1381–1388, 2017.
- [21] Yuanzhen Shao. A family of parameter-dependent diffeomorphisms acting on function spaces over a Riemannian manifold and applications to geometric flows. *NoDEA Nonlinear Differential Equations Appl.*, 22(1):45–85, 2015.
- [22] Yuanzhen Shao and Gieri Simonett. Continuous maximal regularity on uniformly regular Riemannian manifolds. *J. Evol. Equ.*, 14(1):211–248, 2014.
- [23] Glen Wheeler. Lifespan theorem for simple constrained surface diffusion flows. *J. Math. Anal. Appl.*, 375(2):685–698, 2011.

- [24] Glen Wheeler. Surface diffusion flow near spheres. *Calc. Var. Partial Differential Equations*, 44(1-2):131–151, 2012.

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, UNIVERSITY OF RICHMOND, RICHMOND,  
VA USA

*Email address:* `jlecrone@richmond.edu`

*URL:* `http://math.richmond.edu/faculty/jlecrone/`

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN USA

*Email address:* `gieri.simonett@vanderbilt.edu`

*URL:* `http://www.vanderbilt.edu/math/people/simonett`